

294	HNGDQ38	808	<p>Production of MIP1alpha</p>	<p>using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-</p>	<p>dysplasia.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary</p>
-----	---------	-----	--------------------------------	--	-------------------	--

295	HNGDX18	809	<p>2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>Assays for the regulation of transcription through the DMEF1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the DMEF1 response element in a reporter construct (such as that containing the GLUT4 promoter) and to regulate insulin production. The DMEF1 response element is present in the GLUT4 promoter and binds to MEF2 transcription factor and another transcription factor that is required for insulin regulation of Glut4 expression in skeletal muscle. GLUT4 is the primary insulin-responsive glucose transporter in fat and muscle tissue. Exemplary assays that may be used or routinely modified to test for DMEF1 response element activity (in adipocytes and pre-adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Thai, M.V., et al., J Biol Chem,</p>	<p>cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
			<p>A highly preferred indication is diabetes mellitus.</p> <p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include</p>	



295	HNGDX18	809	Activation of transcription through GAS response element in immune cells (such as T-cells).	<p>273(23):14285-92 (1998); Mora, S., et al., J Biol Chem, 275(21):16323-8 (2000); Liu, M.L., et al., J Biol Chem, 269(45):28514-21 (1994); "Identification of a 30-base pair regulatory element and novel DNA binding protein that regulates the human GLUT4 promoter in transgenic mice", J Biol Chem. 2000 Aug 4;275(31):23666-73; Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Adipocytes and pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the mouse 3T3-L1 cell line which is an adherent mouse preadipocyte cell line. Mouse 3T3-L1 cells are a continuous substrain of 3T3 fibroblasts developed through clonal isolation. These cells undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation culture conditions.</p> <p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for</p>	<p>weight loss or alternatively, weight gain. highly preferred indications are complications associated with insulin resistance.</p>	Additional
				<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>		

295	HNGDX18	809	<p>transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
			<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated</p>

296	HNGDY34	810	Production of IL-6	<p>activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or</p>

			<p>Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell</p>	<p>"Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	---	---

297	HNGEA34	811	Production of IL-5	<p>proliferation and functional activities.</p> <p>IL-5 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein</p>	<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma,</p>
-----	---------	-----	--------------------	---	---

298	HNGEQ75	812	Protection from Endothelial Cell Apoptosis.	<p>incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Caspase Apoptosis Rescue. Assays for caspase apoptosis rescue are well known in the art and may be used or routinely modified to assess the ability of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to inhibit caspase protease-mediated apoptosis. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis rescue of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Romeo et al., Cardiovasc Res 45(3): 788-794 (2000); Messmer et al., Br J Pharmacol 127(7): 1633-1640 (1999); and J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine</p>	<p>arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method</p>	

				<p>aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders</p>
--	--	--	--	---	--

299	HNGGA68	813	<p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary</p>	<p>Activation of transcription through cAMP response element in immune cells (such as T-cells).</p>	<p>such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vascularitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
			<p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary</p>		<p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune</p>



299	HNGGA68	813	<p>assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
			<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications</p>

299	HNGGA68	813	<p>agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types.</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies</p>

300	HNGGP65	814	Activation of Hepatocyte ERK Signaling Pathway	<p>Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>(e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating hepatocyte cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting hepatocyte cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating hepatocyte cell differentiation. An alternative highly preferred embodiment of the invention includes a</p>	

				<p>promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Rat liver hepatoma cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary rat liver hepatoma cells that may be used according to these assays include H4IIE cells, which are known to respond to glucocorticoids, insulin, or cAMP derivatives.</p>	<p>method for inhibiting hepatocyte cell differentiation. A highly preferred embodiment of the invention includes a method for activating hepatocyte cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating hepatocyte cells. Highly preferred indications include disorders of the liver and/or endocrine disorders (e.g., as described below under "Endocrine Disorders"). Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section</p>
--	--	--	--	--	---

301	HNGHZ69	815	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and</p>	<p>below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hepatitis, jaundice, gallstones, cirrhosis of the liver, degenerative or necrotic liver disease, alcoholic liver diseases, fibrosis, liver regeneration, metabolic disease, dyslipidemia and cholesterol metabolism. Additional highly preferred indications include neoplasms and cancers, such as, hepatocarcinomas, other liver cancers, and colon and pancreatic cancer. Preferred indications also include prostate, breast, lung, esophageal, stomach, brain, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred</p>
-----	---------	-----	--	--	--

				<p>agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia,</p>
--	--	--	--	--	---

					<p>             bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and           </p>
--	--	--	--	--	---

302	HNGIV64	816	Activation of Adipocyte ERK Signaling Pathway	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation</p>	<p>inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders").</p> <p>Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic</p>
-----	---------	-----	---	--	--



				<p>and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	--	--	--

303	HNGJB41	817	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
304	HNGKT41	818	Activation of transcription through API response element in immune cells (such as T-cells).	Assays for the activation of transcription through the API response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm,	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as,

304	HNGKT41	818	<p>Activation of transcription through CD28 response element in immune cells (such as T-cells).</p>	<p>Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4 responsive suspension-culture cell line.</p>	<p>leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
			<p>Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g.,</p>	

304	HNGKT41	818	<p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of</p>	<p>166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysplastic disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication includes infection (e.g., AIDS, tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
					<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below),</p>

304	HNGKT41	818	<p>the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
			<p>Assays for the activation of transcription through the NFkB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFkB</p>	<p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as</p>

304	HNGKT41	818	<p>transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFkB response element that may be used or routinely modified to test NFkB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
			<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated</p>

				<p>modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
305	HNGMW45	819	Regulation of transcription through the PEPCK promoter in hepatocytes	<p>Assays for the regulation of transcription through the PEPCK promoter are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the PEPCK promoter in a reporter construct and regulate liver gluconeogenesis. Exemplary assays for regulation of transcription</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion,</p>

				<p>through the PEPCK promoter that may be used or routinely modified to test for PEPCK promoter activity (in hepatocytes) of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Lochhead et al., Diabetes 49(6):896-903 (2000); and Yeagley et al., J Biol Chem 275(23):17814-17820 (2000), the contents of each of which is herein incorporated by reference in its entirety. Hepatocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary liver hepatoma cells that may be used according to these assays include H4Ile cells, which contain a tyrosine amino transferase that is inducible with glucocorticoids, insulin, or cAMP derivatives.</p>	<p>drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., an infectious disease or disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include glycogen storage disease (e.g., glycogenoses), hepatitis, gallstones, cirrhosis of the liver, degenerative or necrotic liver disease, alcoholic liver diseases, fibrosis, liver regeneration, metabolic disease, dyslipidemia and cholesterol metabolism, and hepatocarcinomas. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), infection (e.g., an infectious disease and/or disorder as described below under "Infectious Disease"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), and neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"). Additional preferred indications include neoplastic diseases (e.g., as</p>
--	--	--	--	---	--



306	HNGNK44	820	Production of TNF alpha by dendritic cells	<p>TNF<math>\alpha</math> FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNF<math>\alpha</math>), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890</p>	<p>described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, and urinary cancer. A highly preferred indication is liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF<math>\alpha</math> production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF<math>\alpha</math> production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia,</p>
-----	---------	-----	---	--	--

306	HNGNK44	820		<p>(1198); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			<p>Upregulation of CD152 and activation of T cells</p>	<p>CD152 FMAT. CD152 (a.k.a. CTLA-4) expression is restricted to activated T cells. CD152 is a negative regulator of T cell proliferation. Reduced CD152 expression has been linked to hyperproliferative and autoimmune diseases. Overexpression of CD152 may lead to impaired immunoresponses. Assays for immunomodulatory proteins important in the maintenance of T cell homeostasis and expressed almost exclusively on CD4+ and CD8+ T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, maintain T cell homeostasis, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such</p>	<p>A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma,</p>

307	HNGNO53	821	<p>as CD152, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); McCoy et al., Immunol Cell Biol 77(1):1-10 (1999); Oosterveg et al., Curr Opin Immunol 11(3):294-300 (1999); and Saito T, Curr Opin Immunol 10(3):313-321 (1998), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy. An additional preferred indication is infection (e.g., as described below under "Infectious Disease").</p>
			<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple</p>

308	HNGP/25	822	<p>related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon,</p>

				<p>the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
308	HNGP125	822	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications</p>

				<p>the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
308	HNGP25	822	<p>Regulation of transcription of Malic Enzyme in adipocytes</p>	<p>Assays for the regulation of transcription of Malic Enzyme are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as</p>

				agonists or antagonists of the invention) to regulate transcription of Malic Enzyme, a key enzyme in lipogenesis. Malic enzyme is involved in lipogenesis and its expression is stimulated by insulin. ME promoter contains two direct repeat (DR1)-like elements MEp and MEd identified as putative PPAR response elements. ME promoter may also responds to AP1 and other transcription factors. Exemplary assays that may be used or routinely modified to test for regulation of transcription of Malic Enzyme (in adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Streeper, R.S., et al., Mol Endocrinol, 12(11):1778-91 (1998); Garcia-Jimenez, C., et al., Mol Endocrinol, 8(10):1361-9 (1994); Barroso, I., et al., J Biol Chem, 274(25):17997-8004 (1999); Ijpenberg, A., et al., J Biol Chem, 272(32):20108-20117 (1997); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays includes the H4IIE rat liver hepatoma cell line.	described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.
309	HNHEN82	823	Activation of Adipocyte PI3 Kinase Signalling Pathway	A highly preferred embodiment of the invention includes a method for increasing adipocyte survival. An alternative highly preferred embodiment of the invention	

				<p>metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>includes a method for decreasing adipocyte survival. A preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Preferred indications include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"), blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section</p>
--	--	--	--	--	--



310	HNHFE71	824	Activation of Adipocyte PI3 Kinase Signalling Pathway	<p>Kinase assay. Kinase assays, for example an GSK-3 assays, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of</p>	<p>below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternately, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Highly preferred indications include neoplasms and cancer, such as, lipoma, liposarcoma, lymphoma, leukemia and breast, colon, and kidney cancer. Additional highly preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred embodiment of the invention includes a method for increasing adipocyte survival. An alternative highly preferred embodiment of the invention includes a method for decreasing adipocyte survival. A preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A</p>
-----	---------	-----	---	--	---

				<p>the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Preferred indications include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"), blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g. due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the</p>
--	--	--	--	---	--

310	HNHFE71	824	Calcium flux in immune cells (such as monocytes)	Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be	<p>"Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Highly preferred indications include neoplasms and cancer, such as, lipoma, liposarcoma, lymphoma, leukemia and breast, colon, and kidney cancer. Additional highly preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Infection, Inflammation, Atherosclerosis, Hypersensitivity, and Leukemias</p>
-----	---------	-----	--	--	--

311	HNHGK22	825		<p>used or routinely modified to measure calcium flux in immune cells (such as monocytes) include assays disclosed in: Chan, CC, et al., J Pharmacol Exp Ther, 269(3):891-896 (1994); Andersson, K, et al., Cytokine, 12(12):1784-1787 (2000); Scully, SP, et al., J Clin Invest, 74(2) 589-599 (1984); and, Sullivan, E, et al., Methods Mol Biol, 114:125-133 (1999), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the THP-1 monocyte cell line.</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al.,</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally,</p>
-----	---------	-----	--	--	--

312	HNHHB10	826	Activation of transcription through GAS response element in immune cells (such as T-cells).	<p>Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>	

				<p>assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOL T4 cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
313	HNHKS19	827	Protection from Endothelial Cell Apoptosis.	<p>Caspase Apoptosis Rescue. Assays for caspase apoptosis rescue are well known in the art and may be used or routinely modified to assess the ability of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to inhibit caspase protease-mediated apoptosis. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis rescue of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Romeo et al., Cardiovasc Res 45(3): 788-794 (2000); Messmer et al., Br J Pharmacol 127(7): 1633-1640 (1999); and J Atheroscler Thromb 3(2): 75-80 (1996); the contents of</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred</p>

			<p>each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign</p>
--	--	--	---	---

					<p>dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
314	HNTBT17	828	Production of IL-10 and downregulation of immune responses	IL-10 FMAAT. Assays for immunomodulatory proteins produced by activated T cells, B cells, and monocytes	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IL-10. An alternative preferred embodiment of the invention</p>



				<p>that exhibit anti-inflammatory activity and downregulate monocyte/macrophage function and expression of cytokines are well known in the art and may be used or routinely modified to assess the ability of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, and modulate immune cell function and cytokine production. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-10, and the downmodulation of immune responses. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to</p>	<p>includes a method for inhibiting the production of IL-10. Highly preferred indications include inflammation and inflammatory disorders (e.g. inflammatory bowel disease). An additional highly preferred indication includes inflammatory bowel disease. Additional highly preferred indications include blood disorders (e.g., as described below under "Immune Activity" (e.g. autoimmune disorders), "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below). Preferred deficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, Crohn's disease, arthritis, AIDS, granulomatous disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy. An additional preferred indication is infection (e.g., as described below under "Infectious Disease").</p>
--	--	--	--	---	--

315	HNTMH79	829	<p>Activation of transcription through AP1 response element in immune cells (such as T-cells).</p>	<p>enhance responsiveness to immunomodulatory factors.</p> <p>Assays for the activation of transcription through the AP1 response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely modified to test AP1-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the HT2 cell line, which is an IL-2 dependent suspension culture cell line that also responds to IL-4.</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
316	HOABP31	830	<p>Production of IFNgamma using a T cells</p>	<p>IFNgamma FMAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the</p>

				<p>cytokine. IFN<math>\gamma</math> promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFN<math>\gamma</math>), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated</p>	<p>invention includes a method for inhibiting the production of IFN<math>\gamma</math>. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	--	---

317	HOABP31	831		<p>by reference in its entirety. Human T cells that may be used according to these assays herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>IFNgamma FMAT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNγ promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNγ), and the activation of T cells. Such assays that may be used or routinely modified to test</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach,</p>
-----	---------	-----	--	---	--

318	HOACG07	832		<p>immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
				<p>Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease,</p>

319	HODAG07	833	<p>activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 ( Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious</p>
			<p>Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including</p>	
			<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>	

				antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.	disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
320	HODBB70	834	Activation of transcription through cAMP response element in immune cells (such as T-cells).	Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary	Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune

			assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.	response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
320	HODBB70	834	Assays for the regulation of transcription through the PEPCK promoter are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the PEPCK promoter in a reporter construct and regulate liver gluconeogenesis. Exemplary assays for regulation of transcription through the PEPCK promoter that may be used or routinely modified to test for PEPCK promoter activity (in hepatocytes) of polypeptides of the invention (including	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the



				<p>antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Lochhead et al., Diabetes 49(6):896-903 (2000); and Yeagley et al., J Biol Chem 275(23):17814-17820 (2000), the contents of each of which is herein incorporated by reference in its entirety. Hepatocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary liver hepatoma cells that may be used according to these assays include H4Ile cells, which contain a tyrosine amino transferase that is inducible with glucocorticoids, insulin, or cAMP derivatives.</p>	<p>"Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., an infectious diseases or disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include glycogen storage disease (e.g., glycogenoses), hepatitis, gallstones, cirrhosis of the liver, degenerative or necrotic liver disease, alcoholic liver diseases, fibrosis, liver regeneration, metabolic disease, dyslipidemia and cholesterol metabolism, and hepatocarcinomas. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), infection (e.g., an infectious disease and/or disorder as described below under "Infectious Disease"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), and neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"). Additional preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, and urinary cancer.</p>
--	--	--	--	--	---

321	HODB V05	835	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used</p>	<p>A highly preferred indication is liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions,</p>
-----	----------	-----	---------------------	--	---

322	HODCZ32	836	Activation of Natural Killer Cell ERK Signaling Pathway.	<p>according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary natural killer cells that may be used according to these assays include the human natural killer cell lines (for</p>	<p>such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating natural killer cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating natural killer cell differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell differentiation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity") and infections (e.g., as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include cancers such as, kidney, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary cancer, lymphoma and leukemias.</p>	

323	HOEBK60	837	Insulin Secretion	<p>example, NK-YT cells which have cytolytic and cytotoxic activity) or primary NK cells.</p> <p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Shimizu, H., et al., Endocr J, 47(3):261-9 (2000); Salapatek, A.M., et al., Mol Endocrinol, 13(8):1305-17 (1999); Filipsson, K., et al., Ann N Y Acad Sci, 865:441-4 (1998); Olson, L.K., et al., J Biol Chem, 271(28):16544-52 (1996); and, Miraglia S et al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents</p>	<p>Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Other highly preferred indications include, pancytopenia, leukopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), arthritis, asthma, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, immune reactions to transplanted organs and tissues, endocarditis, meningitis, Lyme Disease, and allergies.</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated</p>
-----	---------	-----	-------------------	--	--

			<p>of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATCC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	with insulin resistance.	
324	HOFAA78	838	<p>Activation of Skeletal Muscle Cell PI3 Kinase Signalling Pathway</p>	<p>Kinase assay. Kinase assays, for example an GSK-3 kinase assay, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes</p>	<p>A highly preferred embodiment of the invention includes a method for increasing muscle cell survival. An alternative highly preferred embodiment of the invention includes a method for decreasing muscle cell survival. A preferred embodiment of the invention includes a method for stimulating muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is inhibited. A preferred embodiment of the invention includes a method for stimulating muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is inhibited. Highly preferred indications include disorders</p>

				<p>49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Rat myoblast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary rat myoblast cells that may be used according to these assays include L6 cells. L6 is an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuses to form multinucleated myotubes and striated fibers after culture in differentiation media.</p>	<p>of the musculoskeletal system. Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infections (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly</p>
--	--	--	--	--	---

325	HOFNB74	839	Insulin Secretion	<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Shimizu, H., et al., Endocr J, 47(3):261-9 (2000); Salapatek, A.M., et al., Mol</p>	<p>preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal system including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include: myopathy, atrophy, congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, and vascular disease. Highly preferred indications include neoplasms and cancer, such as, rhabdomyoma, rhabdosarcoma, stomach, esophageal, prostate, and urinary cancer. Preferred indications also include breast, lung, colon, pancreatic, brain, and liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred indication is diabetes mellitus.</p> <p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and</p>
-----	---------	-----	-------------------	--	--

326	HOFNU55	840	Activation of transcription through cAMP response element in immune cells (such as T-cells).	<p>Endocrinol, 13(8):1305-17 (1999); Filipsson, K., et al., Ann N Y Acad Sci, 865:441-4 (1998); Olson, L.K., et al., J Biol Chem, 271(28):16544-52 (1996); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATCC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP, bind to CREB transcription factor, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or</p>	<p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly</p>



			<p>routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
327	HGBF01	841	<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as,</p>

327	HOCBF01	841	Production of IL-6	<p>Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
				<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include</p>

327	HOGBF01	841	Production of IFNgamma using a T cells	<p>Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>IFNgamma FMAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNg promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2</p>	<p>neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNg. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease").</p>	

				<p>helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFN<math>\gamma</math>), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or</p>	<p>Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	--	---

328	HORBS82	842	<p>Activation of Adipocyte ERK Signaling Pathway</p>	<p>cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders").</p> <p>Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity" and "Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic</p>
-----	---------	-----	--	---	--

				<p>and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin)). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	--	--	---

328	HORBS82	842	<p>Regulation of apoptosis of immune cells (such as mast cells).</p>	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate caspase protease-mediated apoptosis in immune cells (such as, for example, in mast cells). Mast cells are found in connective and mucosal tissues throughout the body, and their activation via immunoglobulin E-antigen, promoted by T helper cell type 2 cytokines, is an important component of allergic disease. Dysregulation of mast cell apoptosis may play a role in allergic disease and mast cell tumor survival. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity induced by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Masuda A, et al., J Biol Chem, 276(28):26107-26113 (2001); Yeatman CF 2nd, et al., J Exp Med, 192(8):1093-1103 (2000); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary immune cells that may be used according to these assays include mast cells such as the HMC</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of asthma, allergy, hypersensitivity and inflammation.</p>
-----	---------	-----	--	---	--

329	HORBV76	843	Production of ICAM-1	human mast cell line. Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al., FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
330	HOSDO75	844	Regulation of apoptosis in pancreatic beta cells.	Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Apoptosis in pancreatic beta is associated with induction and progression of diabetes. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia,



				<p>assays disclosed in: Loweth, AC, et al., FEBS Lett, 400(3):285-8 (1997); Saini, KS, et al., Biochem Mol Biol Int, 39(6):1229-36 (1996); Krauthelm, A., et al., Br J Pharmacol, 129(4):687-94 (2000); Chandra J, et al., Diabetes, 50 Suppl 1:S44-7 (2001); Suk K, et al., J Immunol, 166(7):4481-9 (2001); Tejedo J, et al., FEBS Lett, 459(2):238-43 (1999); Zhang, S., et al., FEBS Lett, 455(3):315-20 (1999); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include RIN-m. RIN-m is a rat adherent pancreatic beta cell insulinoma cell line derived from a radiation induced transplantable rat islet cell tumor. The cells produce and secrete islet polypeptide hormones, and produce insulin, somatostatin, and possibly glucagon. ATTC: #CRL-2057 Chick et al. Proc. Natl. Acad. Sci. 1977 74:628; AF et al. Proc. Natl. Acad. Sci. 1980 77:3519.</p>	<p>endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
331	HOSEC25	845	<p>Production of IL-2 and activation of T cells</p>	<p>IL-2 FMAT. IL-2 is the principal T cell factor that allows T cell expansion and differentiation into effector cells. Assays for immunomodulatory proteins secreted by TH1 cells that promote T cell and NK</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. A highly preferred embodiment of the</p>

			<p>cell growth and differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, promote immune cell growth and differentiation, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-2, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Laduda et al., Immunology 94(4):496-502 (1998); and Powell et al., Immunol Rev 165:287-300 (1998), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>invention includes a method for stimulating T cell expansion. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell expansion. A highly preferred embodiment of the invention includes a method for stimulating T cell differentiation. In a specific embodiment, this method stimulates T cell differentiation into effector cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell differentiation. In a specific embodiment, this method inhibits the differentiation of T cells into effector cells. Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, ovarian, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). A highly preferred indication is AIDS and HIV infection. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), organ and tissue transplant rejection. Additional</p>
--	--	--	--	--

					preferred indications include inflammation and inflammatory disorders. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, Non-Hodgkin's lymphoma, Kaposi's sarcoma arthritis, granulomatous disease, inflammatory bowel disease, Hepatitis (e.g. Hepatitis C), sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
332	HOSE181	846	Proliferation, differentiation, and/or cytokine production in immune cells (such as T-cells).	Kinase assays, for example kinase assays for members of the MAP kinase family (including p38, JAK, and ERK) are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, differentiation., and/or cytokine production in immune cells such as T-cells. Exemplary assays for MAP kinase family members that may be used or routinely modified to test polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Rincon M., Curr Opin Immunol; 13(3):339-345 (2001); Wang LH, et al., J Immunol, 162(7):3897-3904 (1999); Sakamoto H, et al., J Biol Chem, 275(46):35857-35862 (2000), the contents of each of which are herein incorporated by reference in its entirety. Exemplary immune cells (for example, T-cells) that may be used according to these assays include the mouse CTLL cell line.	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of inflammation, infection, allergy, asthma, autoimmunity, and cancer.

333	HOSEJ94	847	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to</p>
-----	---------	-----	--------------------	---	---

334	HOUCA21	848	<p>each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>IFNgamma FMAAT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNγ promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNγ), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of</p>	<p>transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications</p>	

			<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neurophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
335	HOUE92	849	<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia,</p>

335	HOUDE2	849	<p>antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
			<p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm,</p>	<p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's</p>

335	HOUE92			<p>Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowsky et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
		849	<p>Activation of transcription through NFAT response in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al.,</p>	<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>



335	HOUDE92	849	Activation of transcription through NFKB response element in immune cells (such as B-cells).	<p>Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p> <p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Gri G, et al., Biol Chem, 273(11):6431-6438 (1998); Pyatt DW, et al., Cell Biol Toxicol 2000;16(1):41-51 (2000); Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA</p>	<p>dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
				<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Autoimmunity, Allergy and Asthma</p>	

335	HOUDE92	849	<p>Activation of transcription through NFKB response element in neuronal cells (such as SKNMC cells).</p>	<p>85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997); Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary immune cells that may be used according to these assays include the Reh B-cell line.</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Neurological Diseases and Disorders (e.g. Alzheimer's Disease, Parkinson's Disease, Brain Cancer, Seizures).</p>
-----	---------	-----	---	---	---

335	HOUDE92	849	<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>	<p>810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Neuronal cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary neuronal cells that may be used according to these assays include the SKNMC neuronal cell line.</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis,</p>
-----	---------	-----	--	--	---

335	HOUE92	849	<p>Activation of transcription through CD28 response element in immune cells (such as T-cells).</p>	<p>cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4 responsive suspension-culture cell line.</p> <p>Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>meningitis, and Lyme Disease.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication includes</p>
-----	--------	-----	---	--	---

335	HOUE92	849	<p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm,</p>	<p>infection (e.g., AIDS, tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
			<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary</p>	

			<p>Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
335	HOUDE92	849	<p>Activation of transcription through STAT6 response element in immune cells (such as T-cells).</p>	<p>A highly preferred indication is allergy.</p> <p>Another highly preferred indication is asthma.</p> <p>Additional highly preferred indications include inflammation and inflammatory disorders.</p> <p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below).</p> <p>Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>

335	HOUDE92	849	Activation of transcription through NFKB response element in immune cells (such as T-cells).	<p>368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Georas et al., Blood 92(12):4529-4538 (1998); Moffatt et al., Transplantation 69(7):1521-1523 (2000); Curiel et al., Eur J Immunol 27(8):1982-1987 (1997); and Masuda et al., J Biol Chem 275(38):29331-29337 (2000), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p> <p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et</p>	<p>conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred</p>	

335	HOUDE92	849	Activation of transcription through serum response element in immune cells (such as natural killer cells).	<p>al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types.</p> <p>Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are</p>	<p>indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
				<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>	



336	HOUDR07	850	Activation of Skeletal Muscle Cell ERK Signalling Pathway	publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.	hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
				Kinase assay. Kinase assays, for example Elk-1 kinase assays, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are	Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders") and disorders of the musculoskeletal system. Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease,

			<p>herein incorporated by reference in its entirety. Rat myoblast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary rat myoblast cells that may be used according to these assays include L6 cells. L6 is an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuses to form multinucleated myotubes and striated fibers after culture in differentiation media.</p>	<p>hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include: myopathy, atrophy, congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, and vascular disease. Highly preferred indications include neoplasms and cancer, such as, rhabdomyoma, rhabdosarcoma, stomach, esophageal, prostate, and urinary cancer. Highly preferred indications also include breast, lung, colon, pancreatic, brain, and liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, hyperplasia, metaplasia, and/or dysplasia.</p>
337	HOUED72	851	<p>Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases</p>

338	HOUFS04	852	<p>other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p> <p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response</p>	<p>(e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p> <p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune</p>
-----	---------	-----	--	--

				<p>element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOLT4 cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
338	HOUFS04	852	<p>Activation of transcription through NFKB response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides</p>	<p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described</p>

				<p>of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOL T4, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
338	HOUFS04	852	<p>Upregulation of CD152 and activation of T cells</p>	<p>CD152 FMAT. CD152 (a.k.a. CTLA-4) expression is restricted to activated T cells. CD152 is a negative regulator of T cell proliferation. Reduced CD152 expression has been linked to hyperproliferative and autoimmune diseases. Overexpression of CD152 may lead to impaired immunoresponses. Assays for immunomodulatory proteins important in the maintenance of T cell homeostasis and expressed almost exclusively on CD4+ and CD8+ T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, maintain T cell homeostasis, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for</p>	<p>A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally,</p>

			immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD152, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); McCoy et al., Immunol Cell Biol 77(1):1-10 (1999); Oostervegal et al., Curr Opin Immunol 11(3):294-300 (1999); and Saito T, Curr Opin Immunol 10(3):313-321 (1998), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy. An additional preferred indication is infection (e.g., as described below under "Infectious Disease").
339	HOUH125	853	Activation of transcription through STAT6 response element in immune cells (such as T-cells).	<p>A highly preferred indication is allergy.</p> <p>Another highly preferred indication is asthma.</p> <p>Additional highly preferred indications include inflammation and inflammatory disorders.</p> <p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Preferred indications include autoimmune diseases (e.g.,</p>

340	HOVBD85	854	<p>regulate STAT6 transcription factors and modulate the expression of multiple genes. Exemplary assays for transcription through the STAT6 response element that may be used or routinely modified to test STAT6 response element activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Georas et al., Blood 92(12):4529-4538 (1998); Moffatt et al., Transplantation 69(7):1521-1523 (2000); Curiel et al., Eur J Immunol 27(8):1982-1987 (1997); and Masuda et al., J Biol Chem 275(38):29331-29337 (2000), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple</p>

				<p>related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
341	HPCAB41	855	<p>Activation of Endothelial Cell p38 or JNK Signaling Pathway.</p>	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred</p>



				<p>the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization.</p>
--	--	--	--	---	---

					<p>Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vascularitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood</p>
--	--	--	--	--	---

342	HPCAL26	856	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that</p>	<p>disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications</p>
-----	---------	-----	--	---	---

				may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.	include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
343	HPEAD23	857	Stimulation of Calcium Flux in pancreatic beta cells.	Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 (Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.

344	HPFBA54	858	Activation of transcription through API response element in immune cells (such as T-cells).	<p>according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>
-----	---------	-----	---	---	--

344	HPFBA54	858	Production of IFNgamma using a T cells	<p>4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p> <p>IFNgamma FMAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNg promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test</p>	<p>dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNg. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach,</p>	

				immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
345	HPFCT36	859	Production of MIP1alpha	MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test	A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as

				<p>described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
			<p>for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
345	HPFC136	859	<p>Stimulation of Calcium Flux in pancreatic beta cells.</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to</p>



				<p>measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 ( Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--	--

345	HPFCI36	859	<p>Activation of Skeletal Muscle Cell PI3 Kinase Signalling Pathway</p>	<p>Kinase assay. Kinase assays, for example an GSK-3 kinase assay, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Rat myoblast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary rat myoblast cells that may be used according to these assays include L6 cells. L6 is an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuses to form multinucleated myotubes and striated fibers after culture in differentiation media.</p>	<p>A highly preferred embodiment of the invention includes a method for increasing muscle cell survival. An alternative highly preferred embodiment of the invention includes a method for decreasing muscle cell survival. A preferred embodiment of the invention includes a method for stimulating muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is inhibited. A preferred embodiment of the invention includes a method for stimulating muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is inhibited. Highly preferred indications include disorders of the musculoskeletal system. Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease,</p>
-----	---------	-----	---	---	--

346	HPFD137	860	Regulation of apoptosis in pancreatic beta cells.	Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to	stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infections (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal system including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include: myopathy, atrophy, congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, and vascular disease. Highly preferred indications include neoplasms and cancer, such as, rhabdomyoma, rhabdosarcoma, stomach, esophageal, prostate, and urinary cancer. Preferred indications also include breast, lung, colon, pancreatic, brain, and liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, hyperplasia, metaplasia, and/or dysplasia.
					A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g. diabetic retinopathy,

			<p>assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Apoptosis in pancreatic beta is associated with induction and progression of diabetes. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Loweth, AC, et al., FEBS Lett, 400(3):285-8 (1997); Saini, KS, et al., Biochem Mol Biol Int, 39(6):1229-36 (1996); Krauthaim, A., et al., Br J Pharmacol, 129(4):687-94 (2000); Chandra J, et al., Diabetes, 50 Suppl 1:S44-7 (2001); Suk K, et al., J Immunol, 166(7):4481-9 (2001); Tejedo J, et al., FEBS Lett, 459(2):238-43 (1999); Zhang, S., et al., FEBS Lett, 455(3):315-20 (1999); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include RIN-m. RIN-m is a rat adherent pancreatic beta cell insulinoma cell line derived from a radiation induced transplantable rat islet</p>	<p>diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	---	---

347	HPTAA80	861		<p>cell tumor. The cells produce and secrete islet polypeptide hormones, and produce insulin, somatostatin, and possibly glucagon. ATTC: #CRL-2057 Chick et al. Proc. Natl. Acad. Sci. 1977 74:628; AF et al. Proc. Natl. Acad. Sci. 1980 77:3519.</p> <p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication</p>
-----	---------	-----	--	--	--

				<p>include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin)). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and</p>
--	--	--	--	---	---

347	HP1AA80	861	Regulation of transcription of Malic Enzyme in adipocytes	<p>Assays for the regulation of transcription of Malic Enzyme are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate transcription of Malic Enzyme, a key enzyme in lipogenesis. Malic enzyme is involved in lipogenesis and its expression is stimulated by insulin. ME promoter contains two direct repeat (DR1)-like elements MEp and MEed identified as putative PPAR response elements. ME promoter may also respond to AP1 and other transcription factors. Exemplary assays that may be used or routinely modified to test for regulation of transcription of Malic Enzyme (in adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Streeter, R.S., et al., Mol Endocrinol, 12(11):1778-91 (1998); Garcia-Jimenez, C., et al., Mol Endocrinol, 8(10):1361-9 (1994); Barroso, I., et al., J Biol Chem, 274(25):17997-8004 (1999); Ijpenberg, A., et al., J Biol Chem, 272(32):20108-20117 (1997); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used</p>	<p>liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
-----	---------	-----	---	--	--

348	HPJBJ51	862	Production of IL-6	<p>according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays includes the H4IIE rat liver hepatoma cell line.</p> <p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention)</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
-----	---------	-----	--------------------	---	---



349	HPJB151	863	Production of IL-6	<p>include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function.</p>	<p>Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include</p>	

350	HPJBU43	864	Upregulation of CD152 and activation of T cells	<p>Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>CD152 FMAT. CD152 (a.k.a. CTLA-4) expression is restricted to activated T cells. CD152 is a negative regulator of T cell proliferation. Reduced CD152 expression has been linked to hyperproliferative and autoimmune diseases. Overexpression of CD152 may lead to impaired immunoresponses. Assays for immunomodulatory proteins important in the maintenance of T cell homeostasis and expressed almost exclusively on CD4+ and</p>	<p>neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or</p>	

				<p>CD8+ T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, maintain T cell homeostasis, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD152, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); McCoy et al., Immunol Cell Biol 77(1):1-10 (1999); Oostervegal et al., Curr Opin Immunol 11(3):294-300 (1999); and Saito T, Curr Opin Immunol 10(3):313-321 (1998), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to</p>	<p>"Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy. An additional preferred indication is infection (e.g., as described below under "Infectious Disease").</p>
--	--	--	--	---	--

350	HPJBU43	864	Upregulation of CD69 and activation of T cells	<p>immunomodulatory factors.</p> <p>CD69 FMAT. CD69 is an activation marker that is expressed on activated T cells, B cells, and NK cells. CD69 is not expressed on resting T cells, B cells, or NK cells. CD69 has been found to be associated with inflammation. Assays for immunomodulatory proteins expressed in T cells, B cells, and leukocytes are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD69, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ferenczi et al., J Autoimmun 14(1):63-78 (2000); Werfel et al., Allergy 52(4):465-469 (1997); Taylor-Fishwick and Siegel, Eur J Immunol 25(12):3215-3221 (1995); and Afetra et al., Ann Rheum Dis 52(6):457-460 (1993), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used</p>	<p>A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for activation B cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating B cells. A highly preferred embodiment of the invention includes a method for activating NK cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting activation of and/or inactivation NK cells. Highly preferred indications include inflammation and inflammatory disorders (e.g., as described below under "Immune Activity"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response and alternatively suppressing a T cell-mediated immune response, and boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. An additional highly preferred indication includes infection (e.g., as described below under "Infectious Disease"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia,</p>
-----	---------	-----	--	---	---

351	HPJCW58	865	<p>according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, asthma, and allergies. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
			<p>Assays for the regulation of transcription through the FAS promoter element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the FAS promoter element in a reporter construct and to regulate transcription of FAS, a key enzyme for lipogenesis. FAS promoter is regulated by many transcription factors including SREBP. Insulin increases FAS gene transcription in livers of diabetic mice. This stimulation of transcription is also somewhat glucose dependent. Exemplary assays that may be used or routinely modified to test for FAS promoter element activity (in hepatocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Xiong, S., et al., Proc Natl Acad Sci U.S.A., 97(8):3948-53 (2000); Roder, K., et al., Eur J Biochem, 260(3):743-51 (1999); Oskouian B, et al., Biochem J, 317</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include</p>

				<p>(Pt 1):257-65 (1996); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used according to these assays, such as H4IIE cells, are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays include rat liver hepatoma cell line(s) inducible with glucocorticoids, insulin, or cAMP derivatives.</p>	<p>weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
352	HPMBX22	866	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described</p>

				production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
353	HPMCJ84	867	Activation of transcription through CD28 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or	A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred

			<p>antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. An additional highly preferred indication includes infection (e.g., AIDS, and/or as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication is infection (e.g., tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease,</p>
--	--	--	--	---



354	HPMCV30	868	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.  Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
355	HPMFH77	869	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated

356	HPQAX38	870	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in</p>	<p>activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis,</p>
-----	---------	-----	--	---	---	---

357	HPQAX38	871	<p>growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
357	HPQAX38	871	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under</p>

			antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.	<p>"Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the</p>
358	HPQCB83	872	Activation of Endothelial Cell p38 or JNK Signaling	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are</p>

			<p>well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such</p>
--	--	--	---	---

					<p>as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment</p>
--	--	--	--	--	---

					<p>/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
359	HPQCC53	873	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes</p>

360	HPRBH85	874	<p>assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
			<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in:</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious</p>



				<p>Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p>	<p>diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
361	HPRCA64	875	Activation of Adipocyte ERK Signaling Pathway	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention)</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as</p>

				<p>include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss</p>
--	--	--	--	---	---

				<p>or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
362	HPRCD35	876	<p>Stimulation of Calcium Flux in pancreatic beta cells.</p>	<p>Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of</p>
				<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and</p>

363	HPTRM02	877	Regulation of viability and proliferation of pancreatic beta cells.	<p>the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 ( Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar</p>	

				<p>quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ohtani KI, et al., Endocrinology, 139(1):172-8 (1998); Krautheim A, et al, Exp Clin Endocrinol Diabetes, 107 (1):29-34 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
363	HPTRM02	877	Activation of transcription through GAS response element in immune cells (such as T-cells).		Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's

				<p>antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
364	HPWBA29	878	Activation of transcription through NFAT response in immune cells (such as T-cells).	<p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of</p>	<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below),</p>

				<p>the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
364	HPWBA29	878	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon,</p>	

			the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOLT4 cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).	pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
365	HPWDK06	879	CD152 FMAT. CD152 (a.k.a. CTLA-4) expression is restricted to activated T cells. CD152 is a negative regulator of T cell proliferation. Reduced CD152 expression has been linked to hyperproliferative and autoimmune diseases. Overexpression of CD152 may lead to impaired immunoresponses. Assays for	A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for stimulating T cell



				<p>immunomodulatory proteins important in the maintenance of T cell homeostasis and expressed almost exclusively on CD4+ and CD8+ T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, maintain T cell homeostasis, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD152, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); McCoy et al., Immunol Cell Biol 77(1):1-10 (1999); Oostervegal et al., Curr Opin Immunol 11(3):294-300 (1999); and Saito T, Curr Opin Immunol 10(3):313-321 (1998), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These</p>	<p>proliferation. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy. An additional preferred indication is infection (e.g., as described below under "Infectious Disease").</p>
--	--	--	--	--	---

366	HRAAD30	880	Activation of Adipocyte ERK Signaling Pathway	cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.  Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as</p>
-----	---------	-----	---	--	---

				<p>cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p> <p>described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin)). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>
--	--	--	--	--

367	HRADA42	881	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>dysplasia.</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis,</p>
-----	---------	-----	--	--	---

368	HRADF49	882	Activation of Adipocyte ERK Signaling Pathway	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast</p>	<p>meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as</p>
-----	---------	-----	---	---	---

				<p>cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin)). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>
--	--	--	--	---	--

368	HRADF49	882	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>dysplasia.</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis,</p>
-----	---------	-----	--	--	---

369	HRADN25	883	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia,</p>
-----	---------	-----	--	--	---



370	HRADT25	884	Production of IFN $\gamma$ using a T cells	<p>neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFN<math>\gamma</math>. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFN<math>\gamma</math>. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications</p>
				<p>IFN<math>\gamma</math> FMT. IFN<math>\gamma</math> plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFN<math>\gamma</math> promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFN<math>\gamma</math>), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed</p>

371	HRDA117	885	Production of ICAM-1	<p>in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly</p>	<p>include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p> <p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke</p>
-----	---------	-----	----------------------	--	--

372	HRDDQ39	886	Regulation of transcription via DMEF1 response element in adipocytes and pre-adipocytes	<p>available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).</p> <p>Assays for the regulation of transcription through the DMEF1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the DMEF1 response element in a reporter construct (such as that containing the GLUT4 promoter) and to regulate insulin production. The DMEF1 response element is present in the GLUT4 promoter and binds to MEF2 transcription factor and another transcription factor that is required for insulin regulation of Glut4 expression in skeletal muscle. GLUT4 is the primary insulin-responsive glucose transporter in fat and muscle tissue. Exemplary assays that may be used or routinely modified to test for DMEF1 response element activity (in adipocytes and pre-adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Thai, M.V., et al., J Biol Chem, 273(23):14285-92 (1998); Mora, S., et al., J Biol Chem, 275(21):16323-8 (2000); Liu, M.L., et al., J Biol Chem, 269(45):28514-21 (1994); "Identification of a 30-base pair regulatory element and novel DNA binding protein that regulates the human GLUT4 promoter in transgenic mice", J</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
-----	---------	-----	---	---	---

372	HRDDQ39	886	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Biol Chem. 2000 Aug 4;275(31):23666-73; Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Adipocytes and pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the mouse 3T3-L1 cell line which is an adherent mouse preadipocyte cell line. Mouse 3T3-L1 cells are a continuous substrain of 3T3 fibroblasts developed through clonal isolation. These cells undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation culture conditions.</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases</p>
-----	---------	-----	--	--	--

372	HRDDQ39	886	<p>Stimulation of insulin secretion from pancreatic beta cells.</p>	<p>Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>(e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia,</p>

			secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.	endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.
373	HRDER22	887	Activation of transcription through serum response element in immune cells (such as T-cells).	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies

			used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.	(e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
374	HRDEX93	888	Assays for the regulation of transcription through the FAS promoter element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the FAS promoter element in a reporter construct	<p>A highly preferred indication is diabetes mellitus.</p> <p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease,</p>

			<p>and to regulate transcription of FAS, a key enzyme for lipogenesis. FAS promoter is regulated by many transcription factors including SREBP. Insulin increases FAS gene transcription in livers of diabetic mice. This stimulation of transcription is also somewhat glucose dependent. Exemplary assays that may be used or routinely modified to test for FAS promoter element activity (in hepatocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Xiong, S., et al., Proc Natl Acad Sci U.S.A., 97(8):3948-53 (2000); Roder, K., et al., Eur J Biochem, 260(3):743-51 (1999); Oskouian B, et al., Biochem J, 317 (Pt 1):257-65 (1996); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used according to these assays, such as H4IIE cells, are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays include rat liver hepatoma cell line(s) inducible with glucocorticoids, insulin, or cAMP derivatives.</p>	<p>stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
375	HRDFK37	889	<p>Activation of transcription through serum response element in immune cells (such as natural killer cells).</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under</p>



			<p>antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types.</p> <p>Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>"Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
376	HRGBD54	890	<p>IFNgamma FMT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the</p>

			<p>cytokine. IFN<math>\gamma</math> promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFN<math>\gamma</math>), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated</p>	<p>invention includes a method for inhibiting the production of IFN<math>\gamma</math>. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	---

				by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	
377	HROEA08	891	Activation of Adipocyte ERK Signaling Pathway	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders").</p> <p>Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity</p>

				<p>be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p> <p>and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin)). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and</p>
--	--	--	--	--

378	HSAVA08	892	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells	breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include benign liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
				A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia,	

378	HSAVA08	892	Production of IL-5	with cytotoxic activity.	<p>thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred</p>
					<p>IL-5 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed</p>

			<p>in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
379	HSA/VW42	893	<p>Stimulation of Calcium Flux in pancreatic beta cells.</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia,</p>

380	HSAWN53	894	Production of ICAM-1	<p>routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995);Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 ( Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p> <p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely</p>	<p>endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Vascular Disease, Atherosclerosis, Restenosis, Stroke, and Asthma.</p>	



381	HSWZ40	895	Production of ICAM-1	<p>modified to measure ICAM-1 expression include assays disclosed in: Rolfe BE, et al., Atherosclerosis, 149(1):99-110 (2000); Panettieri RA Jr, et al., J Immunol, 154(5):2358-2365 (1995); and, Grunstein MM, et al., Am J Physiol Lung Cell Mol Physiol, 278(6):L1154-L1163 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include Aortic Smooth Muscle Cells (AOSMC); such as bovine AOSMC.</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke</p>
				<p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al., FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).</p>	

382	HSAYC41	896	<p>Stimulation of insulin secretion from pancreatic beta cells.</p>	<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAIT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
-----	---------	-----	---	--	---

383	HSDZM54	897	<p>characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types.</p> <p>Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia,</p>
-----	---------	-----	--	--

384	HSHBF76	898	Endothelial Cell Apoptosis	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of</p>	<p>neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic</p>
-----	---------	-----	----------------------------	--	--

				<p>endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly</p>
--	--	--	--	--	--

384	HSHBF76	898	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate</p>	<p>preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory</p>
-----	---------	-----	--------------------	---	---

			<p>immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
385	HSIFG47	899	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly</p>

				<p>of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma,</p>
--	--	--	--	--	--



					<p>angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic</p>
--	--	--	--	--	---

386	HSJBY32	900	Endothelial Cell Apoptosis	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels</p>
-----	---------	-----	----------------------------	--	---

					<p>themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization.</p> <p>Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions.</p>
--	--	--	--	--	---

387	HSKDR27	901	Endothelial Cell Apoptosis	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used</p>	<p>Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and</p>
-----	---------	-----	----------------------------	--	---

				<p>according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also</p>
--	--	--	--	--	--

					include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
387	HSKDR27	901	Proliferation, differentiation, and/or cytokine production in immune cells (such as T-cells).	Kinase assays, for example kinase assays for members of the MAP kinase family (including p38, JAK, and ERK) are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, differentiation, and/or cytokine production in immune cells such as T-cells. Exemplary assays for MAP kinase family members that may be used or routinely modified to test polypeptides	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of inflammation, infection, allergy, asthma, autoimmunity, and cancer.

387	HSKDR27	901	Production of IL-4	<p>of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Rincon M., Curr Opin Immunol; 13(3):339-345 (2001); Wang LH, et al., J Immunol, 162(7):3897-3904 (1999); Sakamoto H, et al., J Biol Chem, 275(46):35857-35862 (2000), the contents of each of which are herein incorporated by reference in its entirety. Exemplary immune cells (for example, T-cells) that may be used according to these assays include the mouse CTLL cell line.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-4 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-4 production. A highly preferred indication includes asthma. A highly preferred indication includes allergy. A highly preferred indication includes rhinitis. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic</p>
-----	---------	-----	--------------------	---	--

				<p>Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):277-283 (1994); Yssel et al., Res Immunol 144(8):610-616 (1993); Bagley et al., Nat Immunol 1(3):257-261 (2000); and van der Graaff et al., Rheumatology (Oxford) 38(3):214-220 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
388	HSLHG78	902	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-</p>



				<p>of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>		<p>mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
388	HSLHG78	902	Production of MCP-1	<p>MCP-1 FMA7. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as</p>	

				<p>modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An</p>
388	HSLHG78	902	Production of MIP1alpha	<p>MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by</p>		

				<p>activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture,</p>	<p>alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP 1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	--	---	---

389	HSLHX15	903	<p>Upregulation of HLA-DR and activation of T cells</p>	<p>which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>HLA-DR FMAT. MHC class II is essential for correct presentation of antigen to CD4+ T cells. Deregulation of MHC class II has been associated with autoimmune diseases (e.g., diabetes, rheumatoid arthritis, systemic lupus erythematosus, and multiple sclerosis). Assays for immunomodulatory proteins expressed on MHC class II expressing T cells and antigen presenting cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of MHC class II products, such as HLA-DR antigens, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Lamour et al., Clin Exp Immunol 89(2):217-222 (1992); Hurme and Sihvola, Immunol Lett 20(3):217-222 (1989); Gansbacher and Zier, Cell Immunol</p>	<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and alternatively, suppressing a T cell-mediated immune response. A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly</p>
-----	---------	-----	---	--	--

390	HSNAP85	904	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in</p> <p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>117(1):22-34 (1988); and Itoh et al., J Histochem Cytochem 40(11):1675-1683, the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. An additional preferred indication is infection (e.g., AIDS, and/or as described below under "Infectious Disease"). Preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"), and neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancer, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy.</p>
					<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis,</p>

				<p>growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
391	HSNAZ09	905	<p>Upregulation of CD154 and activation of T cells</p>	<p>CD154 FMAT, CD154 (a.k.a., CD40L) expression is induced following activation of T cells. Interraction between CD154 and CD40 on B cells is required for correct antibody class switching and germinal center formation. Mutations in CD154 are</p>	<p>A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. Highly preferred indications include blood disorders (e.g., as described below under "Immune</p>

				<p>linked to immunodeficiencies and increased susceptibility to infections. Assays for immunomodulatory proteins important for antibody class switching and TH1 function and expressed on activated T helper lymphocytes are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, modulate antibody class switching, mediate TH1 function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD154, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Mackey et al., J Leukoc Biol 63(4):418-428 (1998); and Skov et al., 164(7):3500-3505 (2000), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and</p>	<p>Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., AIDS). Preferred indications include boosting a T cell-mediated immune response, and alternatively, suppressing a T cell-mediated immune response. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy.</p>
--	--	--	--	--	--

392	HSNBM34	906	<p>express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Assays for the regulation of transcription through the DMEF1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the DMEF1 response element in a reporter construct (such as that containing the GLUT4 promoter) and to regulate insulin production. The DMEF1 response element is present in the GLUT4 promoter and binds to MEF2 transcription factor and another transcription factor that is required for insulin regulation of Glut4 expression in skeletal muscle. GLUT4 is the primary insulin-responsive glucose transporter in fat and muscle tissue. Exemplary assays that may be used or routinely modified to test for DMEF1 response element activity (in adipocytes and pre-adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Thai, M.V., et al., J Biol Chem, 273(23):14285-92 (1998); Mora, S., et al., J Biol Chem, 275(21):16323-8 (2000); Liu, M.L., et al., J Biol Chem, 269(45):28514-21 (1994); "Identification of a 30-base pair regulatory element and novel DNA binding protein that regulates the human GLUT4 promoter in transgenic mice", J</p>	<p>express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
-----	---------	-----	---	---	---



393	H50AH16	907	Production of IFN $\gamma$ using a T cells	<p>Biol Chem. 2000 Aug 4;275(31):23666-73; Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Adipocytes and pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the mouse 3T3-L1 cell line which is an adherent mouse preadipocyte cell line. Mouse 3T3-L1 cells are a continuous substrain of 3T3 fibroblasts developed through clonal isolation. These cells undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation culture conditions.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFN<math>\gamma</math>. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFN<math>\gamma</math>. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory</p>
-----	---------	-----	--	--	--

			<p>helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFN<math>\gamma</math>), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
394	HSQBF66	908	<p>IL-5 FMA7. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil</p>	<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g.,</p>

				<p>function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are</p>	<p>increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
--	--	--	--	--	--

395	HSQDO85	909	<p>primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the</p>
-----	---------	-----	---	---

				<p>cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p> <p>cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also</p>
--	--	--	--	---

					include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vascularitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
396	HSQES57	910	Production of TNF alpha by dendritic cells	TNFa FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary	A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and

			<p>assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1198); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
396	HSQES57	910	<p>Activation of Natural Killer Cell ERK Signaling Pathway.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating natural killer cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell proliferation. A highly preferred embodiment of the invention includes a method for</p>

			<p>invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary natural killer cells that may be used according to these assays include the human natural killer cell lines (for example, NK-YT cells which have cytolytic and cytotoxic activity) or primary NK cells.</p>	<p>stimulating natural killer cell differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell differentiation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity") and infections (e.g., as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include cancers such as, kidney, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary cancer, lymphoma and leukemias. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Other highly preferred indications include, pancytopenia, leukopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), arthritis, asthma, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, immune reactions to transplanted organs and tissues, endocarditis, meningitis, Lyme Disease, and allergies.</p>
397	HSRBE06	911	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g.,</p>



			as T-cells).	ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.	increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
398	HSSDI26	912	Production of IL-6	IL-6 FMAT. IL-6 is produced by T cells	A highly preferred embodiment of the invention

			<p>and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by</p>	<p>includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia,</p>
--	--	--	---	---

398	HSSDI26	912	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.  Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2):495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that	hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
399	HSSDI26	912	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.  Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2):495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation

				<p>may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema;</p>
--	--	--	--	--	---

399	HSSEA64	913	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be</p>	<p>and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vascularitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
					<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies</p>

400	HSSEF77	914	Proliferation, differentiation, and/or cytokine production in immune cells (such as T-cells).	used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.	(e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
					Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of inflammation, infection, allergy, asthma, autoimmunity, and cancer.

400	HSSEF77	914		<p>proliferation, differentiation., and/or cytokine production in immune cells such as T-cells. Exemplary assays for MAP kinase family members that may be used or routinely modified to test polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Rincon M., Curr Opin Immunol; 13(3):339-345 (2001); Wang LH, et al., J Immunol, 162(7):3897-3904 (1999); Sakamoto H, et al., J Biol Chem, 275(46):35857-35862 (2000), the contents of each of which are herein incorporated by reference in its entirety. Exemplary immune cells (for example, T-cells) that may be used according to these assays include the mouse CTLL cell line.</p> <p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in:</p>	<p>A highly preferred indication is diabetes mellitus.</p> <p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious</p>
-----	---------	-----	--	--	--

				<p>Shimizu, H., et al., Endocr J, 47(3):261-9 (2000); Salapatek, A.M., et al., Mol Endocrinol, 13(8):1305-17 (1999); Filipsson, K., et al., Ann N Y Acad Sci, 865:441-4 (1998); Olson, L.K., et al., J Biol Chem, 271(28):16544-52 (1996); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
401	HSSF38	915	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple</p>



401	HSSF38	915	Upregulation of CD152 and activation of T cells	transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.	sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
					<p>A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. An alternative highly preferred embodiment</p>

				<p>immunoresponses. Assays for immunomodulatory proteins important in the maintenance of T cell homeostasis and expressed almost exclusively on CD4+ and CD8+ T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, maintain T cell homeostasis, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD152, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); McCoy et al., Immunol Cell Biol 77(1):1-10 (1999); Oosterveeg et al., Curr Opin Immunol 11(3):294-300 (1999); and Saito T, Curr Opin Immunol 10(3):313-321 (1998), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell</p>	<p>of the invention includes a method for stimulating T cell proliferation. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy. An additional preferred indication is infection (e.g., as described below under "Infectious Disease").</p>
--	--	--	--	--	---

402	HSSGJ58	916	Production of ICAM-1	<p>receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke</p>
403	HSWBE76	917	Production of IL-5	<p>IL-5 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to</p>	<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly</p>

				<p>mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>preferred indication includes asthma. A highly preferred indication includes rhinitis. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
--	--	--	--	---	--

404	HSXCP38	918	<p>Activation of transcription through serum response element in immune cells (such as natural killer cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and</p>
-----	---------	-----	---	---	--

405	HSYBI06	919	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Rolfe BE, et al., Atherosclerosis, 149(1):99-110 (2000); Panettieri RA Jr, et al., J Immunol, 154(5):2358-2365 (1995); and, Grunstein MM, et al., Am J Physiol Lung Cell Mol Physiol, 278(6):L1154-L1163 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include Aortic Smooth Muscle Cells (AOSMC); such as bovine AOSMC.	asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Vascular Disease, Atherosclerosis, Restenosis, Stroke, and Asthma.
406	HT1SC27	920	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method

				<p>activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid</p>
--	--	--	--	---	---

					<p>tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include</p>
--	--	--	--	--	--



407	HT3BF49	921	Production of IL-4	<p>IL-4 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells that stimulate B cells, T cells, macrophages and mast cells and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cells, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-4, and the stimulation of immune cells, such as B cells, T cells, macrophages and mast cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):277-283 (1994); Yssel et al., Res Immunol 144(8):610-616 (1993); Bagley et al., Nat</p>	<p>autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-4 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-4 production. A highly preferred indication includes asthma. A highly preferred indication includes allergy. A highly preferred indication includes rhinitis. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS,</p>
-----	---------	-----	--------------------	---	---

407	HT3BF49	921	Production of IL-5	<p>Immunol 1(3):257-261 (2000); and van der Graaff et al., Rheumatology (Oxford) 38(3):214-220 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>IL-5 FMA T. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test</p>	<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and</p>

408	HT4FV41	922	Stimulation of insulin secretion from pancreatic beta cells.	<p>immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain</p>	<p>immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
				<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar</p>	

409	HT5FX79	923	Activation of Adipocyte ERK Signaling Pathway	<p>proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., <i>Am J Physiol</i>, 277(4 Pt 2):R959-66 (1999); Li, M., et al., <i>Endocrinology</i>, 138(9):3735-40 (1997); Kim, K.H., et al., <i>FEBS Lett</i>, 377(2):237-9 (1995); and, Miraglia S et. al., <i>Journal of Biomolecular Screening</i>, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Afari et al. <i>Endocrinology</i> 1992 130:167.</p>	<p>coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred</p>	

				agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.	embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine
--	--	--	--	--	--

410	HT5GR59	924	<p>Assays for the activation of transcription through the AP1 response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely</p>	<p>Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
			<p>Assays for the activation of transcription through the AP1 response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional</p>

				<p>modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
410	HT5GR59	924	Stimulation of insulin secretion from pancreatic beta cells.	<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia,</p>

411	HTAEI78	925	Upregulation of HLA-DR and activation of T cells	<p>secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p> <p>HLA-DR FMAT. MHC class II is essential for correct presentation of antigen to CD4+ T cells. Deregulation of MHC class II has been associated with autoimmune diseases (e.g., diabetes, rheumatoid arthritis, systemic lupus erythematosus, and multiple sclerosis). Assays for immunomodulatory proteins expressed on MHC class II expressing T cells and antigen presenting cells are well known in the art and may be used or routinely modified to assess the</p>	<p>endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and alternatively, suppressing a T cell-mediated immune response. A highly preferred indication is diabetes mellitus. An additional highly preferred indication</p>	



					<p>ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of MHC class II products, such as HLA-DR antigens, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Lamour et al., Clin Exp Immunol 89(2):217-222 (1992); Hurme and Sihvola, Immunol Lett 20(3):217-222 (1989); Gansbacher and Zier, Cell Immunol 117(1):22-34 (1988); and Itoh et al., J Histochem Cytochem 40(11):1675-1683, the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to</p>	<p>is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. An additional preferred indication is infection (e.g., AIDS, and/or as described below under "Infectious Disease"). Preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"), and neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancer, such as, for example, leukemia, lymphoma, and prostate, breast,</p>
--	--	--	--	--	---	--

412	HTDAA78	926			immunomodulatory factors.	lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy.
			Production of ICAM-1		Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Rolfe BE, et al., Atherosclerosis, 149(1):99-110 (2000); Panettieri RA Jr, et al., J Immunol, 154(5):2358-2365 (1995); and, Grunstein MM, et al., Am J Physiol Lung Cell Mol Physiol, 278(6):L1154-L1163 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include Aortic Smooth Muscle Cells (AOSMC);	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Vascular Disease, Atherosclerosis, Restenosis, Stroke, and Asthma.

412	HTDAA78	926	<p>Activation of Natural Killer Cell ERK Signaling Pathway.</p>	<p>such as bovine AOSMC.</p> <p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary natural killer cells that may be used according to these assays include the human natural killer cell lines (for example, NK-YT cells which have cytolytic and cytotoxic activity) or primary NK cells.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating natural killer cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating natural killer cell differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell differentiation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity") and infections (e.g., as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include cancers such as, kidney, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary cancer, lymphoma and leukemias. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Other highly preferred indications include, pancytopenia, leukopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), arthritis, asthma, AIDS, granulomatous disease, inflammatory bowel disease,</p>
-----	---------	-----	---	---	---

413	HTEAG62	927	Activation of Adipocyte PI3 Kinase Signalling Pathway	<p>Kinase assay. Kinase assays, for example an GSK-3 assays, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>sepsis, psoriasis, immune reactions to transplanted organs and tissues, endocarditis, meningitis, Lyme Disease, and allergies.</p> <p>A highly preferred embodiment of the invention includes a method for increasing adipocyte survival. An alternative highly preferred embodiment of the invention includes a method for decreasing adipocyte survival. A preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. A highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Preferred indications include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"), blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to</p>
-----	---------	-----	---	---	---

413	HTEAG62	927	Regulation of transcription of Malic	Assays for the regulation of transcription of Malic Enzyme are well-known in the art	<p>diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Highly preferred indications include neoplasms and cancer, such as, lipoma, liposarcoma, lymphoma, leukemia and breast, colon, and kidney cancer. Additional highly preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred indication is diabetes mellitus.</p> <p>An additional highly preferred indication is a complication</p>
-----	---------	-----	--------------------------------------	--	---

			<p>Enzyme in adipocytes</p>	<p>and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate transcription of Malic Enzyme, a key enzyme in lipogenesis. Malic enzyme is involved in lipogenesis and its expression is stimulated by insulin. ME promoter contains two direct repeat (DRL)-like elements MEp and MEEd identified as putative PPAR response elements. ME promoter may also responds to AP1 and other transcription factors. Exemplary assays that may be used or routinely modified to test for regulation of transcription of Malic Enzyme (in adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Streeter, R.S., et al., Mol Endocrinol, 12(11):1778-91 (1998); Garcia-Jimenez, C., et al., Mol Endocrinol, 8(10):1361-9 (1994); Barroso, I., et al., J Biol Chem, 274(25):17997-8004 (1999); Ippenber, A., et al., J Biol Chem, 272(32):20108-20117 (1997); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays includes the H4IIE rat liver hepatoma cell line.</p>	<p>associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	-----------------------------	--	---

414	HTECB02	928	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al., FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
415	HTECC15	929	Activation of transcription through NFKB response element in neuronal cells (such as SKNMC cells).	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of neuronal genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Gill JS, et al.,	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Neurological Diseases and Disorders (e.g. Alzheimer's Disease, Parkinson's Disease, Brain Cancer, Seizures).

416	HTEDF18	930	Production of MIP1alpha	<p>Neurobiol Dis, 7(4):448-461 (2000); Tamatani M, et al., J Biol Chem, 274(13):8531-8538 (1999); Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997); Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Neuronal cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary neuronal cells that may be used according to these assays include the SKNMC neuronal cell line.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS,</p>
				<p>MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and</p>	



417	HTEDJ28	931	<p>chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
			<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly</p>

				<p>agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity.</p>
--	--	--	--	---	---

417	HTEDJ28	931	Activation of transcription through NFAT response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention)	Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
					Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example,

418	HTEDS12	932	Activation of Adipocyte ERK Signaling Pathway	<p>include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Aramburu et al., J Exp Med 182(3):801-810 (1995); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. NK cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human NK cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
				<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic</p>

				<p>(1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with</p>
--	--	--	--	---	---

419	HTEED26	933	Activation of transcription through GAS response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992);	insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
				Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under	

420	HTEED26	934	Activation of transcription through GAS response element in immune cells (such as T-cells).	<p>Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>"Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
			<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or</p>	

421	HTEEF26	935		<p>85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>"Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
			Production of MIP1alpha	<p>MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention)</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues,</p>



422	HTEEF26	936	Production of MIP1alpha	<p>include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such</p>	<p>hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute</p>	

423	HTEEW69	937	Activation of transcription through cAMP response element in immune cells (such as T-cells).	assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
				Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP, bind to CREB transcription factor, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or	Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly

				<p>routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
423	HTEW69	937	Activation of transcription through GAS response element in immune cells (such as T-cells).	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and</p>

				<p>antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOLT4 cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
423	HTEEW69	937	<p>Activation of transcription through NFKB response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm,</p>	<p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon,</p>

423	HTEEW69	937	Activation of transcription through NFKB response element in immune cells (such as B-cells).	<p>Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOLT4, that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Gri G, et al., Biol Chem, 273(11):6431-6438 (1998); Pyatt DW, et al., Cell Biol Toxicol 2000;16(1):41-51 (2000); Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997);</p>	<p>pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p> <p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Autoimmunity, Allergy and Asthma</p>
-----	---------	-----	--	---	--

423	HTEW69	937		<p>Aramburu et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary immune cells that may be used according to these assays include the Reh B-cell line.</p> <p>CD152 FMA1. CD152 (a.k.a. CTLA-4) expression is restricted to activated T cells. CD152 is a negative regulator of T cell proliferation. Reduced CD152 expression has been linked to hyperproliferative and autoimmune diseases. Overexpression of CD152 may lead to impaired immunoresponses. Assays for immunomodulatory proteins important in the maintenance of T cell homeostasis and expressed almost exclusively on CD4+ and CD8+ T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, maintain T cell homeostasis, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD152, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or</p>	<p>A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
-----	--------	-----	--	--	---

			antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); McCoy et al., Immunol Cell Biol 77(1):1-10 (1999); Oostervegal et al., Curr Opin Immunol 11(3):294-300 (1999); and Saito T, Curr Opin Immunol 10(3):313-321 (1998), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy. An additional preferred indication is infection (e.g., as described below under "Infectious Disease").
424	HTEGS07	938	Activation of transcription through serum response element in immune cells (such as T-cells).	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and

			<p>include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
425	HTEGS11	939	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated</p>



426	HTEHA56	940	<p>activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly</p>
			<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation,</p>	
			<p>Activation of Adipocyte ERK Signaling Pathway</p>	

				<p>activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and</p>
--	--	--	--	---	---

426	HTEHA56	940	Activation of Adipocyte PI3 Kinase Signalling Pathway	<p>Kinase assay. Kinase assays, for example an GSK-3 assays, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to</p>	<p>impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred embodiment of the invention includes a method for increased adipocyte survival. An alternative highly preferred embodiment of the invention includes a method for decreasing adipocyte survival. A preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. Highly</p>
-----	---------	-----	---	--	---

				<p>test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Preferred indications include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"), blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications</p>
--	--	--	--	---	--

427	HTEHU59	941	Calcium flux in chondrocytes	<p>Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux in chondrocytes include assays disclosed in: Asada S, et al., Inflamm Res, 50(1):19-23 (2001);</p>	<p>associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Highly preferred indications include neoplasms and cancer, such as, lipoma, liposarcoma, lymphoma, leukemia and breast, colon, and kidney cancer. Additional highly preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Bone and Cartilage Diseases, including but not limited to Arthritis, Cartilage repair, Bone Repair, Osteoporosis, and related tumors including chondrosarcomas, chondroblastomas, and chondromas.</p>
-----	---------	-----	------------------------------	---	---

427	HTEHU59	941	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Schwartz Z, et al., J Bone Miner Res, 6(7):709-718 (1991); Iannotti JP, et al., J Bone Joint Surg Am, 67(1): 113-120 (1985); Sullivan E., et al., Methods Mol Biol 1999; 114:125-133 (1999), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include bovine chondrocytes.</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis,</p>
-----	---------	-----	--	--	---

428	HTEJD29	942	Stimulation of Calcium Flux in pancreatic beta cells.	<p>4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the SUPT cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include</p>
				<p>and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>

429	HTEKM46	943	Calcium flux in chondrocytes	<p>(1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p> <p>Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux in chondrocytes include assays disclosed in: Asada S, et al., Inflamm Res, 50(1):19-23 (2001); Schwartz Z, et al. J Bone Miner Res.</p>	<p>weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Bone and Cartilage Diseases, including but not limited to Arthritis, Cartilage repair, Bone Repair, Osteoporosis, and related tumors including chondrosarcomas, chondroblastomas, and chondromas.</p>
-----	---------	-----	------------------------------	---	--



430	HTEMQ17	944		<p>6(7):709-718 (1991); Iannotti JP, et al., J Bone Joint Surg Am, 67(1): 113-120 (1985); Sullivan E., et al., Methods Mol Biol 1999; 114:125-133 (1999), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include bovine chondrocytes.</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of asthma, allergy, hypersensitivity and inflammation.</p>
			<p>Regulation of apoptosis of immune cells (such as mast cells).</p>	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate caspase protease-mediated apoptosis in immune cells (such as, for example, in mast cells). Mast cells are found in connective and mucosal tissues throughout the body, and their activation via immunoglobulin E -antigen, promoted by T helper cell type 2 cytokines, is an important component of allergic disease. Dysregulation of mast cell apoptosis may play a role in allergic disease and mast cell tumor survival. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity induced by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Masuda A, et al., J Biol Chem, 276(28):26107-26113 (2001); Yeatman CF 2nd, et al., J Exp</p>	

431	HTENR63	945		<p>Med, 192(8):1093-1103 (2000); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary immune cells that may be used according to these assays include mast cells such as the HMC human mast cell line.</p>	<p>Assays for the regulation of transcription through the PEPCK promoter are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the PEPCK promoter in a reporter construct and regulate liver gluconeogenesis. Exemplary assays for regulation of transcription through the PEPCK promoter that may be used or routinely modified to test for PEPCK promoter activity (in hepatocytes) of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Lochhead et al., Diabetes 49(6):896-903 (2000); and Yeagley et al., J Biol Chem 275(23):17814-17820 (2000), the contents of each of which is herein incorporated by</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., an infectious diseases or disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity.</p>
-----	---------	-----	--	--	--	--

432	HTGGM44	946	<p>Regulation of viability and proliferation of pancreatic beta cells.</p>	<p>reference in its entirety. Hepatocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary liver hepatoma cells that may be used according to these assays include H4Ile cells, which contain a tyrosine amino transferase that is inducible with glucocorticoids, insulin, or cAMP derivatives.</p>	<p>Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include glycogen storage disease (e.g., glycogenoses), hepatitis, gallstones, cirrhosis of the liver, degenerative or necrotic liver disease, alcoholic liver diseases, fibrosis, liver regeneration, metabolic disease, dyslipidemia and cholesterol metabolism, and hepatocarcinomas. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), infection (e.g., an infectious disease and/or disorder as described below under "Infectious Disease"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), and neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"). Additional preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, and urinary cancer. A highly preferred indication is liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic</p>
-----	---------	-----	--	--	---

			<p>the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Friedrichsen BN, et al., Mol Endocrinol, 15(1):136-48 (2001); Huotari MA, et al., Endocrinology, 139(4):1494-9 (1998); Hugl SR, et al., J Biol Chem 1998 Jul 10;273(28):17771-9 (1998), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p>	<p>neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
433	HTHBZ06	947	<p>Stimulation of insulin secretion from pancreatic beta cells.</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy,</p>

			<p>of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., <i>Am J Physiol</i>, 277(4 Pt 2):R959-66 (1999); Li, M., et al., <i>Endocrinology</i>, 138(9):3735-40 (1997); Kim, K.H., et al., <i>FEBS Lett</i>, 377(2):237-9 (1995); and, Miraglia S et. al., <i>Journal of Biomolecular Screening</i>, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al.</p>	<p>diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	---

434	HTLAP64	948	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Endocrinology 1992 130:167.</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis,</p>
-----	---------	-----	--	---	---

435	HTLBT80	949	Activation of Adipocyte ERK Signaling Pathway	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast</p>	<p>meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders").</p> <p>Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as</p>
-----	---------	-----	---	---	--

				<p>cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin)). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>
--	--	--	--	---	--



435	HTLBT80	949	<p>Activation of transcription through the EGR (Early Growth Response) element in immune cells (such as B-cells).</p>	<p>Assays for the activation of transcription through the EGR response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate EGR transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the EGR response element that may be used or routinely modified to test EGR response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Richards JD, et al., J Immunol, 166(6):3855-3864 (2001); Dinkel, A, et al., J Exp Med, 188(12):2215-2224 (1998); and, Newton, JS, et al., Eur J Immunol 1996 Apr;26(4):811-816 (1996), the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary immune cells that may be used according to these assays include the Raji B-cell line.</p>	<p>dysplasia.</p> <p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Autoimmunity, Allergy and Asthma.</p>
435	HTLBT80	949	<p>Activation of transcription through the EGR (Early Growth Response) element in immune cells (such as B-cells).</p>	<p>Assays for the activation of transcription through the EGR response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate EGR transcription factors and modulate expression of immunomodulatory genes.</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Autoimmunity, Allergy and Asthma.</p>

435	HTLBT80	949	<p>expression of immunomodulatory genes. Exemplary assays for transcription through the EGR response element that may be used or routinely modified to test EGR response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Richards JD, et al., <i>J Immunol</i>, 166(6):3855-3864 (2001); Dinkel, A, et al., <i>J Exp Med</i>, 188(12):2215-2224 (1998); and, Newton, JS, et al., <i>Eur J Immunol</i> 1996 Apr;26(4):811-816 (1996), the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary immune cells that may be used according to these assays include the Raji B-cell line.</p> <p>Assays for the activation of transcription through the NFkB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFkB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFkB response element that may be used or routinely modified to test NFkB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Gri G, et al., <i>Biol Chem</i>, 273(11):6431-6438 (1998);</p>	<p>expression of immunomodulatory genes. Exemplary assays for transcription through the EGR response element that may be used or routinely modified to test EGR response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Richards JD, et al., <i>J Immunol</i>, 166(6):3855-3864 (2001); Dinkel, A, et al., <i>J Exp Med</i>, 188(12):2215-2224 (1998); and, Newton, JS, et al., <i>Eur J Immunol</i> 1996 Apr;26(4):811-816 (1996), the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary immune cells that may be used according to these assays include the Raji B-cell line.</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Autoimmunity, Allergy and Asthma</p>
-----	---------	-----	---	--	---

436	HTLDA84	950		<p>Pyatt DW, et al., Cell Biol Toxicol 2000;16(1):41-51 (2000); Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997); Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary immune cells that may be used according to these assays include the Reh B-cell line.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFN<math>\gamma</math>. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFN<math>\gamma</math>. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred</p>
			Production of IFN $\gamma$ using a T cells	<p>IFN<math>\gamma</math> plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFN<math>\gamma</math> promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity.</p>	

				<p>Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
437	HTLDN29	951	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as</p>

437	HTLDN29	951	Production of MIP1alpha	<p>modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An</p>
-----	---------	-----	-------------------------	---	---	--

			<p>activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture.</p>	<p>alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	---	--

438	HTLDU78	952	Regulation of viability and proliferation of pancreatic beta cells.	<p>which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Friedrichsen BN, et al., Mol Endocrinol, 15(1):136-48 (2001); Huotari MA, et al., Endocrinology, 139(4):1494-9 (1998); Hugi SR, et al., J Biol Chem 1998 Jul 10;273(28):17771-9 (1998), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an</p>	<p>A highly preferred indication is diabetes mellitus.</p> <p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
-----	---------	-----	---	---	--

438	HTLDU78	952	Proliferation, differentiation, and/or cytokine production in immune cells (such as T-cells).	<p>X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p> <p>Kinase assays, for example kinase assays for members of the MAP kinase family (including p38, JAK, and ERK) are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, differentiation., and/or cytokine production in immune cells such as T-cells. Exemplary assays for MAP kinase family members that may be used or routinely modified to test polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Rincon M., Curr Opin Immunol; 13(3):339-345 (2001); Wang LH, et al., J Immunol, 162(7):3897-3904 (1999); Sakamoto H, et al., J Biol Chem, 275(46):35857-35862 (2000), the contents of each of which are herein incorporated by reference in its entirety. Exemplary immune cells (for example, T-cells) that may be used according to these assays include the mouse CTLL cell line.</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of inflammation, infection, allergy, asthma, autoimmunity, and cancer.</p>
438	HTLDU78	952	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred</p>	



			<p>cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the</p>	<p>indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	---	---

439	HTLEC82	953	<p>art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>Assays for the activation of transcription through the AP1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely modified to test AP1-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4 responsive suspension-culture cell line.</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
-----	---------	-----	---	---

440	HTLEM16	954	<p>Regulation of apoptosis of immune cells (such as mast cells).</p>	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate caspase protease-mediated apoptosis in immune cells (such as, for example, in mast cells). Mast cells are found in connective and mucosal tissues throughout the body, and their activation via immunoglobulin E -antigen, promoted by T helper cell type 2 cytokines, is an important component of allergic disease. Dysregulation of mast cell apoptosis may play a role in allergic disease and mast cell tumor survival. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity induced by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Masuda A, et al., J Biol Chem, 276(28):26107-26113 (2001); Yeatman CF 2nd, et al., J Exp Med, 192(8):1093-1103 (2000); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary immune cells that may be used according to these assays include mast cells such as the HMC</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of asthma, allergy, hypersensitivity and inflammation.</p>
-----	---------	-----	--	--	--

440	HTLEM16	954	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	human mast cell line. Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2):495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic</p>
-----	---------	-----	--	---	--

					<p>overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly</p>
--	--	--	--	--	--

441	HTLEV48	955	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these	preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
					Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke

442	HTLFA13	956	Regulation of viability and proliferation of pancreatic beta cells.	assays include microvascular endothelial cells (MVEC). Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Friedrichsen BN, et al., Mol Endocrinol, 15(1):136-48 (2001); Huotari MA, et al., Endocrinology, 139(4):1494-9 (1998); Hugl SR, et al., J Biol Chem 1998 Jul 10;273(28):17771-9 (1998), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.
-----	---------	-----	---	---	--

443	HTLF173	957		<p>insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p> <p>TNF<math>\alpha</math> FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor <math>\alpha</math> (TNF<math>\alpha</math>), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and</p>	<p>Production of TNF <math>\alpha</math> by dendritic cells</p>	<p>insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p> <p>TNF<math>\alpha</math> FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor <math>\alpha</math> (TNF<math>\alpha</math>), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and</p>	<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF <math>\alpha</math> production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF <math>\alpha</math> production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS,</p>
-----	---------	-----	--	--	---	--	---



444	HTLGI89	958	Stimulation of Calcium Flux in pancreatic beta cells.	<p>Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 ( Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec:10(8):535-41</p>	<p>granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			Stimulation of Calcium Flux in pancreatic beta cells.	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include</p>	

445	HTLIF11	959	Protection from Endothelial Cell Apoptosis.	<p>(1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p> <p>Caspase Apoptosis Rescue. Assays for caspase apoptosis rescue are well known in the art and may be used or routinely modified to assess the ability of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to inhibit caspase protease-mediated apoptosis. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis rescue of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Romeo et al., Cardiovasc Res 45(3): 788-794 (2000); Messmer et al., Br J Pharmacol 127(7): 1633-1640 (1999); and J Atheroscler Thromb 3(2): 75-80 (1996); the contents of</p>	<p>weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred</p>
-----	---------	-----	---	--	--

				<p>each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (BAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign</p>
--	--	--	--	---	---

446	HTLIF12	960	Activation of transcription through GAS response element	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-	<p>dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred</p>
-----	---------	-----	--	--	---

			known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.	indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.	
447	HTLIF12	961	Activation of transcription through GAS response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or	Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for

448	HTLIF12	962	<p>as T-cells).</p> <p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p> <p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma,</p>
-----	---------	-----	--	---	--

449	HTLIF12	963	<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
			<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's</p>

450	HTLIF12	964	<p>antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
				<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon,</p>
			<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of</p>	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>



451	HTLIF12	965	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT</p>	<p>the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p> <p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary</p>
-----	---------	-----	--	---	---	--

			transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.	cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
452	HTNAM63	966	Activation of transcription through cAMP response element in immune cells (such as T-cells).	Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune

				<p>wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
453	HTNBK13	967	<p>Regulation of viability and proliferation of pancreatic beta cells.</p>	<p>Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension,</p>

453	HTNBK13	967	Production of TNF alpha by dendritic cells	<p>active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Friedrichsen BN, et al., Mol Endocrinol, 15(1):136-48 (2001); Huotari MA, et al., Endocrinology, 139(4):1494-9 (1998); Hugl SR, et al., J Biol Chem 1998 Jul 10;273(28):17771-9 (1998), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Afari et al. Endocrinology 1992 130:167.</p> <p>TNFα FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of</p>	<p>stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus,</p>	

454	HTOAI50 • 968	Production of IFNgamma using a T cells	<p>the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>IFNgamma FMAAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory</p>	<p>Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the</p>
-----	---------------	--	---	---

			<p>cytokine. IFN<math>\gamma</math> promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFN<math>\gamma</math>), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated</p>	<p>invention includes a method for inhibiting the production of IFN<math>\gamma</math>. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	---

455	HTOAM11	969		<p>by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory</p>
-----	---------	-----	--	--	---

455	HTOAM11	969	Activation of transcription through cAMP response element in immune cells (such as T-cells).	assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.  Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is a suspension	bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.  Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis,
-----	---------	-----	--	--	---



455	HTOAM11	969	Activation of transcription through GAS response element in immune cells (such as T-cells).	<p>culture of IL-2 dependent cytotoxic T cells.</p> <p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>meningitis, Lyme Disease, and asthma and allergy.</p> <p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and</p>
-----	---------	-----	---	--	---

455	HTOAM11	969	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>allergy.</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis,</p>
-----	---------	-----	--	---

455	HTOAM11	969	Stimulation of insulin secretion from pancreatic beta cells.	<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1</p>	<p>meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
-----	---------	-----	--	--	---

455	HTOAM11	969		<p>cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p> <p>Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays</p>	<p>cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. An additional highly preferred indication includes infection (e.g., AIDS, and/or as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma),</p>
-----	---------	-----	--	---	--	---

455	HTOAM11	969	Production of IFNgamma using a T cells	<p>include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication is infection (e.g., tuberculosis, and infections associated with granulomatous disease, and osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
					<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFN<math>\gamma</math>. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFN<math>\gamma</math>. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease</p>

				<p>art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFN<math>\gamma</math>), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be</p>	<p>(e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	--	--

456	HTODH57	970	<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>	<p>preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysplastic disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
457	HTODH83	971	<p>Activation of transcription through serum response element</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the</p>

			<p>be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	--	---



457	HTODH83	971	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
458	HTOE16	972	Production of IL-6	IL-6 FMT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability	A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-

459	HTOGR38	973	<p>of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the</p>	<p>mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating natural killer cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell proliferation. A highly preferred embodiment of the invention includes a method for</p>

460	HTOHO21	974	Activation of Adipocyte ERK Signaling Pathway	<p>invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary natural killer cells that may be used according to these assays include the human natural killer cell lines (for example, NK-YT cells which have cytolytic and cytotoxic activity) or primary NK cells.</p>	<p>stimulating natural killer cell differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell differentiation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity") and infections (e.g., as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include cancers such as, kidney, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary cancer, lymphoma and leukemias. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Other highly preferred indications include, pancytopenia, leukopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), arthritis, asthma, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, immune reactions to transplanted organs and tissues, endocarditis, meningitis, Lyme Disease, and allergies.</p>
460	HTOHO21	974	Activation of Adipocyte ERK Signaling Pathway	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte</p>

				<p>and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension,</p>
--	--	--	--	--	---

460	HTOH021	974	<p>stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
			<p>A highly preferred embodiment of the invention includes a method for increasing muscle cell survival. An alternative highly preferred embodiment of the invention includes a method for decreasing muscle cell survival. A preferred embodiment of the invention includes a method for stimulating muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is</p>
			<p>Kinase assay. Kinase assays, for example an GSK-3 kinase assay, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including</p>
			<p>Activation of Skeletal Muscle Cell PI3 Kinase Signalling Pathway</p>

			<p>antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Rat myoblast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary rat myoblast cells that may be used according to these assays include L6 cells. L6 is an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuses to form multinucleated myotubes and striated fibers after culture in differentiation media.</p>	<p>stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is inhibited. A preferred embodiment of the invention includes a method for stimulating muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is inhibited. Highly preferred indications include disorders of the musculoskeletal system. Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia,</p>
--	--	--	---	--

461	HTOHQ05	975	Production of MIP1alpha	MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate	endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infections (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal system including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include: myopathy, atrophy, congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, and vascular disease. Highly preferred indications include neoplasms and cancer, such as, rhabdomyoma, rhabdosarcoma, stomach, esophageal, prostate, and urinary cancer. Preferred indications also include breast, lung, colon, pancreatic, brain, and liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, hyperplasia, metaplasia, and/or dysplasia. A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications
-----	---------	-----	-------------------------	--	---

			<p>chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
462	HTOJL95	976	<p>Upregulation of CD71 and activation of T cells</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. Preferred indications include blood disorders (e.g., as</p>



				<p>proliferating. Assays for immunomodulatory proteins expressed on activated T cells, B cells, and most proliferating cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD71, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Afetra et al., Ann Rheum Dis 52(6):457-460 (1993), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include infection. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
--	--	--	--	---	---

463	HTOJL95	977	<p>Upregulation of CD71 and activation of T cells</p>	<p>CD71 FMAT. CD71 is the transferrin receptor. Transferrin is a major iron carrying protein that is essential for cell proliferation. CD71 is expressed predominantly on cells that are actively proliferating. Assays for immunomodulatory proteins expressed on activated T cells, B cells, and most proliferating cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD71, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Afetra et al., Ann Rheum Dis 52(6):457-460 (1993), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include infection. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
-----	---------	-----	---	---	---

464	HTPDU17	978	Production of IL-4	<p>express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>IL-4 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells that stimulate B cells, T cells, macrophages and mast cells and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cells, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-4, and the stimulation of immune cells, such as B cells, T cells, macrophages and mast cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):277-283 (1994); Yssel et al., Res Immunol 144(8):610-616 (1993); Bagley et al., Nat Immunol 1(3):257-261 (2000); and van der Graaff et al., Rheumatology (Oxford)</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-4 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-4 production. A highly preferred indication includes asthma. A highly preferred indication includes allergy. A highly preferred indication includes rhinitis. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of</p>
-----	---------	-----	--------------------	---	---

465	HTSFJ32	979	Activation of transcription through serum response element in immune cells (such as T-cells).	38(3):214-220 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
			Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal,	

465	HTSFJ32	979	<p>assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			<p>Kinase assay. Kinase assays, for example an GSK-3 kinase assay, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein</p>	<p>A highly preferred embodiment of the invention includes a method for increasing muscle cell survival. An alternative highly preferred embodiment of the invention includes a method for decreasing muscle cell survival. A preferred embodiment of the invention includes a method for stimulating muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is inhibited. A preferred embodiment of the invention includes a method for stimulating muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is inhibited. Highly preferred indications include disorders of the musculoskeletal system. Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), endocrine disorders (e.g.,</p>

				<p>incorporated by reference in its entirety.</p> <p>Rat myoblast cells that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>Exemplary rat myoblast cells that may be used according to these assays include L6 cells. L6 is an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuses to form multinucleated myotubes and striated fibers after culture in differentiation media.</p>	<p>as described below under "Endocrine Disorders"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infections (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture).</p> <p>An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal system</p>
--	--	--	--	---	--

466	HTTCB60	980			<p>including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include: myopathy, atrophy, congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, and vascular disease. Highly preferred indications include neoplasms and cancer, such as, rhabdomyoma, rhabdosarcoma, stomach, esophageal, prostate, and urinary cancer. Preferred indications also include breast, lung, colon, pancreatic, brain, and liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with</p>
					<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of</p>

467	HTTEE41	981	Stimulation of Calcium Flux in pancreatic beta cells.	<p>Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p> <p>Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin I.S. et al., Endocrinology.</p>	<p>obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious</p>	



468	HTTEZ02	982	Endothelial Cell Apoptosis	<p>136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 ( Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p> <p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely</p>	<p>Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing)</p>	

				<p>modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary</p>
--	--	--	--	--	---

469	HTWEH94	983	Stimulation of insulin secretion from	Assays for measuring secretion of insulin are well-known in the art and may be used	<p>cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication</p>
-----	---------	-----	---------------------------------------	---	--

			pancreatic beta cells.	<p>or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible</p>	<p>associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	------------------------	---	---

470	HTXBD09	984	Activation of transcription through NFKB response element in epithelial cells (such as HELA cells).	<p>insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p> <p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of epithelial genes.</p> <p>Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Kaltschmidt B, et al., Oncogene, 18(21):3213-3225 (1999); Beetz A, et al., Int J Radiat Biol, 76(11):1443-1453 (2000); Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997); Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Epithelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary epithelial cells that may be used according to these assays include the HELA cell line.</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Wound Healing, and Inflammation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include include inflammation and inflammatory disorders.</p>
471	HTXDB22	985	Activation of	<p>Assays for the activation of transcription</p>	Highly preferred indications include neoplastic diseases

		transcription through GAS response element in immune cells (such as T-cells).	through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the SUPT cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).	<p>(e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p> <p>Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
472	HTXDC38	986	Assays for the activation of transcription through the Gamma Interferon Activation	Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below

			<p>GAS response element in immune cells (such as T-cells).</p>	<p>Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the SUPT cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
473	HTXDC77	987	<p>Activation of transcription through serum response element</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of</p>

			<p>in immune cells (such as natural killer cells).</p>	<p>be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm. Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	--	---	---



474	HTXDD61	988	Upregulation of CD69 and activation of T cells	<p>CD69 FMAT. CD69 is an activation marker that is expressed on activated T cells, B cells, and NK cells. CD69 is not expressed on resting T cells, B cells, or NK cells. CD69 has been found to be associated with inflammation. Assays for immunomodulatory proteins expressed in T cells, B cells, and leukocytes are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD69, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ferenczi et al., J Autoimmun 14(1):63-78 (2000); Werfel et al., Allergy 52(4):465-469 (1997); Taylor-Fishwick and Siegel, Eur J Immunol 25(12):3215-3221 (1995); and Afetra et al., Ann Rheum Dis 52(6):457-460 (1993), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated</p>	<p>A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for activation B cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating B cells. A highly preferred embodiment of the invention includes a method for activating NK cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting activation of and/or inactivation NK cells. Highly preferred indications include inflammation and inflammatory disorders (e.g., as described below under "Immune Activity"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response and alternatively suppressing a T cell-mediated immune response, and boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. An additional highly preferred indication includes infection (e.g., as described below under "Infectious Disease"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis,</p>
-----	---------	-----	--	--	--

475	HTXDG92	989			<p>using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Kinase assays, for example kinase assays for members of the MAP kinase family (including p38, JAK, and ERK) are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, differentiation, and/or cytokine production in immune cells such as T-cells. Exemplary assays for MAP kinase family members that may be used or routinely modified to test polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Rincon M., Curr Opin Immunol; 13(3):339-345 (2001); Wang LH, et al., J Immunol, 162(7):3897-3904 (1999); Sakamoto H, et al., J Biol Chem, 275(46):35857-35862 (2000), the contents of each of which are herein incorporated by reference in its entirety. Exemplary immune cells (for example, T-cells) that may be used according to these assays include the mouse CTLL cell line.</p>	<p>meningitis, Lyme Disease, inflammation and inflammatory disorders, asthma, and allergies. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of inflammation, infection, allergy, asthma, autoimmunity, and cancer.</p>
-----	---------	-----	--	--	---	--

475	HTXDG92	989	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the SUPT cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
476	HTXET11	990	<p>Proliferation,</p>	<p>Kinase assays, for example kinase assays</p>	<p>Preferred embodiments of the invention include using</p>

		differentiation, and/or cytokine production in immune cells (such as T-cells).	for members of the MAP kinase family (including p38, JAK, and ERK) are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, differentiation, and/or cytokine production in immune cells such as T-cells. Exemplary assays for MAP kinase family members that may be used or routinely modified to test polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Rincon M., Curr Opin Immunol; 13(3):339-345 (2001); Wang LH, et al., J Immunol, 162(7):3897-3904 (1999); Sakamoto H, et al., J Biol Chem, 275(46):35857-35862 (2000), the contents of each of which are herein incorporated by reference in its entirety. Exemplary immune cells (for example, T-cells) that may be used according to these assays include the mouse CTLL cell line.	polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of inflammation, infection, allergy, asthma, autoimmunity, and cancer.
477	HTXFA72	991	Stimulation of Calcium Flux in pancreatic beta cells.	<p>A highly preferred indication is diabetes mellitus.</p> <p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease,</p>

478	HTXJY08	992	<p>activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 ( Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
			<p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of</p>	<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic</p>

			killer cells).	<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Aramburu et al., J Exp Med 182(3):801-810 (1995); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. NK cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human NK cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
478	HTXJY08	992	Activation of transcription through serum response element in immune cells (such as natural killer cells).	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications</p>	

479	HTXKF95	993	Activation of Skeletal Muscle Cell ERK	<p>(including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
				<p>Kinase assay. Kinase assays, for examplek Elk-1 kinase assays, for ERK signal</p>	<p>Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine</p>

			<p>transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Rat myoblast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary rat myoblast cells that may be used according to these assays include L6 cells. L6 is an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuses to form multinucleated myotubes and striated fibers after culture in differentiation media.</p>	<p>Disorders") and disorders of the musculoskeletal system. Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	---	--



480	HTXMZ07	994	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-</p>	<p>Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include: myopathy, atrophy, congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, and vascular disease. Highly preferred indications include neoplasms and cancer, such as, rhabdomyoma, rhabdosarcoma, stomach, esophageal, prostate, and urinary cancer. Highly preferred indications also include breast, lung, colon, pancreatic, brain, and liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma</p>
-----	---------	-----	--	---	---

481	HUFCL31	995		<p>3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>(e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			Production of GM-CSF	<p>GM-CSF/FMAT. GM-CSF is expressed by activated T cells, macrophages, endothelial cells, and fibroblasts. GM-CSF regulates differentiation and proliferation of granulocytes- macrophage progenitors and enhances antimicrobial activity in neutrophils, monocytes and macrophage. Additionally, GM-CSF plays an important role in the differentiation of dendritic cells and monocytes, and increases antigen presentation. GM-CSF is considered to be a proinflammatory cytokine. Assays for immunomodulatory proteins that promote the production of GM-CSF are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of GM-CSF. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of GM-CSF. Highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., as described below under "Infectious Disease". Highly preferred indications include blood disorders (e.g., neutropenia (and the prevention of neutropenia (e.g., in HIV infected patients), and/or as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications also include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include asthma. Highly preferred indications include neoplastic diseases (e.g., leukemia (e.g., acute</p>

			mediate immunomodulation and modulate the growth and differentiation of leukocytes. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as GM-CSF, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Ye et al., J Leukoc Biol (58(2):225-233, the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC) or may be isolated using techniques disclosed herein or otherwise known in the art. Natural killer (NK) cells are large granular lymphocytes that have cytotoxic activity but do not bind antigen. NK cells show antibody-independent killing of tumor cells and also recognize antibody bound on target cells, via NK Fc receptors, leading to cell-mediated cytotoxicity.	lymphoblastic leukemia, and acute myelogenous leukemia), lymphoma (e.g., non-Hodgkin's lymphoma and Hodgkin's disease), and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications include: suppression of immune reactions to transplanted organs and tissues (e.g., bone marrow transplant); accelerating myeloid recovery; and mobilizing hematopoietic progenitor cells. Preferred indications include boosting a T cell-mediated immune response, and alternatively, suppressing a T cell-mediated immune response. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and allergy.
482	HUKBT67	996	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for</p>

				<p>proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis</p>
--	--	--	--	---	---

					<p>and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, stenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune</p>
--	--	--	--	--	--

483	HUKDF20	997	Regulation of transcription via DMEF1 response element in adipocytes and pre-adipocytes	<p>Assays for the regulation of transcription through the DMEF1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the DMEF1 response element in a reporter construct (such as that containing the GLUT4 promoter) and to regulate insulin production. The DMEF1 response element is present in the GLUT4 promoter and binds to MEF2 transcription factor and another transcription factor that is required for insulin regulation of Glut4 expression in skeletal muscle. GLUT4 is the primary insulin-responsive glucose transporter in fat and muscle tissue. Exemplary assays that may be used or routinely modified to test for DMEF1 response element activity (in adipocytes and pre-adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Thai, M.V., et al., J Biol Chem, 273(23):14285-92 (1998); Mora, S., et al., J Biol Chem, 275(21):16323-8 (2000); Liu, M.L., et al., J Biol Chem, 269(45):28514-</p>	<p>Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
-----	---------	-----	---	---	---

483	HUKDF20	997	<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>	<p>21 (1994); "Identification of a 30-base pair regulatory element and novel DNA binding protein that regulates the human GLUT4 promoter in transgenic mice", J Biol Chem. 2000 Aug 4;275(31):23666-73; Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Adipocytes and pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the mouse 3T3-L1 cell line which is an adherent mouse preadipocyte cell line. Mouse 3T3-L1 cells are a continuous substrain of 3T3 fibroblasts developed through clonal isolation. These cells undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation culture conditions.</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under</p>
-----	---------	-----	--	---	---

483	HUKDF20	997	<p>assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>“Hyperproliferative Disorders”). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
			<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under “Immune Activity”, “Blood-Related Disorders”, and/or “Cardiovascular Disorders”), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below</p>



483	HUKDF20	997	Production of ICAM-1	<p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly</p>	<p>85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke</p>
-----	---------	-----	----------------------	--	---	--

483	HUKDF20	997	<p>Activation of transcription through serum response element in immune cells (such as natural killer cells).</p>	<p>available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types.</p> <p>Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis.</p> <p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous</p>
-----	---------	-----	---	--	---

484	HUKDY82	998	Activation of transcription through GAS response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOLT4 cell line, that may be used according to these assays are publicly	activity.  disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
				Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL),	

485	HUSCJ14	999	Regulation of transcription through the FAS promoter element in hepatocytes	<p>available (e.g., through the ATCC).</p> <p>Assays for the regulation of transcription through the FAS promoter element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the FAS promoter element in a reporter construct and to regulate transcription of FAS, a key enzyme for lipogenesis. FAS promoter is regulated by many transcription factors including SREBP. Insulin increases FAS gene transcription in livers of diabetic mice. This stimulation of transcription is also somewhat glucose dependent. Exemplary assays that may be used or routinely modified to test for FAS promoter element activity (in hepatocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Xiong, S., et al., Proc Natl Acad Sci U.S.A., 97(8):3948-53 (2000); Roder, K., et al., Eur J Biochem, 260(3):743-51 (1999); Oskouian B, et al., Biochem J, 317 (Pt 1):257-65 (1996); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety.</p>	<p>plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
-----	---------	-----	---	---	--

485	HUSCJ14	999	Upregulation of CD71 and activation of T cells	<p>Hepatocytes that may be used according to these assays, such as H4IIE cells, are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays include rat liver hepatoma cell line(s) inducible with glucocorticoids, insulin, or cAMP derivatives.</p> <p>CD71 FMAT. CD71 is the transferrin receptor. Transferrin is a major iron carrying protein that is essential for cell proliferation. CD71 is expressed predominantly on cells that are actively proliferating. Assays for immunomodulatory proteins expressed on activated T cells, B cells, and most proliferating cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD71, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include infection. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's</p>
-----	---------	-----	--	--	--

486	HUSGL67	1000		<p>(2000); and Afetra et al., Ann Rheum Dis 52(6):457-460 (1993), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T</p>	<p>disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
				<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors,</p>	

487	HUSGU40	1001	Activation of transcription through serum response element in immune cells (such as T-cells).	cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.	and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
			Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al.,	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally,	

488	HUSIR.18	1002	Activation of transcription through serum response element in immune cells (such as T-cells).	<p>Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
			<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm-</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis.</p>	



				<p>Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
488	HUSIR18	1002	<p>Production of IL-10 and downregulation of immune responses</p>	<p>IL-10 FMAT. Assays for immunomodulatory proteins produced by activated T cells, B cells, and monocytes that exhibit anti-inflammatory activity and downregulate monocyte/macrophage function and expression of cytokines are well known in the art and may be used or routinely modified to assess the ability of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, and modulate immune cell function and cytokine production.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating the production of IL-10. An alternative preferred embodiment of the invention includes a method for inhibiting the production of IL-10. Highly preferred indications include inflammation and inflammatory disorders (e.g. inflammatory bowel disease). An additional highly preferred indication includes inflammatory bowel disease. Additional highly preferred indications include blood disorders (e.g., as described below under "Immune Activity" (e.g. autoimmune disorders), "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as</p>	

489	HUVDJ48	1003	<p>Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-10, and the downmodulation of immune responses. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent</p>	<p>described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, Crohn's disease, arthritis, AIDS, granulomatous disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy. An additional preferred indication is infection (e.g., as described below under "Infectious Disease").</p>
				<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or</p>

490	HWAA112	1004	Upregulation of CD71 and activation of T cells	<p>cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Friedrichsen BN, et al., Mol Endocrinol, 15(1):136-48 (2001); Huotari MA, et al., Endocrinology, 139(4):1494-9 (1998); Hugl SR, et al., J Biol Chem 1998 Jul 10;273(28):17771-9 (1998), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Afari et al. Endocrinology 1992 130:167.</p> <p>CD71 FMAT. CD71 is the transferrin receptor. Transferrin is a major iron carrying protein that is essential for cell proliferation. CD71 is expressed predominantly on cells that are actively proliferating. Assays for</p>	<p>blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-</p>	

				<p>immunomodulatory proteins expressed on activated T cells, B cells, and most proliferating cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD71, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Afetra et al., Ann Rheum Dis 52(6):457-460 (1993), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include infection. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
491	HWBBQ70	1005	Production of IL-6	IL-6 FMAT. IL-6 is produced by T cells	A highly preferred embodiment of the invention

				<p>and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by</p>	<p>includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia,</p>
--	--	--	--	---	---

492	HWBCN36	1006	Production of ICAM-1	reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities. Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
493	HWBDJ08	1007	Production of IL-6	IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
					A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood

				<p>disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture,</p>	<p>disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	--	--	--

493	HWBD108	1007	<p>Activation of transcription through NFAT response element in immune cells (such as T-cells).</p>	<p>which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays</p>	<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, liver and urinary pancreatic, esophageal, stomach, brain, lung, colon, cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
-----	---------	------	---	--	---



494	HWBFX16	1008	Activation of transcription through cAMP response element in immune cells (such as T-cells).	include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells. Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP, regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the HT2 cell line, which is a suspension culture of IL-2 dependent T cells that also respond to IL-4.	Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
495	HWDAC26	1009	Activation of	Kinase assay. JNK kinase assays for	A highly preferred embodiment of the invention includes

			<p>signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK kinase activity that may be used or routinely modified to test JNK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect</p>
--	--	--	--	---

					<p>vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vascularitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment</p>
--	--	--	--	--	--

496	HWDAG96	1010	Activation of transcription through API response element in immune cells (such as T-cells).	Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the	<p>/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia,</p>
-----	---------	------	---	---	--

497	HWDJ01	1011	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>contents of each of which are herein incorporated by reference in its entirety. Mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the HT2 cell line, which is an IL-2 dependent suspension culture cell line that also responds to IL-4.</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-</p>	<p>thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
				<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred</p>	

498	HWHPB78	1012	<p>Activation of transcription through NFKB response element in immune cells (such as natural killer cells).</p>	<p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997); Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its</p>	<p>2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS,</p>
-----	---------	------	--	---	---	---

499	HYABC84	1013	Stimulation of Calcium Flux in pancreatic beta cells.	<p>entirety. NK cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary NK cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p> <p>Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 ( Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly</p>	<p>granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
				<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>	

500	HYABC84	1014	Stimulation of Calcium Flux in pancreatic beta cells.	<p>available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>A highly preferred indication is diabetes mellitus.</p> <p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's</p>
-----	---------	------	---	--	--



				<p>Richardson SB, et al., Biochem J, 288 ( Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777</p> <p>Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--	--

TABLE 2

Clone ID	Contig ID:	SEQ ID NO:X	Analysis Method	PFam/NR Description	PFam/NR Accession Number	Score/Percent Identity	NT From	NT To
H6BSF56	762968	11	HMMER 2.1.1	PFAM: Zinc-binding dehydrogenases	PF00107	35.6	176	415
			blastx.2	CGI-63 PROTEIN.	sp Q9Y373 Q9Y373	94% 72% 44%	53 48 25	427 80 78
H6EDM64	841331	12	blastx.2	ANG2.	sp Q9UID3 Q9UID3	99% 100% 35% 34% 36%	922 107 2235 874 203	2451 922 2459 1038 310
H6EEC72	889401	13	blastx.2	hypothetical protein DKFZp434L061.1 - human	pir T43456 T43456	51% 70% 65% 53%	1459 1448 1484 137	365 807 927 96
HACBS22	847113	16	blastx.2	adenylate cyclase (EC 4.6.1.1) type III - rat	pir A39833 A39833	93% 99% 31% 25% 23%	416 6 2078 1445 917	2446 416 2299 1987 1111
HADDE71	839187	17	blastx.2	(AF002460) gene_id:F1D9.26~unknown protein [Arabidopsis thaliana]	dbj BA97098.1	91% 80% 90% 100% 100% 100% 100% 100% 100% 100%	499 480 506 520 520 520 520 520 520 520	666 665 667 666 666 666 666 666 666 666

[illegible]

[illegible][illegible]

[illegible]

[illegible][illegible]

090316Z-091801Z

[illegible]

[illegible]



[illegible]







HAGFS7	847120	24	blastx.2	X-LIKE 1 PROTEIN.	sp Q9Y485 Q9Y485	58%	9	872
HAGHN57	773286	25	blastx.2	MITOCHONDRIAL PROCESSING PEPTIDASE BETA SUBUNIT PRECURSOR 1	sp O75439 MPPB_HUM AN	98% 99% 100%	65 89 39	1444 1444 89
HAHEA15	847013	26	blastx.2	HYPOTHETICAL 31.4 KDA PROTEIN.	sp Q9NWD5 Q9NWD5	99% 76%	30 455	560 832
HAA47	534670	27	blastx.2	CDA14.	sp Q9NZA3 Q9NZA3	100%	17	157
HAY92	845601	28	blastx.2	ORF2-LIKE PROTEIN (FRAGMENT).	sp O00549 O00549	31% 52% 39% 25%	1721 2226 1602 661	2242 2333 1769 915
HJBV67	866415	29	blastx.2	SM-11044 BINDING PROTEIN (FRAGMENT).	sp Q9UHW8 Q9UHW8	99% 100%	116 13	1681 126
HAOAG15	852204	31	HMMER 2.1.1	PFAM: von Willebrand factor type A domain	PF00092	180.1	506	1057
			blastx.2	INTEGRIN ALPHA-10 PRECURSOR.	sp O75578 ITAG_HUMA N	98%	8	3508
HAQCE11	633730	33	blastx.2	(AY012159) virion-associated nuclear-shuttling protein [Mus musculus]	gb AAG42155.1	96% 32%	48 464	134 547
HATC103	580805	37	blastx.2	CDNA: FLJ121463 fis, clone COL04765.	sp BAB15071 BAB15071	68%	924	688
HBAGD86	838799	39	blastx.2	HYPOTHETICAL PROTEIN (FRAGMENT).	sp Q14287 Q14287	37%	801	559
HBDAB91	789532	41	blastx.2	PUTATIVE P150.	sp O00370 O00370	36% 40%	529 587	5 513
HBDAB91	864374	42	blastx.2	PUTATIVE P150.	sp O00370 O00370	36% 40%	849 907	307 833
HGBBC29	691473	43	blastx.2	BETA-1,4- GALACTOSYLTRANSFERASE.	sp O60513 O60513	98% 100%	65 1	1021 78
HBHAA05	603174	45	blastx.2	PRO2550.	sp AAG35515 AAG35515	71%	676	386
HBHAA81	846465	46	blastx.2	CDNA FLJ10724 FIS, CLONE NT2RP3001176.	sp Q9NVH9 Q9NVH9	27% 35%	49 1341	1356 1460
HBIAC29	831751	48	blastx.2	CDNA FLJ11730 fis, clone	sp BAB13898 BAB13898	100%	25	597

HBAB02	837309	50	blastx.2	HEMBA1005403. CDNA FLJ20062 FIS, CLONE COL01508.	sp Q9NXT6 Q9NXT6	100% 38% 35% 28%	2 1627 1616 1625	1267 1680 1675 1687
HBICK46	815649	53	HMMER 2.1.1 blastx.2	PFAM: WD domain, G-beta repeat DJ703H14.1 PROTEIN (FRAGMENT).	PF00400	36.6	790	867
HBIDS79	813588	54	WUblastx.6 4	(AK011059) putative [Mus musculus]	dbj BAB27367.1	92% 89% 93% 100% 66%	1119 1322 1032 1509 2	1325 1519 1127 1532 1075
HBIEL16	847030	56	blastx.2	PROTEIN ZERO RELATED PROTEIN.	sp O95297 O95297	98%	285	491
HBIG20	866159	58	HMMER 2.1.1 blastx.2	PFAM: Cytochrome c oxidase subunit III CYTOCHROME C OXIDASE POLYPEPTIDE III (EC 1.9.3.1).	PF00510 sp P00414 COX3_HUMA N	162.6 95%	321 9	551 617
HBKDI6	853358	59	blastx.2	CDNA FLJ20080 FIS, CLONE COL03184.	sp Q9NXS4 Q9NXS4	99%	8	1528
HBMBM96	561935	60	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	75%	814	518
HBMTX26	695704	63	blastx.2	HYPOTHETICAL PROTEIN (FRAGMENT).	sp Q14288 Q14288	44% 47% 61% 54% 44% 53% 66%	546 985 272 611 985 839 136	139 608 156 507 863 795 101
HBMTY48	637521	64	blastx.2	CDNA: FLJ23235 fis, clone CAS04980.	sp BAB15583 BAB15583	100%	54	941
HBMUH74	866160	65	blastx.2	(AL136843) hypothetical protein [Homo sapiens]	emb CAB66777.1	100%	11	427

HBMWE61	778066	66	blastx.2	(AF319977) mage-d3 [Mus musculus]	gb AAK01205.1	92%	302	586
						72%	585	863
						67%	585	863
						64%	579	863
						61%	585	851
						60%	588	863
						47%	576	836
						42%	576	863
						43%	585	851
						46%	585	851
						44%	585	863
						43%	585	863
						47%	594	845
						46%	579	824
						46%	594	836
						43%	588	851
						45%	594	854
						42%	579	851
						44%	585	854
						42%	579	845
						42%	588	863
						41%	579	854
						42%	585	827
						42%	591	842
						40%	573	845
						41%	585	875
						41%	585	836
						40%	594	848
						39%	579	851
						43%	579	851
						43%	579	851
						40%	588	857
						45%	579	821
						44%	600	821
						37%	576	863
						42%	591	851
						41%	579	848

[illegible]



					transposon MER37					60%	564	758
HCE3B04	831151	78		blastx.2	HYPOTHETICAL 31.3 KDA PROTEIN (FRAGMENT).		sp O43466 O43466			77%	430	564
HCEDR26	771144	80		blastx.2	CDNA FLJ20489 FIS, CLONE KAT08285.		sp Q9NX17 Q9NX17			49%	217	972
HCREEQ25	531784	82		blastx.2	SODIUM CHANNEL 2.		sp P78349 P78349			53%	300	758
HCCEU18	688041	83		blastx.2	UNNAMED PORTEIN PRODUCT.		sp Q9N083 Q9N083			98%	836	1003
HCEZF82	831745	84		blastx.2	(BC001698) Similar to lipase protein [Homo sapiens]		gb AAH01698.1 AAH01698			90%	401	490
HCFLN88	610000	86		blastx.2	BCL7B protein - human		pir S58284 S58284			69%	1417	1115
HCHAB84	834326	88		blastx.2	(AX061649) unnamed protein product [Homo sapiens]		emb CAC25009.1			95%	311	433
HCONSD29	862314	91		blastx.2	HUNTINGTIN-INTERACTING PROTEIN HYP/AFBP11 (FRAGMENT).		sp O75400 O75400			100%	658	714
HCRAY10	695709	96		blastx.2	HSPC244.		sp Q9P0N5 Q9P0N5			93%	433	480
HCRBF72	828945	97		blastx.2	MITOTIC SPINDLE ASSEMBLY CHECKPOINT PROTEIN MAD2B 1		sp Q9UI95 MD22_HUMAN			56%	1223	933
HCUCF89	637986	100		blastx.2	PRO2822.		sp Q9P147 Q9P147			49%	186	10
HCUCK44	790277	101		blastx.2	hypothetical protein DKFZp564J157.1 - human (fragment)		pir T34520 T34520			100%	17	604
HCWFU39	651316	104		blastx.2	CDNA FLJ20366 FIS, CLONE HEP18008.		sp Q9NX95 Q9NX95			95%	594	782
HDPDI72	897277	109		blastx.2	adult-specific brush border protein - rabbit		pir C45665 C45665			86%	278	475
HDPDJ58	587265	110		blastx.2	hypothetical protein DKFZp434D2328.1 - human		pir T42691 T42691			100%	82	744
										99%	337	1605
										40%	192	437
										100%	191	823
										84%	503	426
										100%	421	398
										99%	29	529
										100%	266	9
										84%	427	236
										67%	11	223
										76%	746	1450
										100%	307	627

09-08-2009

	(fragment)								
	100%							14	307
	87%							621	785
	33%							785	1393
	30%							656	1327
	38%							307	627
	35%							307	624
	36%							325	621
	30%							848	1402
	36%							259	624
	31%							854	1447
	33%							259	624
	96%							1137	1220
	37%							322	585
	37%							92	307
	30%							319	621
	37%							89	307
	31%							337	618
	26%							902	1402
	29%							307	615
	33%							307	618
	36%							101	313
	32%							86	307
	35%							83	307
	35%							92	286
	37%							134	307
	100%							1405	1464
	41%							624	779
	34%							89	316
	40%							827	1042
	40%							630	785
	26%							848	1144
	35%							89	274
	36%							101	307
	33%							131	307
	37%							125	307
	32%							340	594
	33%							307	540

HDPFF10	853513	111	HMMER 2.1.1	PFAM: Leucine Rich Repeat	PF00560	65.1	74	30%				316
							803	28%				1402
			WUblastx.6 4	garp [Homo sapiens]	emb CAA80847.1	38%	285	41%				624
							1153	29%				618
						37%	89	41%				1117
							624	27%				307
						50%	1229	33%				779
							1265	34%				1402
						50%	660	36%				1411
							645	36%				785
						50%	137	39%				773
							660	33%				307
						50%	633	36%				785
							188	33%				881
						50%	654	37%				316
							645	31%				785
						50%	645	35%				785
							624	35%				785
						50%	624	31%				794
							624	34%				776
						50%	89	42%				172
							454	35%				606
						50%	630	34%				785
							666	34%				785
						50%	543	31%				671
							553	34%				618
						50%	1134	45%				618
							1137	33%				1223
						50%	1125	35%				1220
							1149	28%				1220
						50%	1125	37%				1220
							1184	50%				1184
HDPFF10	853513	111	HMMER 2.1.1	PFAM: Leucine Rich Repeat	PF00560	65.1	729	65.1				800
			WUblastx.6 4	garp [Homo sapiens]	emb CAA80847.1	38%	285	38%				965
						42%	1153	42%				1593
						32%	1306	32%				1641
						29%	1147	29%				1446

HDPFU43	790189	112	blastx.2	PROTEIN-TYROSINE SULFOTRANSFERASE 2 (EC 2.8.2.20) 1	sp O60704 TPS2_HUMA N	31% 26% 30% 33% 30% 31% 33% 28% 27% 37%	1614 1159 1174 1306 1174 1162 1183 468 246 1629	1898 1512 1536 1632 1494 1539 1500 893 965 2111
HDPGE24	801947	114	blastx.2	NEURONAL THREAD PROTEIN AD7C-NTP.	sp O60448 O60448	56% 52% 55% 77% 52% 77% 62% 62% 53% 66% 39% 40% 57% 44% 29% 51% 50% 56% 51% 39% 53% 66% 48% 60%	1418 1413 1452 2528 2594 2525 2533 1858 1314 1813 1301 1457 2625 2242 2527 1857 1857 1364 1836 1305 2461 1800 2522 1380	1179 1237 1291 2448 2394 2433 2453 1763 1186 1742 1098 1248 2524 2078 2144 1723 1750 1290 1714 1177 2366 1747 2442 1321

100-443887-100

[illegible]

[illegible]

[illegible]

[illegible]



[illegible][illegible]

HDPPQ30	684292	120	blastx.2	CDNA FLJ20378 FIS, CLONE KAJA0536.	sp Q9NX85 Q9NX85	66%	549	701
HDPPW82	778405	121	blastx.2	hypothetical protein UL126 - human cytomegalovirus (strain AD169)	pir S09875 S09875	94%	6	155
HDQHM36	852328	123	blastx.2	UNNAMED PORTEIN PRODUCT.	sp Q9N083 Q9N083	66%	965	1153
HDTAU35	838139	124	blastx.2	NADH dehydrogenase (ubiquinone) (EC 1.6.5.3) chain 3 - human mitochondrion	pir A00422 DNHUN3	90%	11	340
HDTAV54	801898	125	blastx.2	(AL136938) hypothetical protein [Homo sapiens]	emb CAB66872.1	100%	13	303
HDTGW48	827285	127	blastx.2	SIRP-B2.	sp Q9P1W8 Q9P1W8	100%	783	1100

HE2CA60	888705	130		blastx.2		CISPLATIN RESISTANCE- ASSOCIATED OVEREXPRESSED PROTEIN.	sp Q9P2S7 Q9P2S7	81% 79%	1347 1658	1757 1744
HE2CM39	553651	132		blastx.2		SIMILAR TO RING-H2 FINGER PROTEIN RHA1A. CDNA FLJ10814 FIS, CLONE NT2RP4000984.	sp Q9SNH1 Q9SNH1	61%	565	449
HE2HC60	753265	133		blastx.2		CDNA FLJ10814 FIS, CLONE NT2RP4000984.	sp Q9NVC4 Q9NVC4	99%	125	1300
HE6CS65	762960	136		blastx.2		CDNA: FLJ21047 fis, clone CAS00253.	sp BAB14967 BAB14967	98%	938	1378
HE6DO92	562767	137		blastx.2		gag polyprotein - human endogenous virus S71	pir A46312 A46312	80% 64%	19 623	633 928
HE6EY13	847058	138		blastx.2		HYPOTHETICAL 28.3 KDA PROTEIN.	sp Q95476 Q95476	99%	5	490
HE6FU11	827236	139		HMMER 2.1.1		PFAM: von Willebrand factor type A domain	PF00092	184.7	244	771
				WUblastx.6 4		(AJ007581) matrilin-4 [Homo sapiens]	emb CAA07569.1	98% 38% 77% 45% 33% 47%	782 863 145 782 190 782	1642 1501 783 907 741 907
HE8FC45	843781	141		blastx.2		CDNA FLJ20378 FIS, CLONE KAIA0536.	sp Q9NX85 Q9NX85	72% 75% 46%	1845 1672 1312	1663 1553 1172
HE8FC45	845672	142		blastx.2		CDNA FLJ20378 FIS, CLONE KAIA0536.	sp Q9NX85 Q9NX85	72% 75% 46%	1845 1672 1312	1663 1553 1172
HE8FD92	856544	145		blastx.2		HYPOTHETICAL 105.9 KDA PROTEIN.	sp Q9UJ19 Q9UJ19	75% 85% 87% 84% 48%	2 2 389 2 2	1414 1102 1345 613 1267

HE8FD92	869847	146	blastx.2	HYPOTHETICAL 105.9 KDA PROTEIN.	sp Q9UIJ9 Q9UIJ9	48% 41% 39% 38%	167 47 605 101	1102 1075 1345 331
						95% 83% 45% 57% 56% 65% 40% 62% 62% 60% 50% 43%	4 4 1 1 1 31 4 331 331 331 316 40	609 540 462 297 297 270 462 540 540 540 540 258
HE8FD92	901142	147	blastx.2	HYPOTHETICAL 105.9 KDA PROTEIN.	sp Q9UIJ9 Q9UIJ9	94% 78% 48% 48% 62% 62% 60% 50% 58% 43% 41%	25 25 61 61 202 202 202 187 25 25 58	480 411 408 408 411 411 411 411 141 168 129
HE8SG96	862016	148	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	69% 66%	1860 2020	1696 1868
HE8TY46	899528	149	blastx.2	ZINC FINGER PROTEIN 289.	sp Q9JIH1 Q9JIH1	94%	318	938
HE9EA10	827796	151	blastx.2	laminin alpha-1 chain precursor - human	pir S14458 S14458	99% 28% 55%	761 878 736	1891 1840 789
HEBFR46	847064	157	blastx.2	CDNA FLJ20378 FIS, CLONE KAJA0536.	sp Q9NX85 Q9NX85	81% 80%	1304 1111	1110 1022
HEBGE07	798096	158	blastx.2	CDNA FLJ20378 FIS, CLONE KAJA0536.	sp Q9NX85 Q9NX85	81%	1863	1720

HELAT35	693175	160	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	69% 90%	2110 2111	1802 2079
HELBUS4	637624	161	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	59%	1255	1031
HEMEY47	834491	164	blastx.2	PRO2550.	sp AAG35515 AAG35515	63%	509	838
HEPBA14	855935	166	blastx.2	peptidylprolyl isomerase (EC 5.2.1.8) A - bovine	pir A01852 CSBOAB	92% 83% 81%	85 423 30	426 515 77
HEQAH80	701984	167	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	58%	1107	805
HEQBF89	786205	168	blastx.2	CDNA FLJ20378 FIS, CLONE KAlA0536.	sp Q9NX85 Q9NX85	61% 70% 70% 70% 70%	827 858 859 855 856	516 829 830 826 827
HETCI16	844543	169	blastx.2	BOG25.	sp Q9P0V3 Q9P0V3	99%	3	356
HETDW58	790557	170	blastx.2	(AF251296) TPA018 [Homo sapiens]	gb AAG44596.1 AF251296.1	100%	324	1058
HFCFE20	701985	175	blastx.2	EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT 10 1	sp Q14152 IF3A_HUMAN	89% 87% 44% 34% 29% 30%	438 1083 1071 1083 1083 1083	581 1205 1178 1205 1205 1172
HFEAY59	658685	176	blastx.2	C29.	sp Q9Z320 Q9Z320	79% 83% 39% 21%	50 403 486 608	1153 495 584 799
HFGAJ16	580824	177	blastx.2	CDM protein - human	pir S44279 S44279	97%	263	403
HFIJA29	839206	179	blastx.2	PRO1902 PROTEIN.	sp Q9UHT1 Q9UHT1	59% 46%	1026 889	880 806
HFIJA68	847074	180	blastx.2	SIX TRANSMEMBRANE EPITHELIAL ANTIGEN OF PROSTATE.	sp Q9UHE8 STEA_HUMAN	100%	13	399
HFKE505	827572	181	blastx.2	SREBP cleavage activating protein - Chinese hamster	pir T18526 T18526	85% 74%	241 9	1722 509

HFKEU12	634006	182	blastx.2	hypothetical protein 3 - rat	pir S21347 S21347	62%	387	692
						38%	756	1031
HFPDS07	821646	185	blastx.2	(AF327434) glutaminase [Homo sapiens]	gb AAG47842.1 AF32743 4_1	51%	757	933
HFVGK35	731868	189	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	53%	695	778
HFVHW43	570948	190	blastx.2	CDNA FLJ11786 fis, clone HEMBA1006036.	sp BAB13911 BAB13911	84%	2	436
						79%	343	528
HFXAV37	626595	191	blastx.2	NEURONAL THREAD PROTEIN AD7C-NTP.	sp O60448 O60448	60%	528	833
						67%	1223	1068
						59%	1453	1187
						64%	607	407
						69%	606	406
						52%	1472	1218
						69%	607	461
						58%	1454	1275
						68%	473	333
						56%	1295	1173
						37%	558	346
						45%	1342	1232
						43%	1408	1169
						47%	1366	1232
						42%	1412	1275
						78%	402	361
						60%	1240	1166
						31%	549	355
						75%	1404	1357
						43%	1454	1386
						40%	1467	1402
						59%	1287	1222
						52%	1229	1179
						47%	607	539
						34%	583	461
						60%	398	354
						46%	1187	1110
HFXBT66	580831	193	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	63%	535	807
						54%	809	913

HGBER72	826710	195	blastx.2	CDNA FLJ20489 FIS, CLONE KAT08285.	sp Q9NX17 Q9NX17	72%	1284	1063
HGBHP91	693011	198	blastx.2	PUTATIVE P150.	sp O00372 O00372	55%	1082	942
HGCAC19	851527	199	blastx.2	WUGSC:H_DJ0687K01.2 PROTEIN.	sp Q9UIE9 Q9UIE9	45%	537	52
HGCAC19	801999	200	blastx.2	CDNA: FLJ22454 fis, clone HRC09703 (Fragment).	sp BAB15362 BAB15362	52%	541	491
HGCAC19	842540	201	blastx.2	CDNA: FLJ22454 fis, clone HRC09703 (Fragment).	sp BAB15362 BAB15362	30%	94	5
HHEAK45	765278	202	blastx.2	DJ202121.1 (NOVEL PROTEIN) (CDNA FLJ11101 FIS, CLONE 1	sp Q9NPB0 Q9NPB0	100%	361	2712
HHEOW19	886174	204	blastx.2	RAB5 GDP/GTP EXCHANGE FACTOR HOMOLOGUE.	sp Q9UI41 Q9UI41	99%	184	1473
HHEFF87	778071	205	blastx.2	coatamer zeta chain - bovine	pir A49465 A49465	99%	182	1471
HHEFL34	753230	206	blastx.2	NEURONAL THREAD PROTEIN AD7C-NTP.	sp O60448 O60448	53%	890	1495
						64%	1049	1495
						81%	998	1495
						91%	1052	1525
						92%	1052	1540
						64%	956	1615
						100%	1052	1648
						99%	1949	1401
						89%	166	426
						81%	465	656
						91%	611	715
						92%	129	167
						64%	420	470
						100%	50	145
						43%	2597	1872
						71%	2631	2407
						66%	2048	1818
						63%	2029	1787
						63%	2631	2461
						58%	2464	2348
						50%	1963	1829
						44%	2631	2530
						41%	2541	2392
						66%	2410	2348
						60%	2475	2416

[illegible]



HKABU43	838573	227	blastx.2	CG6756 PROTEIN. (AB055298) hypothetical protein [Macaca fascicularis]	sp Q9VKC8 Q9VKC8	42%	95	1582
HKACI79	833361	228	blastx.2		dbj BAB21923.1	72%	886	1104
HKAFF50	790192	229	blastx.2	PRO1777.	sp Q9P1G7 Q9P1G7	93%	1753	1400
HKGBF25	738797	230	blastx.2	CDNA FLJ20489 FIS, CLONE KAT08285.	sp Q9NX17 Q9NX17	64%	1995	1630
HKMLK03	734213	232	blastx.2	UNNAMED PORTEIN PRODUCT.	sp Q9N083 Q9N083	65% 44%	856 1008	701 832
HKTAB41	695732	234	blastx.2	CDNA: FLJ22976 fis, clone KAT11222 (Fragment).	sp BAB15513 BAB15513	92% 92% 80% 77%	714 714 693 693	797 797 797 797
HLDQU79	740755	237	WUblastx.6 4	(AF064093) KE04p [Homo sapiens]	gb AAC26658.1	90%	105	1142
HLDQU79	837599	512	blastx.2	KE04P.	sp O75477 O75477	99%	81	1118
HLDRT09	830544	238	blastx.2	ATP-binding cassette half- transporter.	sp AAG33617 AAG33617	99%	2	469
HLHAP05	638476	239	blastx.2	CDNA FLJ20489 FIS, CLONE KAT08285.	sp Q9NX17 Q9NX17	72% 88% 55% 87% 87%	1833 1837 1585 1832 1831	1585 1811 1526 1809 1808
HLHCS23	560663	240	blastx.2	LINE-1 ELEMENT ORF2.	sp O62658 O62658	42%	1098	1412
HLIBO72	883431	241	blastx.2	(AF140225) unknown [Homo sapiens]	gb AAG48521.1	97%	32	547
HLICE88	840321	242	blastx.2	fibrinogen gamma-A chain precursor [validated] - human	pir A90470 FGHUG	100%	3	584
HLMBW89	701996	245	blastx.2	BETA-1,4 MANNOSYLTRANSFERASE	sp Q9P2Y2 Q9P2Y2	100%	10	90
HLMGP50	647603	246	blastx.2	Hypothetical 9.0 kDa protein.	sp BAB12374 BAB12374	72% 63%	935 765	807 706
HLQAS12	886180	249	blastx.2	LECTIN-LIKE OXIDIZED LDL RECEPTOR.	sp P78380 P78380	100% 75%	364 651	711 851
HLQCL64	864966	250	HMMER 2.1.1	PFAM: Major intrinsic protein	PF00230	87.3	87	449

HLQCX36	584786	251	blastx.2	aquaporin 9 - human PRO0478 PROTEIN.	pir JC5973 JC5973	98%	18	548
HLWDB73	838453	258	blastx.2	CDNA: FLJ21016 fis, clone CAE05735.	sp Q9UI59 Q9UI59	87%	1100	1216
HL YGB19	838083	261	blastx.2	(AL136703) hypothetical protein [Homo sapiens]	sp BAB14955 BAB14955	98%	1	657
HL YGY91	658703	263	blastx.2	CDNA FLJ13386 fis, clone PLACE1001104, weakly similar to 1	emb CAB66638.1	100%	660	872
HMC AZ04	839783	264	WUblastx.6 4	(AF042284) unknown [Homo sapiens]	sp BAB14578 BAB14578	97%	204	518
HMC AZ04	858210	265	blastx.2	HYPOTHETICAL 50.0 KDA PROTEIN.	gb AAD41160.1 AF04228 4_1	94%	221	391
HMC AZ04	867910	266	blastx.2	HYPOTHETICAL 50.0 KDA PROTEIN.	sp Q9Y6N5 Q9Y6N5	100%	106	1455
HMC AZ04	887445	267	blastx.2	HYPOTHETICAL 50.0 KDA PROTEIN.	sp Q9Y6N5 Q9Y6N5	100%	106	1455
HMC AZ04	668249	268	blastx.2	CGI-44 PROTEIN.	sp Q9Y6N5 Q9Y6N5	100%	107	1456
HMDAB29	584789	270	blastx.2	CDNA FLJ20489 FIS, CLONE KAT08285.	sp Q9UQM8 Q9UQM8	100%	9	1055
HMEBB82	783077	272	blastx.2	MITOCHONDRIAL ISOLEUCINE TRNA SYNTHETASE (FRAGMENT).	sp Q9NX17 Q9NX17	72%	1186	890
HMEDE24	837027	273	blastx.2	CG1837 PROTEIN.	sp Q9NSE4 Q9NSE4	99%	2	2206
HMED190	840077	274	blastx.2	Rab3 interacting protein variant 4 (Fragment).	sp Q9YVYV3 Q9YVYV3	46%	104	1159
HMELM75	587307	275	blastx.2	CDNA FLJ10468 FIS, CLONE NT2RP2000007.	sp AAG23165 AAG23165	38%	101	757
HMICP65	847403	279	blastx.2	Guanine nucleotide binding protein beta subunit 5L.	sp Q9NVW5 Q9NVW5	40%	794	1135
HMSBE04	709672	281	blastx.2	CDNA: FLJ22969 fis, clone KAT10759.	sp BAB15511 BAB15511	21%	2629	2823
						100%	81	794
						100%	137	391
						99%	8	892
						23%	269	943
						85%	182	3

HMSCL38	801919	282	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	55%	1155	1472
						71%	2841	2653
HMSCR69	843059	283	HMMER 2.1.1	PFAM: Zinc finger present in dystrophin, CBP/p300	PF00569	74%	2935	2855
			blastx.2	POTASSIUM CHANNEL MODULATORY FACTOR.	sp Q9P0J7 Q9P0J7	48.2	113	250
HMSHC86	840402	284	blastx.2	UNNAMED PORTEIN PRODUCT.	sp Q9N083 Q9N083	99%	107	1249
HMSHU20	847410	285	WUblastx.6 4	(AK025116) unnamed protein product [Homo sapiens]	dbj BAB15071.1	67%	1674	1420
			blastx.2	matrin 3 - rat	pir A40016 A40016	54%	1398	1234
HMTAB77	847411	287				70%	1724	1674
						47%	1722	1453
						87%	1024	2520
						100%	630	1055
						98%	242	628
						53%	2311	2760
						98%	3258	3428
						26%	2584	2763
						31%	2596	2709
						35%	1705	1797
						35%	3312	3404
HMUAE26	747403	288	blastx.2	SEVEN TRANSMEMBRANE DOMAIN ORPHAN RECEPTOR.	sp Q9P2R4 Q9P2R4	99%	577	1272
						100%	153	575
HMUAN45	833072	289	blastx.2	UNNAMED PORTEIN PRODUCT.	sp Q9N092 Q9N092	61%	239	1516
						70%	684	1238
						39%	346	1080
						30%	2022	2231
HMVBC31	825598	290	blastx.2	PRENYLCYSTEINE CARBOXYL METHYLTRANSFERASE.	sp O60725 O60725	100%	88	789
						80%	747	938
HMVDU15	801969	291	blastx.2	CGI-30 PROTEIN.	sp Q9Y319 Q9Y319	94%	75	917
HMWBL03	822861	292	blastx.2	hypothetical protein DKFZp762L0311.1 - human (fragment)	pir T50635 T50635	56%	617	1237
HMWJF53	758158	293	blastx.2	Nuclear LIM interactor-interacting factor.	sp AAG15402 AAG15402	100%	154	720
						91%	3	170

HNEAK81	722235	294	blastx.2	UNNAMED PORTEIN PRODUCT.	sp Q9N083 Q9N083	56%	770	1087
HNECL22	799541	295	blastx.2	HT015 PROTEIN.	sp Q9NYZ2 Q9NYZ2	100% 100% 26%	1756 2204 1762	2202 2359 2289
HNEHD88	815675	297	blastx.2	(AB055283) hypothetical protein [Macaca fascicularis]	dbj BAB21907.1	43% 64% 57% 40% 44% 53%	1667 58 9 53 9 59	1825 99 50 247 95 97
HNFAC50	815676	298	blastx.2	(AF308287) serologically defined breast cancer antigen NY-BR-20 [Homo sapiens]	gb AAG48255.1 AF30828 7_1	100%	425	282
HNFHF34	722237	300	blastx.2	(AF205218) NS1-binding protein-like protein [Homo sapiens]	gb AAG43485.1	100% 34% 35% 33% 32%	9 9 3 9 129	431 404 407 407 422
HNGAK51	603910	301	blastx.2	NEURONAL THREAD PROTEIN AD7C-NTP.	sp O60448 O60448	74% 66% 67% 45% 55% 43% 62% 62% 37%	714 702 733 631 530 542 629 626 527	914 878 915 903 631 709 715 706 751
HNGAM58	688114	302	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	65% 85% 73%	818 1081 1020	1018 1143 1076
HNGGP65	597449	310	blastx.2	Hypothetical 14.1 kDa protein.	sp BAB12154 BAB12154	47% 32% 100%	398 69 128	541 302 9
HNGJB41	852178	313	blastx.2	probable oxysterol-binding protein DJ430N08.1 - human (fragment)	pir T02435 T02435			
HNHFE71	834487	320	blastx.2	hypothetical protein	pir T47135 T47135	67%	822	583

HNHGK22	597451	321	blastx.2	DKFZp761L0812.1 - human (fragment)	sp P11369 POL2_MOUSE	48%	733	485
				RETROVIRUS-RELATED POL POLYPROTEIN [CONTAINS: REVERSE TRANSCRIPTASE (EC 2.7.7.49); ENDONUCLEASE].		37%	483	10
						33%	483	37
						55%	236	177
						36%	337	248
HNHBB10	634589	322	blastx.2	NEURONAL THREAD PROTEIN AD7C-NTP.	sp O60448 O60448	64%	894	691
						62%	888	706
						50%	854	561
						66%	719	594
						40%	887	711
						63%	109	14
						33%	885	586
						56%	106	14
						68%	718	653
						30%	130	5
						55%	639	586
						38%	868	794
						77%	80	54
HNTBT17	855957	324	blastx.2	CYCLIN L ANIA-6A.	sp Q9UK58 Q9UK58	59%	556	1440
						43%	743	1342
						78%	729	947
						41%	589	660
HOACG07	792928	328	blastx.2	CDNA FLJ14158 fis, clone NT2RM1001112.	sp BAB14854 BAB14854	99%	183	704
HODBV05	825283	331	blastx.2	94 KDA B-RAF PROTEIN (FRAGMENT).	sp Q13878 Q13878	100%	566	661
HODCZ32	836069	332	blastx.2	WD-REPEAT PROTEIN 9 (FRAGMENT).	sp Q9NSI6 WDR9_HUMAN	86%	8	331
HOEBK60	789396	333	blastx.2	CDNA FLJ13081 fis, clone NT2RP3002033.	sp BAB14427 BAB14427	98%	132	1916
						100%	14	109
						88%	106	159
HOFAA78	836646	334	blastx.2	CDNA FLJ20084 FIS, CLONE COL03526.	sp Q9NXS2 Q9NXS2	88%	29	529
						90%	529	792
						50%	9	80
HOFNB74	762821	335	blastx.2	CDNA FLJ20374 FIS, CLONE	sp Q9NX88 Q9NX88	44%	172	510

[illegible]

[illegible]

095008-091201

[illegible]



00000000000000000000000000000000

[illegible]

[illegible][illegible]

[illegible][illegible]

[illegible]

[illegible]

039008053601

[illegible]

**06-08-2017**

[illegible]

**096030**

[illegible]



095032-09101

[illegible]

**09-06-01**

[illegible]

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99

[illegible]



050909-02101

[illegible]

[illegible]

HPJBU43	862058	360	blastx.2	KAT05170.		sp Q9P1E1 Q9P1E1	54%	187	44
HPMCI84	562779	363	blastx.2	PRO2221.		sp AAG35515 AAG35515	65%	646	467
HPMCV30	612870	364	blastx.2	PRO2550.		sp Q99770 Q99770	56%	786	613
HPQAX38	843592	366	blastx.2	HYPOTHETICAL 15.4 KDA PROTEIN.		sp Q99770 Q99770	65%	614	384
HPQAX38	845752	367	blastx.2	CDNA FLJ20378 FIS, CLONE KAIA0536.		sp Q9NX85 Q9NX85	76%	384	334
HPRBH85	695752	370	blastx.2	CDNA FLJ20378 FIS, CLONE KAIA0536.		sp Q9NX85 Q9NX85	56%	885	607
HPRCA64	824074	371	blastx.2	CDNA FLJ20378 FIS, CLONE KAIA0536.		sp Q9NX85 Q9NX85	58%	605	513
HPRCD35	853551	372	blastx.2	Hypothetical 49.2 kDa protein.		sp BAB17282 BAB17282	54%	891	607
HPTRM02	812879	373	blastx.2	NCK-ASSOCIATED PROTEIN 1 (NAP 1) (P125NAP1) 1		sp Q9Y2A7 NCP1_HUM AN	58%	605	513
HRAAD30	866187	376	blastx.2	hypothetical protein DKFZp762L1710.1 - human (fragment)		pir T50629 T50629	97%	534	977
HRADA42	827302	377	blastx.2	SRC HOMOLOG 3 DOMAIN-CONTAINING PROTEIN HIP-55 (DREBRIN F).		sp Q9Y2A7 NCP1_HUM AN	63%	2	616
HRADF49	866481	378	blastx.2	CDNA: FLJ21839 fis, clone HEP01794.		pir T50629 T50629	64%	214	549
HRADN25	800628	379	blastx.2	hypothetical protein C11D2.4 - Caenorhabditis elegans		sp Q9Y2A7 NCP1_HUM AN	59%	935	1015
HRDAI17	560720	381	blastx.2	CDNA: FLJ22169 fis, clone HRC00632.		pir T50629 T50629	100%	1021	1926
HRDDQ39	840405	382	blastx.2	CDNA: FLJ21839 fis, clone HEP01794.		sp Q9Y2A7 NCP1_HUM AN	91%	387	1019
				hypothetical protein C11D2.4 - Caenorhabditis elegans		pir T50629 T50629	94%	11	481
				CDNA: FLJ22169 fis, clone HRC00632.		sp Q9Y2A7 NCP1_HUM AN	98%	296	613
				MYG1 homolog.		sp Q9Y2A7 NCP1_HUM AN	100%	2	250
				CDNA: FLJ21463 fis, clone COL04765.		sp Q9Y2A7 NCP1_HUM AN	99%	98	940
				CDNA FLJ20378 FIS, CLONE		sp Q9Y2A7 NCP1_HUM AN	35%	2	448
						sp Q9Y2A7 NCP1_HUM AN	99%	23	1393
						sp Q9Y2A7 NCP1_HUM AN	74%	668	931
						sp Q9Y2A7 NCP1_HUM AN	48%	387	668
						sp Q9Y2A7 NCP1_HUM AN	98%	13	825
						sp Q9Y2A7 NCP1_HUM AN	98%	813	1379
						sp Q9Y2A7 NCP1_HUM AN	77%	1291	1593
						sp Q9Y2A7 NCP1_HUM AN	34%	1590	1685
						sp Q9Y2A7 NCP1_HUM AN	99%	47	1174
						sp Q9Y2A7 NCP1_HUM AN	70%	1495	1334
						sp Q9Y2A7 NCP1_HUM AN	66%	775	578

HRDER22	688056	383	blastx.2	KAIA0536. CDNA FLJ10390 FIS, CLONE NT2RM4000104, MODERATELY SIMILAR TO 1	sp Q9NWX7 Q9NWX7	53%	582	436
HRDEX93	816046	384	blastx.2	PEFLN.	sp Q9UBV8 Q9UBV8	100%	13	864
HRDFK37	840381	385	blastx.2	UNNAMED PROTEIN PRODUCT.	sp Q9N032 Q9N032	57%	487	642
HRGBD54	828436	386	blastx.2	HPK/GCK-LIKE KINASE HGK.	sp O95819 O95819	78% 69% 87% 27%	379 253 32 6	2019 831 253 149
HSAVA08	580870	388	blastx.2	(AB055293) hypothetical protein [Macaca fascicularis]	dbj BAB21918.1	66% 46% 57% 63%	1059 941 949 796	934 792 896 764
HSAVW42	637660	389	blastx.2	SIMILAR TO RING-H2 FINGER PROTEIN RHA1A.	sp Q9SNH1 Q9SNH1	81% 73%	595 594	497 493
HSAWZ40	634000	391	blastx.2	ORF2-LIKE PROTEIN (FRAGMENT).	sp O00549 O00549	64% 64%	613 951	8 610
HSDZM54	637870	393	blastx.2	NADH dehydrogenase (ubiquinone) (EC 1.6.5.3) chain 3 - human mitochondrion	pir A00422 DNHUN3	86%	226	516
HSHBF76	715838	394	blastx.2	(AF194407) unknown [Homo sapiens]	gb AAG42221.1	66% 50% 72%	1267 762 834	836 460 748
HSJBY32	702020	396	blastx.2	(AJ278118) neuronal nicotinic acetylcholine alpha10 subunit [Homo sapiens]	emb CAC20435.1	96% 84%	215 466	514 684
HSKDR27	580874	397	blastx.2	HYDROXYPROLINE-RICH GLYCOPROTEIN (FRAGMENT).	sp Q39866 Q39866	33% 75% 46%	86 668 466	352 691 510
HSLHX15	777861	399	blastx.2	(AY012159) virion-associated nuclear-shuttling protein [Mus musculus]	gb AAG42155.1	100% 58%	180 487	263 522
HSNBM34	635131	402	blastx.2	acyl-CoA dehydrogenase (EC 1.3.99.-) very-long-chain	pir S54183 S54183	99% 100%	113 1548	1546 1979



HSOAH16	827038	403		blastx.2	specific - human CDNA FLJ20489 FIS, CLONE KAT08285.	sp Q9NX17 Q9NX17	35%	2053	2178
HSQDO85	853393	405		blastx.2	CG10161 PROTEIN.	sp Q9VCK0 Q9VCK0	64% 61% 56%	485 60 10	1021 521 57
HSQES57	831222	406		WUblastx.6 4	(AF151850) CGI-92 protein [Homo sapiens]	gb AAD34087.1 AF15185 0_1	93%	195	980
HSRBE06	871264	407		blastx.2	PRO2550.	sp AAG35515 AAG35515	70%	1626	1327
HSSDI26	560722	408		blastx.2	HYPOTHETICAL 15.4 KDA PROTEIN.	sp Q99770 Q99770	66%	1398	1264
HSSEA64	853395	409		blastx.2	Hypothetical 17.2 kDa protein.	sp AAG17210 AAG17210	98%	7	243
HSSEF77	658725	410		blastx.2	WW DOMAIN BINDING PROTEIN-1.	sp Q95637 Q95637	100% 59%	296 10	829 468
HSSFE38	742512	411		HMMER 2.1.1	PFAM: Ribonuclease HII	PF01351	76.3	184	-142
HSWBE76	751308	413		blastx.2	CDNA FLJ10375 FIS, CLONE NT2RM2001950.	sp Q9NW15 Q9NW15	100%	84	266
HSXCP38	895392	414		blastx.2	hydroxymethylglutaryl-CoA lyase (EC 4.1.3.4) - chicken	pir B45470 B45470	73%	17	895
HSYBI06	740766	415		blastx.2	(AB055298) hypothetical protein [Macaca fascicularis]	dbj BAB21923.1	78% 69%	821 916	913 954
HT4FV41	853400	418		blastx.2	collagen alpha 1(I) chain - chicken (tentative sequence) 1	pir A90458 CGCH1S	30%	1388	30
HT5GR59	801930	420		blastx.2	DOCKING PROTEIN.	sp O60496 O60496	73% 100%	70 687	1284 1031
HTAEI78	637684	421		blastx.2	EMDC II PROTEIN.	sp Q9Y3S0 Q9Y3S0	90%	85	174
HTDAA78	566861	422		blastx.2	(AL137800) dI127C7.3 (actin related protein 2/3 complex, subunit 5 (16 kD)) [Homo sapiens]	emb CAC19687.1	58%	84	302
HTEAG62	812332	423		blastx.2	HOST CELL FACTOR 2.	sp Q9Y5Z7 Q9Y5Z7	98%	14	2011

[illegible]

HTLEM16	779133	450	blastx.2	WW DOMAIN BINDING PROTEIN-2.	sp O95638 O95638	99%	50	841
HTLFA13	535937	452	blastx.2	CDNA FLJ20489 FIS, CLONE KAT08285.	sp Q9NX17 Q9NX17	56%	1159	839
HTLGI89	835069	454	blastx.2	AP47 protein - mouse	pir S19693 S19693	98%	675	1370
HTLFI11	843506	455	blastx.2	ORNITHINE DECARBOXYLASE-2.	sp Q9I8S4 Q9I8S4	98%	104	682
HTLFI12	834946	456	blastx.2	proline-rich protein PRB1/2S (EA) - human (fragment)	pir D40750 D40750	59%	353	1687
HTLFI12	842691	457	blastx.2	proline-rich protein PRB1/2S (EA) - human (fragment)	pir D40750 D40750	68%	309	356
HTLFI12	870167	458	blastx.2	proline-rich protein PRB1/2S (EA) - human (fragment)	pir D40750 D40750	52%	768	866
HTLFI12	886780	459	blastx.2	proline-rich protein PRB1/2S (EA) - human (fragment)	pir D40750 D40750	35%	129	323
HTLFI12	891533	460	blastx.2	proline-rich protein PRB1/2S (EA) - human (fragment)	pir D40750 D40750	52%	770	868
HTLFI12	901225	461	blastx.2	proline-rich protein PRB1/2S (EA) - human (fragment)	pir D40750 D40750	35%	131	325
HTNBK13	831967	463	blastx.2	proline-rich protein PRB1/2S (EA) - human (fragment)	pir D40750 D40750	52%	770	868
HTOAI50	638623	464	blastx.2	(AL136686) hypothetical protein [Homo sapiens]	emb CAB66621.1	35%	131	325
HTOAM11	664508	465	blastx.2	PRO1438.	sp Q9PIH3 Q9PIH3	100%	123	500
HTOEVI6	853616	468	blastx.2	CDNA: FLJ21463 fis, clone COL04765.	sp BAB15071 BAB15071	68%	1251	1138
HTOHO21	732808	470	blastx.2	1-ACYL-SN-GLYCEROL-3- PHOSPHATE	sp Q9NRZ5 PLCD_HUM AN	75%	603	433
HTOHQ05	853621	471	blastx.2	ACYLTRANSFERASE DELTA (EC 1.1.1.1)	sp Q9UII4 Q9UII4	68%	448	308
HTOJL95	806212	472	blastx.2	P47 LBC oncogene - human	sp Q15605 Q15605	99%	379	1164
HTOJL95	762851	473	blastx.2	CYCLIN-E BINDING PROTEIN 1.	sp Q15605 Q15605	98%	201	383
HTOJL95	762851	473	blastx.2	ORF1 CODES FOR A 40 KDA PRODUCT.	sp Q15605 Q15605		751	161
HTOJL95	762851	473	blastx.2	L1 ELEMENT L1.24 P40.	sp O00373 O00373	63%	192	61
HTOJL95	762851	473	blastx.2	L1 ELEMENT L1.24 P40.	sp O00373 O00373	86%	876	730
HTOJL95	762851	473	blastx.2	L1 ELEMENT L1.24 P40.	sp O00373 O00373	57%	607	248

HTPDU17	840596	474		blastx.2		CDNA FLJ10404 FIS, CLONE NT2RM4000486.	sp Q9NW00 Q9NW00	63% 32%	279 683	85 609
HTSFI32	637720	475		blastx.2		VESICLE ASSOCIATED MEMBRANE PROTEIN 2B.	sp Q9WUW2 Q9WUW2	94% 60% 36% 36%	553 1098 467 908	1308 1664 1444 1444
HTTCB60	853401	476		blastx.2		RNA polymerase III transcription initiation factor BRFU.	sp AAG30222 AAG30222	86% 75% 99%	448 690 6	627 803 881
HTTEE41	840950	477		blastx.2		T-COMPLEX PROTEIN 1, BETA SUBUNIT (TCP-1-BETA) (CCT-BETA).	sp P78371 TCPB_HUMAN	100%	92	1696
HTTEZ02	702027	478		blastx.2		(AL136812) hypothetical protein [Homo sapiens]	emb CAB66746.1	98% 73% 36% 48% 53%	6 278 54 54 78	314 391 185 128 116
HTWEH94	561680	479		blastx.2		UNNAMED PORTEIN PRODUCT.	sp Q9N083 Q9N083	55% 58%	912 856	1151 906
HTXDC38	801935	482		blastx.2		HSPC171.	sp Q9NZZ7 Q9NZZ7	78% 96%	100 437	486 514
HTXDC77	844258	483		HMMER 2.1.1		PFAM: Class I Histocompatibility antigen, domains alpha 1 and 2	PF00129	103.3	137	259
				blastx.2		HLA-B*5501 - human	pir I72752 I72752	89% 90% 90%	216 65 256	947 256 285
HTXFA72	853410	487		blastx.2		UNNAMED PORTEIN PRODUCT.	sp Q9N083 Q9N083	63% 59% 66%	1854 1688 75	1681 1557 1469
HTXMZ07	834881	490		blastx.2		hypothetical protein Sand - Fugu rubripes	pir T30808 T30808			
HUFCL31	801938	491		blastx.2		BG:DS01219.1 PROTEIN.	sp Q9V441 Q9V441	35%	601	1128
HUKBT67	844446	492		blastx.2		(AL136895) hypothetical protein [Homo sapiens]	emb CAB66829.1	100% 100% 32%	1040 8 80	1216 61 241

[illegible]

[illegible]

[illegible]

[illegible]



[illegible]

[illegible]

[illegible]

[illegible]

**SECRET**

[illegible]

054032-091201

[illegible]

[illegible]

[illegible]



[illegible]

[illegible]

[illegible]

[illegible]

[123] Table 2 further characterizes certain encoded polypeptides of the invention, by providing the results of comparisons to protein and protein family databases. The first column provides a unique clone identifier, "Clone ID NO:", corresponding to a cDNA clone disclosed in Table 1A and/or Table 1B. The second column provides the unique contig identifier, "Contig ID:" which allows correlation with the information in Table 1B. The third column provides the sequence identifier, "SEQ ID NO:", for the contig polynucleotide sequences. The fourth column provides the analysis method by which the homology/identity disclosed in the Table was determined. The fifth column provides a description of the PFAM/NR hit identified by each analysis. Column six provides the accession number of the PFAM/NR hit disclosed in the fifth column. Column seven, score/percent identity, provides a quality score or the percent identity, of the hit disclosed in column five. Comparisons were made between polypeptides encoded by polynucleotides of the invention and a non-redundant protein database (herein referred to as "NR"), or a database of protein families (herein referred to as "PFAM"), as described below.

[124] The NR database, which comprises the NBRF PIR database, the NCBI GenPept database, and the SIB SwissProt and TrEMBL databases, was made non-redundant using the computer program nrdb2 (Warren Gish, Washington University in Saint Louis). Each of the polynucleotides shown in Table 1B, column 3 (e.g., SEQ ID NO:X or the 'Query' sequence) was used to search against the NR database. The computer program BLASTX was used to compare a 6-frame translation of the Query sequence to the NR database (for information about the BLASTX algorithm please see Altshul et al., J. Mol. Biol. 215:403-410 (1990), and Gish and States, Nat. Genet. 3:266-272 (1993). A description of the sequence that is most similar to the Query sequence (the highest scoring 'Subject') is shown in column five of Table 2 and the database accession number for that sequence is provided in column six. The highest scoring 'Subject' is reported in Table 2 if (a) the estimated probability that the match occurred by chance alone is less than  $1.0e-07$ , and (b) the match was not to a known repetitive element. BLASTX returns alignments of short polypeptide segments of the Query and Subject sequences which share a high degree of similarity; these segments are known as High-Scoring Segment Pairs or HSPs. Table 2 reports the degree of similarity between the Query and the Subject for each HSP as a percent identity in Column 7. The percent identity is determined by dividing the number of exact matches between the two aligned sequences in the HSP, dividing by the number of Query amino acids in the HSP and multiplying by 100. The polynucleotides of SEQ ID

NO:X which encode the polypeptide sequence that generates an HSP are delineated by columns 8 and 9 of Table 2.

[125] The PFAM database, PFAM version 2.1, (Sonnhammer, Nucl. Acids Res., 26:320-322, 1998))consists of a series of multiple sequence alignments; one alignment for each protein family. Each multiple sequence alignment is converted into a probability model called a Hidden Markov Model, or HMM, that represents the position—specific variation among the sequences that make up the multiple sequence alignment (see, e.g., Durbin, et al., *Biological sequence analysis: probabilistic models of proteins and nucleic acids*, Cambridge University Press, 1998 for the theory of HMMs). The program HMMER version 1.8 (Sean Eddy, Washington University in Saint Louis) was used to compare the predicted protein sequence for each Query sequence (SEQ ID NO:Y in Table 1B) to each of the HMMs derived from PFAM version 2.1. A HMM derived from PFAM version 2.1 was said to be a significant match to a polypeptide of the invention if the score returned by HMMER 1.8 was greater than 0.8 times the HMMER 1.8 score obtained with the most distantly related known member of that protein family. The description of the PFAM family which shares a significant match with a polypeptide of the invention is listed in column 5 of Table 2, and the database accession number of the PFAM hit is provided in column 6. Column 7 provides the score returned by HMMER version 1.8 for the alignment. Columns 8 and 9 delineate the polynucleotides of SEQ ID NO:X which encode the polypeptide sequence which show a significant match to a PFAM protein family.

[126] As mentioned, columns 8 and 9 in Table 2, “NT From” and “NT To”, delineate the polynucleotides of “SEQ ID NO:X” that encode a polypeptide having a significant match to the PFAM/NR database as disclosed in the fifth column. In one embodiment, the invention provides a protein comprising, or alternatively consisting of, a polypeptide encoded by the polynucleotides of SEQ ID NO:X delineated in columns 8 and 9 of Table 2. Also provided are polynucleotides encoding such proteins, and the complementary strand thereto.

[127] The nucleotide sequence SEQ ID NO:X and the translated SEQ ID NO:Y are sufficiently accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, the nucleotide sequences of SEQ ID NO:X are useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the cDNA contained in Clone ID NO:Z. These probes will also hybridize to nucleic acid molecules in biological samples, thereby

enabling immediate applications in chromosome mapping, linkage analysis, tissue identification and/or typing, and a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used to generate antibodies which bind specifically to these polypeptides, or fragments thereof, and/or to the polypeptides encoded by the cDNA clones identified in, for example, Table 1A and/or 1B.

[128] Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

[129] Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the generated nucleotide sequence identified as SEQ ID NO:X, and a predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing cDNA Clone ID NO:Z (e.g., as set forth in columns 2 and 3 of Table 1A and/or as set forth, for example, in Table 1B, 6, and 7). The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with known methods. Further, techniques known in the art can be used to verify the nucleotide sequences of SEQ ID NO:X.

[130] The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.

#### ***RACE Protocol For Recovery of Full-Length Genes***

[131] Partial cDNA clones can be made full-length by utilizing the rapid amplification of cDNA ends (RACE) procedure described in Frohman, M.A., et al., Proc. Nat'l. Acad. Sci. USA, 85:8998-9002 (1988). A cDNA clone missing either the 5' or 3' end can be reconstructed to include the absent base pairs extending to the translational start or stop

codon, respectively. In some cases, cDNAs are missing the start codon of translation, therefor. The following briefly describes a modification of this original 5' RACE procedure. Poly A+ or total RNA is reverse transcribed with Superscript II (Gibco/BRL) and an antisense or complementary primer specific to the cDNA sequence. The primer is removed from the reaction with a Microcon Concentrator (Amicon). The first-strand cDNA is then tailed with dATP and terminal deoxynucleotide transferase (Gibco/BRL). Thus, an anchor sequence is produced which is needed for PCR amplification. The second strand is synthesized from the dA-tail in PCR buffer, Taq DNA polymerase (Perkin-Elmer Cetus), an oligo-dT primer containing three adjacent restriction sites (XhoI, Sall and ClaI) at the 5' end and a primer containing just these restriction sites. This double-stranded cDNA is PCR amplified for 40 cycles with the same primers as well as a nested cDNA-specific antisense primer. The PCR products are size-separated on an ethidium bromide-agarose gel and the region of gel containing cDNA products the predicted size of missing protein-coding DNA is removed. cDNA is purified from the agarose with the Magic PCR Prep kit (Promega), restriction digested with XhoI or Sall, and ligated to a plasmid such as pBluescript SKII (Stratagene) at XhoI and EcoRV sites. This DNA is transformed into bacteria and the plasmid clones sequenced to identify the correct protein-coding inserts. Correct 5' ends are confirmed by comparing this sequence with the putatively identified homologue and overlap with the partial cDNA clone. Similar methods known in the art and/or commercial kits are used to amplify and recover 3' ends.

[132] Several quality-controlled kits are commercially available for purchase. Similar reagents and methods to those above are supplied in kit form from Gibco/BRL for both 5' and 3' RACE for recovery of full length genes. A second kit is available from Clontech which is a modification of a related technique, SLIC (single-stranded ligation to single-stranded cDNA), developed by Dumas et al., *Nucleic Acids Res.*, 19:5227-32 (1991). The major differences in procedure are that the RNA is alkaline hydrolyzed after reverse transcription and RNA ligase is used to join a restriction site-containing anchor primer to the first-strand cDNA. This obviates the necessity for the dA-tailing reaction which results in a polyT stretch that is difficult to sequence past.

[133] An alternative to generating 5' or 3' cDNA from RNA is to use cDNA library double-stranded DNA. An asymmetric PCR-amplified antisense cDNA strand is synthesized with an antisense cDNA-specific primer and a plasmid-anchored primer. These primers are removed and a symmetric PCR reaction is performed with a nested cDNA-specific antisense primer and the plasmid-anchored primer.



***RNA Ligase Protocol For Generating The 5' or 3' End Sequences To Obtain Full Length Genes***

[134] Once a gene of interest is identified, several methods are available for the identification of the 5' or 3' portions of the gene which may not be present in the original cDNA plasmid. These methods include, but are not limited to, filter probing, clone enrichment using specific probes and protocols similar and identical to 5' and 3' RACE. While the full length gene may be present in the library and can be identified by probing, a useful method for generating the 5' or 3' end is to use the existing sequence information from the original cDNA to generate the missing information. A method similar to 5' RACE is available for generating the missing 5' end of a desired full-length gene. (This method was published by Fromont-Racine et al., Nucleic Acids Res., 21(7):1683-1684 (1993)). Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcript and a primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest, is used to PCR amplify the 5' portion of the desired full length gene which may then be sequenced and used to generate the full length gene. This method starts with total RNA isolated from the desired source, poly A RNA may be used but is not a prerequisite for this procedure. The RNA preparation may then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase if used is then inactivated and the RNA is treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase. This modified RNA preparation can then be used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction can then be used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the relevant gene.

[135] The present invention also relates to vectors or plasmids which include such DNA sequences, as well as the use of the DNA sequences. The material deposited with the ATCC (e.g., as described in columns 2 and 3 of Table 1A, and/or as set forth in Table 1B,

Table 6, or Table 7) is a mixture of cDNA clones derived from a variety of human tissue and cloned in either a plasmid vector or a phage vector, as described, for example, in Table 1A and Table 7. These deposits are referred to as "the deposits" herein. The tissues from which some of the clones were derived are listed in Table 7, and the vector in which the corresponding cDNA is contained is also indicated in Table 7. The deposited material includes cDNA clones corresponding to SEQ ID NO:X described, for example, in Table 1A and/or 1B (Clone ID NO:Z). A clone which is isolatable from the ATCC Deposits by use of a sequence listed as SEQ ID NO:X, may include the entire coding region of a human gene or in other cases such clone may include a substantial portion of the coding region of a human gene. Furthermore, although the sequence listing may in some instances list only a portion of the DNA sequence in a clone included in the ATCC Deposits, it is well within the ability of one skilled in the art to sequence the DNA included in a clone contained in the ATCC Deposits by use of a sequence (or portion thereof) described in, for example Tables 1A and/or 1B or 2, by procedures hereinafter further described, and others apparent to those skilled in the art.

[136] Also provided in Table 1A and 7 is the name of the vector which contains the cDNA clone. Each vector is routinely used in the art. The following additional information is provided for convenience.

[137] Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., *Nucleic Acids Res.* 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., *Nucleic Acids Res.* 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., *Strategies* 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Phagemid pBS may be excised from the Lambda Zap and Uni-Zap XR vectors, and phagemid pBK may be excised from the Zap Express vector. Both phagemids may be transformed into *E. coli* strain XL-1 Blue, also available from Stratagene.

[138] Vectors pSport1, pCMVSPORT 1.0, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, also available from Life Technologies. See, for instance, Gruber, C. E., et al., *Focus* 15:59- (1993). Vector lafmid BA (Bento Soares, Columbia University, New York, NY) contains an ampicillin resistance gene and can be transformed into *E. coli*

strain XL-1 Blue. Vector pCR<sup>®</sup>2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, available from Life Technologies. See, for instance, Clark, J. M., *Nuc. Acids Res.* 16:9677-9686 (1988) and Mead, D. *et al.*, *Bio/Technology* 9: (1991).

[139] The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, and/or the deposited clone (Clone ID NO:Z). The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include preparing probes or primers from the disclosed sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

[140] Also provided in the present invention are allelic variants, orthologs, and/or species homologs. Procedures known in the art can be used to obtain full-length genes, allelic variants, splice variants, full-length coding portions, orthologs, and/or species homologs of genes corresponding to SEQ ID NO:X or the complement thereof, polypeptides encoded by genes corresponding to SEQ ID NO:X or the complement thereof, and/or the cDNA contained in Clone ID NO:Z, using information from the sequences disclosed herein or the clones deposited with the ATCC. For example, allelic variants and/or species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for allelic variants and/or the desired homologue.

[141] The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

[142] The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below). It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification, such as multiple histidine residues, or an additional sequence for stability during recombinant production.

[143] The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially purified using

techniques described herein or otherwise known in the art, such as, for example, by the one-step method described in Smith and Johnson, Gene 67:31-40 (1988). Polypeptides of the invention also can be purified from natural, synthetic or recombinant sources using techniques described herein or otherwise known in the art, such as, for example, antibodies of the invention raised against the polypeptides of the present invention in methods which are well known in the art.

**[144]** The present invention provides a polynucleotide comprising, or alternatively consisting of, the nucleic acid sequence of SEQ ID NO:X, and/or the cDNA sequence contained in Clone ID NO:Z. The present invention also provides a polypeptide comprising, or alternatively, consisting of, the polypeptide sequence of SEQ ID NO:Y, a polypeptide encoded by SEQ ID NO:X or a complement thereof, a polypeptide encoded by the cDNA contained in Clone ID NO:Z, and/or the polypeptide sequence encoded by a nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1C. Polynucleotides encoding a polypeptide comprising, or alternatively consisting of, the polypeptide sequence of SEQ ID NO:Y, a polypeptide encoded by SEQ ID NO:X, a polypeptide encoded by the cDNA contained in Clone ID NO:Z, and/or a polypeptide sequence encoded by a nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1C are also encompassed by the invention. The present invention further encompasses a polynucleotide comprising, or alternatively consisting of, the complement of the nucleic acid sequence of SEQ ID NO:X, a nucleic acid sequence encoding a polypeptide encoded by the complement of the nucleic acid sequence of SEQ ID NO:X, and/or the cDNA contained in Clone ID NO:Z.

**[145]** Moreover, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in Table 1C column 6, or any combination thereof. Additional, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the complementary strand(s) of the sequences delineated in Table 1C column 6, or any combination thereof. In further embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in Table 1C, column 6, and have a nucleic acid sequence which is different from that of the BAC fragment having the sequence disclosed in SEQ ID NO:B (see Table 1C, column 5). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in Table 1C, column 6, and have a nucleic



**[147]** Further, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in column 6 of Table 1C which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1C, column 2), or any combination thereof. Additional, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the complementary strand(s) of the sequences delineated in column 6 of Table 1C which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1C, column 2), or any combination thereof. In further embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in column 6 of Table 1C which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1C, column 2) and have a nucleic acid sequence which is different from that of the BAC fragment having the sequence disclosed in SEQ ID NO:B (see Table 1C, column 5). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in column 6 of Table 1C which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1C, column 2) and have a nucleic acid sequence which is different from that published for the BAC clone identified as BAC ID NO:A (see Table 1C, column 4). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in column 6 of Table 1C which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1C, column 2) and have a nucleic acid sequence which is different from that contained in the BAC clone identified as BAC ID NO:A (See Table 1C, column 4). Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides and polypeptides are also encompassed by the invention.

**[148]** Moreover, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in the same row of Table 1C column 6, or any combination thereof. Additional, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the complementary strand(s) of the sequences delineated in the same row of Table 1C column 6, or any combination thereof. In preferred embodiments, the polynucleotides of

**[149]** In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in column 6 of Table 1C, and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1C, column 2) or fragments or variants thereof. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

**[150]** In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in column 6 of Table 1C which correspond to the same Clone ID NO:Z (see Table 1C, column 1), and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1A, 1B, or 1C) or fragments or variants thereof. In preferred embodiments, the delineated sequence(s) and polynucleotide sequence of SEQ ID NO:X correspond to the same Clone ID NO:Z. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

[151] In further specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in the same row of column 6 of Table 1C, and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1A, 1B, or 1C) or fragments or variants thereof. In preferred embodiments, the delineated sequence(s) and polynucleotide sequence of SEQ ID NO:X correspond to the same row of column 6 of Table 1C. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

[152] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1C and the 5' 10 polynucleotides of the sequence of SEQ ID NO:X are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[153] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1C and the 5' 10 polynucleotides of a fragment or variant of the sequence of SEQ ID NO:X are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[154] In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of the sequence of SEQ ID NO:X and the 5' 10 polynucleotides of the sequence of one of the



sequences delineated in column 6 of Table 1C are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

**[155]** In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of a fragment or variant of the sequence of SEQ ID NO:X and the 5' 10 polynucleotides of the sequence of one of the sequences delineated in column 6 of Table 1C are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides, are also encompassed by the invention.

**[156]** In further specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1C and the 5' 10 polynucleotides of another sequence in column 6 are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

**[157]** In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1C and the 5' 10 polynucleotides of

another sequence in column 6 corresponding to the same Clone ID NO:Z (see Table 1C, column 1) are directly contiguous. Nucleic acids which hybridize to the complement of these 20 lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[158] In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of one sequence in column 6 corresponding to the same contig sequence identifier SEQ ID NO:X (see Table 1C, column 2) are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[159] In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1C and the 5' 10 polynucleotides of another sequence in column 6 corresponding to the same row are directly contiguous. In preferred embodiments, the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1C is directly contiguous with the 5' 10 polynucleotides of the next sequential exon delineated in Table 1C, column 6. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[160] Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. Accordingly, for each contig sequence (SEQ ID NO:X) listed in the fifth column of Table 1A and/or the fourth column of Table 1B, preferably excluded are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 and the final nucleotide minus 15 of SEQ ID NO:X, b is an integer of 15 to the final nucleotide of SEQ ID NO:X, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:X, and where b is greater than or equal to a + 14. More specifically, preferably excluded are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a and b are integers as defined in columns 4 and 5, respectively, of Table 3. In specific embodiments, the polynucleotides of the invention do not consist of at least one, two, three, four, five, ten, or more of the specific polynucleotide sequences referenced by the Genbank Accession No. as disclosed in column 6 of Table 3 (including for example, published sequence in connection with a particular BAC clone). In further embodiments, preferably excluded from the invention are the specific polynucleotide sequence(s) contained in the clones corresponding to at least one, two, three, four, five, ten, or more of the available material having the accession numbers identified in the sixth column of this Table (including for example, the actual sequence contained in an identified BAC clone). In no way is this listing meant to encompass all of the sequences which may be excluded by the general formula, it is just a representative example. All references available through these accessions are hereby incorporated by reference in their entirety.

**TABLE 3**

Clone ID	SEQ ID NO: X	Contig ID:	EST Disclaimer		Accession #'s
			Range of a	Range of b	
H6BSF56	11	762968	1 - 591	15 - 605	AW958287, BF027085, AV650800, AV650218, BF689895, BE409727, BE871017, BE278963, BF975253, AA449214, AA150070, BF247445, AA310756, BF337859, AA425098, BE746295, BE732859, BE742068, R48107, BF129114, AA393871, AI707816, AI523073, AW002940, BE672910, BF764476, AW827130, AA468022, AA493695, AW857950, AW275510, AW857971, BF667587, AW021735, R52299, AW965008,

				<p> AI223604, AI254279, BE179557, BG059450,  AW963750, AI445674, AW979031,  AV703942, AV762535, AI687343, BG249643,  AA769402, AW827120, AA484373,  AI345157, AV739452, AW168618,  AW504900, AA467876, BF887977,  AV710066, AV763354, AV762098, AI744826,  BF964993, AA279421, AW302903,  AW872575, AI700109, BF437493, AV764329,  BE253048, AW270343, AL046205,  BE782280, BF677892, AV759437, AV734583,  AV760777, AV760486, BF965007, BE907585,  AV764578, AW131249, BE297262,  BF347740, BF337291, BF679274, AW193265,  AI247199, BF347791, AV764307, AV763183,  AW235497, AA747070, BF760796,  AW872676, AI004704, AW002350,  AI270117, AI311927, BF871137, BE883107,  AL043009, AI754658, AI250083, AV760258,  AW069769, AI370094, AL119691,  AW063143, AI270559, AA372481,  AV760937, AL119713, AW857949,  BF742624, AA720702, BE736829, BF681649,  AI953275, AA490183, AF330238,  AW970871, AU145314, BF977376,  AL138265, AV759172, AV761106,  AV735614, AW953071, AA019312,  AA584167, AV728425, AU121243,  AV763847, AL038799, AV733830,  BF965154, BG026806, AI133164, AA523841,  AV763540, AV762050, AI470646, AI284640,  AI307022, AA635739, AI350211, BE350772,  AI691091, AI251082, AI370074, BF936005,  AI305766, AI732378, AI860013, AV744393,  AW974109, AW500125, AV760378,  AV734666, AV764241, AL037683,  AL038705, AA683238, BG023888,  AK025830, AF151821, AB015724,  AC004760, AC005089, AC005988,  AL049766, AC005257, AL117377, AL109936,  AC009311, AC007383, AC018637,  AL161445, AL034545, L78833, AC005250,  AC006511, AL136223, Z95115, AC006999,  AC005606, AL022322, AC007011,  AC007279, AC007428, AC002476,  AL049795, AC016576, Z98051, AC011508,  AL021393, AC006285, AC005923,  AL137839, AC003101, AL121934,  AC020893, AC004638, AP001753,  AC007620, AC010524, AF215937, AL117332,  AL022163, AC008482, AC006077,  AC022432, AC011559, AL121586,  AC004849, AC019215, AC005071,  AL022238, AL034423, AC006275, AL354720,  AC002314, Z82198, AL137780, AC005694,  AC010422, AF252830, AC004686, AL121751,  AC004814, AC024084, AC005280,  AC007192, AC002430, AL157938, AP001680,  AL031777, AL353748, AL021807, AL136131,  AC002395, AL136969, AL020997, </p>
--	--	--	--	---

					AC005258, AC018633, AL118501, AC018720, AL135818, AC004008, AC006111, AL035462, AC002470, AC016025, AC007384, AC018712, AL136308, AL035667, AC004876, AC005020, AC002425, AL109823, AL034420, AC007216, AC004650, AC007225, AL078611, AF001549, Z69706, AP001725, AL049544, AL121892, U63630, AL499629, AC010271, AL008725, AL021939, AL445687, AF254822, AC002531, AL121601, AL022237, AC004388, AC011310, AL031311, AC005527, AL050349, Z68162, AL121903, AC007546, AC020917, AL139186, Z93241, AC015853, AC020934, AF117829, Z82190, AC022150, AL360227, AL157931, AC004980, AL121897, AC004622, Z83844, AC005288, AX039602, AP001687, AL031286, AC022027, AL117258, AC005821, AC004899, AL391839, AL356414, AP001746, Z97196, AL109921, AL031662, AC004598, AC015555, Z83845, AC008372, AL080242, AL137061, AC012309, AC004659, AC004755, AJ229043, AL137853, AC008770, AF228703, AC011477, AC020916, AC008521, AL021578, M63796, AL009181, AC006312, AC024561, AC005488, AC005274, AC006130, AP001721, AC021752, AL008712, AC004534, AC023105, AC006277, AL133282, AC005529, U63312, AC016395, AC005484, AC005531, AL033520, AC001231, AC004894, AL136137, AC073976, Z98742, AL035079, AF047825, AC002301, Z98752, AC008976, AL359695, AL035424, AC018644, U95742, Z99716, AL031650, AC010609, AC013436, AC011531, AC006501, AC006017, Z98750, AL133289, AL135901, AF317635, AL009183, AL138976, AL023284, AP000962, AF042090, AC000066, AC009481, AC022517, AC007450, AC005600, AL121952, AC007262, AL117382, AL365335, AC008886, AC020906, AF243527, AL049537, AC008088, AP000501, AC006451, AL353807, AP000555, AC008403, AC009277, AF108083, AC005081, AL078590, AC005778, and AL450226.
H6EDM64	12	841331	1 - 2596	15 - 2610	AL529288, AL514648, AL523579, AL523918, AL530571, AL528848, AL523917, AL523578, BE795355, BE614208, AL529287, BE797988, BE747962, BE798201, AL530750, BF689293, BE884814, BF508994, BE798313, BE613450, BE787266, AW131835, AL530749, BG248495, BE386285, BF526775, BE873469, BE299650, AL042569, BE621187, BG168950, AW410458, BE883794, BE869375, BF348689, AW239351, BE737181, BE734276, BF309636, BF129214, BG180549, AW410610, AW601905, BE621858,

					AA689552, BF310547, AW960649, BF953086, AL045821, BE882424, BF724804, BE019151, AW246108, BG179779, AW374338, AW675186, BE279317, BG011956, AI475847, AI394166, AI142042, AW068652, AI539419, AI970048, AI792316, AA536006, AW272491, BG012645, AI827847, BG254459, AI673493, AW007399, AI719374, AA994188, BG176564, AI707847, AW104963, AI220974, AA022523, BF807054, BG012634, BF803094, N24911, AW665019, AI458806, AA689495, AA480131, AI808412, N41812, W17347, BE772562, BG012642, BF807055, BE772573, BG011957, BG012641, AL514647, F22287, AI160580, AI149344, BE772556, AI870582, BE772568, AW801577, BG176616, AW801325, AW068651, AI197831, BE265961, AA483525, BE772566, BE772574, BE693737, AA687509, BE839398, BF799200, AA687451, AI201450, BF896481, BE772569, BE244158, BE772576, BE826728, AI452812, BE772561, AA317941, AA308425, AA745895, AW751437, BG256219, AA782657, BF373198, AA364848, AL039960, AA405870, AW963550, BE300303, N78953, AA112404, BE826586, AI061434, AI143698, AW087863, AI382254, AW731818, BE788591, AW304748, AI589259, AA357514, BF663656, AW673017, AW664622, AA524482, AW246627, BE831243, BE831271, BG055766, AI749023, AA380438, BF746714, BE839346, AW084279, AA113160, BF529848, AI160508, BF764174, BF752908, AA053148, AW842671, F32117, AI190107, BF752929, BE547478, AA977756, AA360528, AA022454, BF808843, BF813892, AI917965, BG011699, BG012316, BF373193, BG122581, AA622680, BF688484, BE772558, AA053706, AA733114, BG012318, AW880294, AA482098, BE256450, BE831281, BF765811, BF803085, BE243388, AA774840, AA576098, BE831236, BE772816, AF024631, AX011724, AX015345, AF096303, U73627, AF061779, AC004923, AF238378, and AC000385.
H6EEC72	13	889401	1 - 1479	15 - 1493	BF034355, BF034892, BE792423, BF338898, BG105853, BE390915, BE613966, AA449897, BE389478, AW857371, AW861388, BE891738, R71843, BF983885, BF739366, AI688525, BF591064, AI589048, AI933344, BE387873, AI660119, AI950422, BF830644, AA250941, W68171, BE613313, BE389218, AA699649, BE612723, AI553767, BG178871, BE966158, AW965656, AI807258, AW606086, W67712, N34048, AA789094, AI160489, AA953906, AA029513, AI798377, AA961141, AI191879, AI277742, BF757878, AI341511, BF941471, T79588, W39291, BF843992, BF761673, BE552032, AW938641, AI684229, AI829091, AI696662, H79702,

					AI803066, AI423727, AW081674, AW014236, AW582288, BE675078, AI760447, BE042621, AA250965, AI991516, AW438983, AW205754, AI658602, AW594379, AA449841, BF926493, AI955308, AI917867, BF063286, AA029448, AI933496, AA350855, R71793, AW578255, BE829073, BE828899, AB014591, AL133647, AF180474, and AF211967.
HACAB68	14	584773	1 - 1286	15 - 1300	BF967733, BF340072, AW058572, BE877116, BF029667, BE221318, BE042897, BF434234, BE966145, BF593609, AW966641, BE549675, AI692588, BF433926, W68167, AW674743, W67708, BG163487, AI802057, AW051536, AW005086, BE073104, AU145008, AI634647, AI743810, N51396, BE218196, AI857811, AI816124, AI802067, AI095027, BE503637, BF669349, AI925492, BE669954, AI813855, AI811403, BG236435, AA833834, BE073105, AA748470, AW975666, BE502705, N56917, AI146547, AI949209, AI492350, AI190896, BE219670, AI167132, AW013890, AI089941, AI810922, AI804940, AI689151, BF699838, AW873589, AA209320, N62725, AI420094, AI221693, BF130415, AI301467, AA808217, AW511885, BE073003, AW166094, AA019916, AI359094, BE073109, AI753256, AW675323, BF671156, AA258518, AA954483, AA324329, BF668455, F13496, AA281446, BF247796, AA487161, BF029971, AA730575, AA121642, AI123192, R49582, AI887042, AA487312, AA364288, AA385769, AW440846, R60975, AW451535, AA972339, AI091153, T74984, R36295, AW118180, R75731, BE003024, AI459209, BG055090, AW895451, H80344, AI984894, AA581815, BF877111, AW805837, BE000523, D57701, T03076, AI767454, F10499, Z40296, R43580, AA081798, R49916, BE928534, Z43703, AA493265, AA526871, AU118452, AI144481, BE540542, AL442081, AR079029, AL354793, AK001029, AF189009, AB015344, AF177346, and AF293385.
HACBJ56	15	847112	1 - 874	15 - 888	AA157001, BE348653, AW027639, AA534339, AW001883, AA363258, AW959379, T71037, AW953765, BE048583, BF878388, T67200, AW393348, AW393350, AW384705, AW386713, AA156760, AW055343, BF892732, and BE140594.
HACBS22	16	847113	1 - 3225	15 - 3239	AU124073, BG253611, AW965191, BG115564, BF032820, AW965203, BF306517, BF304908, BF306593, BF982389, BF033269, AW890326, BF983369, AI302793, BF591498, AI567467, AV711792, AA429582, AI422294, AI859789, AA433987, AA814217, AA252998, AA279301, AA411385, AI342153, BE328862, BF127455, BF223404, BE073310, AW131093, AI061457, AW886431, BF316718, AI418930, AI126382, BF334564,

					BF115589, BF229901, AI199632, AI083697, AI076499, AI565545, AI083759, AW168955, AI075711, AA451848, AI832008, AA278862, AA406270, AA039587, AA603155, AI654833, AI217940, T67815, AA004762, BG056988, AW166397, W48835, AA857111, AA099992, AI283878, AA156539, AI268898, AA311784, AA625982, AW081237, AI431333, AW337580, AA157697, AI142910, AA434158, AW953535, AA281247, AI886152, AA099991, AI222819, AA279346, AA464193, AW902181, H41843, AA292181, T41156, T41125, AA918848, AV725329, AW237010, R20286, AA864669, H53657, BG008052, R22827, AA366602, AA456430, AA323466, AA897397, AA399222, BG002413, AA278539, R15382, BF899835, BE746477, R73996, AW663090, F03040, BG005938, R23319, BG002290, F06777, AA431990, AA884057, AI696614, BG002306, AA743065, AW805802, T40225, AA004763, H41803, T26456, R49943, H78738, AA598528, H78725, BE242201, AW769702, H79499, AW804514, H79512, AW589498, R08207, AA446922, BG008055, R42123, AI040559, H53618, BE938610, R23242, BF943906, T41091, W49816, BG007599, BF746451, T40268, BF943900, BE242322, T26457, BG007600, AA779288, AA620918, AA345630, T67965, BF943907, BF230057, T40261, AL120752, R43626, AA159388, H69584, AA252997, BE247168, R73903, R46015, AW189448, BE241991, AI868154, AI953925, AI421607, BF869086, BE939781, AA894421, AA039586, AA324932, AI498022, AW965271, AW300921, AW965468, BF871875, W68610, BE242008, BF829222, AA333777, R08258, AA400456, BE769384, AA332944, BE267667, AI147721, R49942, BG121728, W42651, BF243188, BG170063, AA815037, BE257390, BE790941, BF808078, AW977805, AW291305, T81704, AB011083, AF033861, AX015348, M55075, AX015524, AF087963, and AF253540.
HADDE71	17	839187	1 - 653	15 - 667	BE646364, BE562975, BE734905, AA227916, BE275558, BE387443, AI568587, BE387535, AW245842, BE857544, AI805978, BE386863, AA530975, AA845548, BF437434, AW627607, BE741623, BE898827, AA393921, AI201926, AI391625, AI199262, AI675180, AI123847, AA463396, AI128152, AI197839, BE791237, BE384118, AA913172, AA505110, AW408817, AI187762, AI076304, BE899199, R52594, AW009600, AA465034, AA913634, AA488109, AA885156, AA452881, AA464960, AA884143, BF340639, AI886462, AI188491, BF446332, BF221728, AA227574, R84997, BE243531, BE903843, AA424231, BF331414, AI675231, BE249813, D45528, R40380, W60397,



				BE670322, F37062, R53393, AI473277, R84954, BE265829, AW407603, W60306, AA322573, AI928674, AI368380, AL535519, AI312011, AI886883, AI471250, BF032040, AA336279, AI357522, F31267, BE910005, BE311908, BE293522, AV681951, AI349772, BE964812, BG108147, BE047859, AW827203, BF054789, AV682330, AL513597, AL514803, AV682809, BG168696, AI868831, AW268253, AV682441, AL047042, AV682266, BF724691, BE047863, BF795712, BG058208, AV711509, AW071349, BE048071, AL514627, AL513907, AL513803, AV758592, AV723772, AL515041, BE613622, AV758110, AV710479, AV762488, BF673434, AV704928, AI349645, AV682249, AI815383, AV755581, AL135661, AV695052, BF968041, BE905408, BG033403, AV733397, AV682521, AV723204, BG179993, AI684265, AV756770, AI349614, AV682051, AV682772, AL119049, AV681668, AV682289, AV706777, AL514791, BG250190, AI207510, BF726322, AL514473, BE785905, BE880190, AV655645, BE881155, AV681630, AV682252, AL121270, AV723062, AL045500, AL513643, AV758668, AI906328, AV758217, AV681857, AV757012, BG108324, AV682082, AL120854, AV682479, AV681872, AL514935, AW080838, AV682074, BG105099, AV682466, AV729890, AV682222, BF981774, BG259801, AI349598, BF343172, AL515047, AL514155, BE048319, AV755613, AW467961, AV682672, AL515373, AI500553, AI907070, BG254754, AI064830, BE964700, AV733385, BF340104, BE777769, BE964486, AI687376, AW166645, AV682697, AV756703, AV682476, AV726951, AV756477, AV682351, BG109125, BG036846, BG259943, AL513631, AL515173, AI909666, AL514261, BG110283, AI340582, BE783707, AI436456, BF971016, BF969662, BE891101, BG178809, AV758806, AV755614, AV708119, AV757096, AL036396, AV681858, BE877769, BE018711, AW162071, BG114104, AV732941, AV723953, BE906959, AV710608, AV705644, AV733326, AV734318, BE048065, BF883916, AV681586, AL513763, AL514691, BG257535, AV704350, AV758738, AV729334, BF339420, AI580190, AI149592, AI345111, AI624859, AV682496, AL514919, BF344705, BE967113, Y11587, I48979, AF116644, AF116602, AF078844, AF116639, AR079032, AF125949, AF130105, AL389978, S78214, AF113691, AF116691, L31396, L31397, AL133640, AF090900, AF130059,
--	--	--	--	--

					<p>AF130075, AX046603, AF118064, AF116646, AF090934, AL442072, AL050393, AL049938, AF118070, AL157431, AB048953, AF090943, AF116631, A93016, AF113013, AL133016, AF130104, AF138861, AL110196, AF130082, AL442082, AF090901, AL050146, AL137527, A08916, AL117457, AB041801, AJ242859, AK026608, AL133606, AL117460, AB050510, AF113694, AF218014, AL122050, AF104032, AF090903, AK000212, AL080060, AL390167, AL049452, AB049758, I89947, AF119878, AF113676, AL359596, AL110221, AL359601, S68736, AF111847, AF116688, AF113690, AF106862, AK026865, AL162006, AF130092, AB047615, I89931, AF090896, AR059958, AB048964, Y16645, U42766, AL049430, AX019230, AF113689, AB019565, AL050149, AL049466, AK026741, AF119875, AL050116, AF113677, AX019229, AK025084, AL122093, AL162083, AK025339, AL050108, X84990, A08913, AL096744, AL049314, AL133075, AL133557, AK025958, AK026045, AL133258, AL133080, AF219137, AK026855, AL050277, AF113019, AF091512, I48978, AL122123, AL080137, AF017152, AR011880, AF113699, AX006092, AL080124, AC007375, AL133093, AK026784, AL137283, AL389982, AF097996, AL133565, E03348, AK026744, AF130099, AF116649, AL137459, X63574, AF119899, AL137557, AK000618, AF158248, Y11254, AL122121, AJ000937, U91329, AF111851, E07361, AL050138, AC007390, AF146568, AK026542, AF119871, AK025772, AK026533, AL117394, AK027096, AF314091, AF091084, AK000614, AC004690, AL359618, AF207829, AK000445, AL110225, E05822, AK000137, AK025092, AF125948, AC022215, AK026452, AC004093, AK000083, AL137550, AF271350, AL359941, AB047904, X82434, AK026353, AC002467, AF119909, AB048954, U00763, AF079765, AK000652, AB048974, AL117585, AF242189, AF177401, AC000111, AL353802, AK026592, E07108, AL359615, AF130087, AC006371, AB051158, AR087170, AC006336, AL133560, AC004686, AK026647, AK026927, AF017437, AK026583, AL117583, X70685, A65341, AK024538, AF067728, AX042059, AK026480, AF177336, AL353940, AP001699, AF061943, AB052191, E02349, AK025491, AC006944, AL049382, AL022147, AC005886, AK026528, AL049300, AK027113, AL049464, AL078630, and AK026504.</p>
HADDJ13	18	827273	1 - 2304	15 - 2318	<p>AW575129, AW022897, AA010299, AU144131, AA669573, F11929, AL138228, AA634252, T66105, AA219059, R91924,</p>

					<p>R51726, AA856981, AA782322, AC004552, Z95118, AF224669, AC018797, AC003035, AC008636, AC009600, AC002565, AC002350, AL390738, AC073593, AC004491, AC011497, AF205588, AC007546, AC005250, AC016594, AC006038, AL031120, Z84469, AC018719, AC006080, AL391259, AL162615, AL034372, AC018639, AL132777, AC004605, AC024168, AL080275, AC004685, AL136000, Z82198, AL121578, AC013734, AL133286, AC002091, AL356575, AC007934, AC005035, AL031311, AC005840, AL117337, AL160313, AL035400, AC007225, AC005208, AC002368, AC011484, AC018637, AL035555, AL136231, AP001714, AC006211, AP000512, AP001732, AC083861, AC087095, AB023051, AC078833, AC005670, AL163267, AL080315, AC022201, AL138721, AC005529, AL136300, AL391114, AC006285, AL079342, AL031293, AC011310, U82671, U52111, AC018812, AL136365, AL356379, AC007358, AC000085, AC007151, AL445669, AL079340, AC007363, Z94056, AC006195, AL139109, AC002312, U85195, AL022164, AC005884, AC008009, AL109964, AL139092, AC010326, AC003950, AC002351, AE000658, AC009484, AL031659, AL138836, AC009247, AC004674, AC007386, AP001731, AL138832, AC007228, AL355530, AP001671, AC015853, AC073607, AL137230, AL031295, AL122125, AC018641, AL137818, AC055740, AC012599, AC015971, AP001730, AC004554, AC083874, AL035089, U91322, AL133500, AL122020, AC007011, AP001713, AL162831, AC007907, AC003029, AL033529, AF015262, AC005822, AC034242, AC008518, AC005902, AC004675, AL096862, AC005703, AL356299, AL121747, AC005722, AC020629, AC012384, AP001728, AL109939, AC000120, AJ229043, AF088219, AL023876, AC006116, AL137141, AC007057, AL158828, AL137800, AL391122, AL035454, AC006064, AC005046, AL008715, AC004253, and AP000030.</p>
HADMB15	19	847116	1 - 316	15 - 330	<p>AW136268, BG056888, AI131328, AI174443, AI091646, AW117296, AW168872, AI082447, AI432175, AI290911, AI741489, AI682685, AI142536, BG059892, AW149659, AW071935, AA233541, AI183690, BG056462, AI689641, AA599916, BF196591, BF196843, AA199743, AW136277, N77910, AA564806, AA243035, AA779709, AV722133, AI032138, AA844525, AI467910, AW965361, AA852418, AI982751, AI282445, AI982761, T03902, AI420648, AW167499, H08108, BE328548, AW068986, C15651,</p>

					D52660, AW665899, AI246702, AI538705, AI271662, AI435112, AI288692, BE466948, AI690048, D55112, AA779042, AL536118, D53747, D54101, AA486941, D53384, W07076, AA232504, AA486765, BF832290, AI038647, AW497637, BF947006, AU155428, and T05461.
HAGBQ12	20	722205	1 - 729	15 - 743	AI332690, AI374724, AI285345, AA876359, AA987498, AI702600, AI079453, AI382918, C04098, R63800, AI697895, H87363, R67068, BE673734, R73892, BF432849, R68633, R68632, R66112, H03322, AA340294, H87907, AP000350, AC007363, AC003969, AC005948, AP000360, AC004066, AC000053, AC002060, AL450169, AL033522, AP001696, AL050309, Y17293, AL138696, and AC004776.
HAGDW20	21	637489	1 - 1270	15 - 1284	AI671549, AA603387, AA614197, AC005629, AC006453, AC020898, AP001610, AC025435, and AC083863.
HAGEG10	22	823543	1 - 5670	15 - 5684	AU117844, AU118506, AU119632, AU142701, AU118316, AU119959, AU119025, BE875471, AU126076, AU137238, BG104575, AU138967, AU119281, BE880144, BE620212, BF527450, AL046250, BG029297, AU117066, AU126818, BG113544, AU128669, AW604129, BG179797, BF346364, BE694058, BE079543, AI872395, AW292851, AW170564, BE079454, AW812853, AV714888, AL042857, BG028602, BG005799, BE614073, AA523866, AW390450, BE856764, AW393917, BE676397, BF375747, AW373523, BG168417, BE874209, BG110102, AW393931, AW393924, BF340623, BE348284, AL048803, AW440289, BE896149, AW603939, AW590802, W72259, AA131497, N37081, BF880848, AA074901, AI126725, BF197419, AV720399, BE185077, AW753218, AU145052, AW935198, W58475, BE185075, BE185029, BF132321, AL048441, BE464666, AW968607, BE544412, AI872418, AU143989, AA188783, BF984545, AW166006, AU150921, AU153212, BG059577, BF001763, AI923075, AI188147, AI687738, AA429541, W58350, BF576448, AI191432, BF434307, AA687272, AW967782, D82378, AA486891, AA074726, AA770549, AI352455, BE677700, AU146166, T62894, AA019898, AI682014, AI141110, AI984576, BE091476, AU159948, AA167708, AI803035, BE220983, AI954034, AW473381, AW170305, AI888001, AV750571, AA514490, AI149197, AW022225, W95447, BF222723, AI160170, AW390442, W88638, AI159801, AA019911, W94007, AA918797, AI149526, N20405, AW022401, BF593100, AI872122, AA129509, AI123869, AI829442, AW340679, BE932213, AW665371,

					BF353920, AI470206, AA471281, AA831456, BF448445, AA923369, AL046249, AI015859, AW444790, AI811146, BE165922, AW272390, AI825915, AA847170, AI393175, AA621242, AI190196, AU144907, AA053669, AA262264, AA663506, AI206842, AW073367, BG236039, BF743168, AI221801, AA548471, AA767426, N31919, AA013224, BE146085, W79591, AW361597, AW361657, W30927, AI090566, AW393576, AW993026, AW390443, BE614509, AW170516, AW069441, BE966031, AW293184, AL121271, AA612903, AA991198, AA525491, AA995766, AA428587, AI619772, BF844737, AA046189, AW103132, BF111589, D82485, AA811517, AA262875, AI989394, AA252802, AA760763, AA486695, AA768156, AA188773, AA664378, AA834574, BG117833, W79810, AI131383, N36423, W76476, H62712, T65953, BF699718, AA149019, AI204502, AW602174, AA205998, W42709, AI880545, AI862197, AU145459, AA826167, AW992357, AA166945, L44407, T65958, W69500, AA022549, AA308301, AA487122, AW362039, AA771935, AA811100, BF851506, AA476695, AI887553, AA136559, AW339567, W02787, D82500, AA486801, W69501, W92367, AW264084, BF361101, BF001501, AA865320, AI347497, AI274491, D50525, AK000749, AK000625, AK025129, AC004542, AF088006, X98066, AC008897, AF067728, AX042059, AF111849, T62744, T64529, T64535, T92713, T92790, R00209, R05403, R05404, R06985, R06986, R18149, R41696, R41696, R63842, R64456, R66071, R67671, H04551, H53471, H53846, H62797, H86336, H86337, N42704, N44298, N46397, N48830, W42902, W88556, W93952, W92368, W95398, AA013319, AA022548, AA046083, AA053771, AA070849, AA079889, AA079890, AA131379, AA166764, AA243403, AA251232, and AA252615.
HAGEQ79	23	828055	1 - 771	15 - 785	AI741487, AA779582, BE674646, AW303577, AI305251, BE219521, AI688718, AI936253, AI093754, AW341275, BE222507, AI692909, BF966664, AI493111, BF525487, AW016639, W92767, AU150022, AW341787, AI278427, BF966817, AA044775, AA910036, AI685015, AI285959, AA719683, BE645673, AW196910, AI432636, AI096735, BE618873, BE541159, H91757, AI702190, BG152855, AI244929, AI681847, BE217959, AI498036, BE467879, BE696146, AI810609, R55798, AA897359, BE464034, AA975324, F09971, W73069, AW594097, AA703815, BF224038, F10760, T72606, AA932659, H46138, AA351671, T31206, AA317283, AI867144, AI681277, AV718692, BE938093, AV718489,

					<p> T31188, AI874073, AV724520, D80253,  D80219, AW582752, D80240, D80210,  AV719783, D80134, D59275, AV742001,  D80391, D51423, AV720464, AV718844,  AV720731, D80227, D51250, AV701332,  AV699550, D59619, D80193, AV722801,  AV699447, AV720220, AW973447,  AV742667, AI306486, AV743601, AV745080,  AV701443, AV744934, D80949, AV701017,  AV742720, AI305252, AV741012, AV701043,  AV701248, AV700229, AV743008, D80196,  D59927, AV701428, AV719000, AV701004,  AV745831, AV744773, D80043, AV701125,  AV701154, AV719913, AV699927,  AV701431, AV721784, AV700889,  AV744771, D80366, M79008, D59787,  AV744768, D81026, AV723927, AV701261,  AV720203, AV745724, AV720812,  AV740535, AV745723, C14227, AV720607,  C14014, AV701012, AV701149, AV701118,  AV737584, AV745366, AV701163,  AV745369, BE702445, AV723097,  AV701166, BE702442, AV744770,  AV701422, D50995, AV701344, AV699479,  AV699669, AV719822, T11051, AV699866,  AV701055, BF224150, AV701227, C75259,  D80045, D80168, AV699746, AV746385,  AV719324, AV701145, AV718707,  AV745488, AV701021, AV701330,  AV746335, AV701121, AV742671,  AV700357, D80022, D80195, AV700622,  AV720211, D58283, AV745392, D59889,  C15076, D81030, AV701335, AV746162,  D80188, D59467, AV745920, AW959570,  D80038, AV701130, F13647, D51799,  AV701419, AV718770, AV745847,  AV701013, D80378, AV745388, T03269,  AV721386, AV745490, AV718800,  AV701415, AV718681, AW904616,  AV700495, AW949642, AV701151,  AV719468, AV738934, C14429, AW978634,  AV745853, AV745197, AW949645,  AV701231, AV701357, AW965158,  AW975618, C14298, AW949633, AW949632,  AW949641, D80212, AV699682, AV745621,  AW949629, AA285331, AI557751, T11417,  AW966531, AV720150, AV701224,  AW949653, AW949658, D50979, AW949656,  AW966062, D59502, AK022817, AL096677,  AX027925, Y17188, A67220, AX021518,  AX020190, X68127, D34614, A25909,  AB012117, AR025207, AX047062, A85396,  AR074141, A85477, AR066482, AR088705,  A86792, AX042372, A44171, AB037923,  AX026824, I19525, A62298, AX020191,  AX033851, AR092424, D26022, A62300,  I18371, AX047063, AR008430, AR062871,  A84916, AR037157, A84772, A84774,  A84776, A84773, A84775, AR062872,  AR017907, AR062873, A20702, AR079804, </p>
--	--	--	--	--	---

10950000 09100

					AR091537, A43189, A43188, AX006821, AR067731, A20700, AR093385, AR067732, A58522, A91750, AX035462, AX026823, AX006825, AX006826, AR070327, AX006822, X67155, AX001082, A80951, AX047064, AJ302649, AF156304, AJ132110, AR018138, D89785, AF260572, A78862, AJ294956, S78798, AX035434, X82626, D88547, S69292, X92518, X93549, AF058696, AR087649, AF156302, AR077702, AR008278, AF156303, AF135125, AB028859, A82595, AX015396, and AX028130.
HAGFS57	24	847120	1 - 860	15 - 874	BF893958, AL079477, BE221875, AL532698, AI299412, R51649, AL040440, AA339493, F12505, F05649, Z43527, F06606, R12847, BF690787, R25251, T74335, AW382934, and AB020663.
HAGHN57	25	773286	1 - 2426	15 - 2440	AL533248, AU118622, AU119331, AU133909, AU119469, AU118182, BE794468, BE791529, BG176702, BE280450, BE729801, BF663566, BF970116, BE257176, BG032912, AL516224, BG121097, BE784191, BG249033, BE727671, BE881192, BE745390, BF792305, BF037862, AV710149, BE617085, AV751361, AW291174, BG163346, AI686123, BG033409, AV762315, AV704873, BE540243, BF344980, AV707943, BF671351, BE394881, AW070780, BE538770, BF303671, BE541947, AW963773, BF303913, AW299817, BE378370, AW299807, BF107096, AW515893, AI338838, BE254836, AW402330, AA455894, AI436127, AL516223, BF001973, AI392820, W31025, W28207, BE535313, BE258523, BF109189, AA182513, BE617702, AW275883, AW674662, BG169977, BE711218, AA134574, AW304388, AA588768, BE868534, AU144819, AA455892, BF802948, BF222585, AW902162, H16095, AI034153, AU145137, AI905391, AI985354, BG011776, AW612879, BE711276, AV659416, AU150558, BE702340, BF055535, BE711244, AA652292, AW271981, AA780056, AI624858, AA319693, AA604113, AV744893, AW771218, AV742941, AA837954, T60588, AA150957, AA151047, AI991761, AI912891, AI628783, AI434787, AW072744, AA716130, BF807693, AA181782, AI554969, AA916968, AA101864, AI473865, AA362607, AW338509, AI525459, BE244147, AI928082, AI433249, BF062859, AI910904, AA285264, BE711295, AI354885, AW006732, AI950274, AU144122, AI990867, AI922170, AA115829, AA806393, BE672240, AU156842, BE243206, AI633602, W01852, BE711219, AI280611, AA707161, AA301320, BF197637, AI695111, AW966603, BF447153, F29695, BE378061, AA336840, AI424341, AA385049, AI307649, N58884, AA131117, AI205138, BF431130, BF807685, N98771, AA602492,

0950082-091201

					BE711204, BF438567, F34557, AA748737, T60437, AA745028, AW891490, AW673414, AI630237, AW378199, AW779341, BE172988, BE172375, AA101187, AA781579, AI478435, BE699167, R57333, AI927982, R92570, BE764834, BF818234, AA648053, BE464290, AK000994, AR034165, AF054182, L12965, D13907, AC004668, AL050216, and AA227675.
HAHEA15	26	847013	1 - 1332	15 - 1346	BF683891, AW977366, AI743651, AW295805, AU144609, AI469869, AI188995, AA236904, AU117919, AI080089, N92489, BG116971, AW956038, W17171, AA872336, H96165, AA236057, BE538565, H96164, AI824681, AA736871, BF089224, R23025, BE934268, AA370017, AW891311, AI356478, AA693355, AI924051, AA766932, AA669094, AI568061, AI348901, AW020619, AL036368, AI436438, AW021717, AL049053, AV681572, BF344734, AK000965, AF072546, AK026843, AF139986, AK027210, AF126247, X98066, AB052176, AL137258, AF116617, AK024855, U62807, AF126372, AL122104, AJ132433, AL050172, A40111, AB048975, and AL133029.
HAJAA47	27	534670	1 - 1223	15 - 1237	BF991208, BF743765, AW021917, T74524, T57767, AI491765, N22058, AA904275, AA228349, AI689019, AA054085, AU131834, BE256101, AW270771, AL119691, AI284543, AU118852, BE062478, AI859946, BF769528, AW873261, AW152178, AC009318, AC011811, AL023799, AL137796, AP000704, AL499628, AC007934, AC005082, AC006111, AP001711, U91323, AC002407, AL031680, AL356244, AC008526, AL132987, AP000103, AL049540, AC013434, AP000269, AK024933, AP000212, AL133211, AC008924, AL035422, AP000280, AC018719, AC005200, AC005000, AC004858, AL133163, AF045555, AC009756, AC007546, Z98044, AP000107, AC008267, AC005520, Z98050, AL121933, AC002994, AL133174, U47924, AP000031, AP000354, AL162430, AL021154, AC011449, AP000039, AC009600, AC000025, AC004526, AP000355, AL356057, AL137798, AC012085, AC004383, AC004998, AL049713, AC004253, AP000065, AP000134, AC008521, AC004477, AC022173, AL031432, AP001727, AC012351, AC011442, AC005920, AC010271, AC010636, AL009183, AL109797, AL022237, AC024078, AC004232, AC007371, AC011470, AC008753, AC005484, AF109907, AC009155, AC004882, AL109827, AC011452, AL121891, AL109804, AC011465, U63721, AC008738, Z81364, AL138878, AL050308, AL117380, AC002487, AL161659, AC008764, AC005480, AC005841, AC003070, AL022163, AJ224877, Z93017, AC005220, AC004821, AC005755,



					AC005944, AF157623, AJ012824, AL353701, AL135783, AL122020, AL133264, AC007366, AP001714, AL352979, U62293, AC003959, AC008403, AC004968, AC011475, AC008747, AC005409, AC027644, AC020916, AL139353, AP001752, AL160397, AL022312, AC005231, AC020904, AC018663, AC010170, AL137073, AC005288, AL049872, AL133405, AC008249, AC002389, AC002492, AL009179, AC007057, AL138707, AL049775, AL135744, AC007345, AL034402, AL160175, AJ277546, AC020552, AC011742, AL353777, AC008805, AC002301, AC010458, AP001719, Z98200, AC007163, AP000167, AP000052, Y15994, AC002430, AL021940, Z98752, AL080242, AP001715, AL121753, AC007384, AL136532, AC007050, AL109743, AC008569, AP001694, AL160162, AC002551, AL161936, AL033521, AC006312, AL390374, AC007917, AF001548, AL353807, AL109952, AC011451, AL022320, AL109798, AC003665, AC005839, AC006088, AC005821, Z97056, Z95115, AC074331, AC008736, AP000556, AL031295, AL109759, AL133231, U62317, AL121897, AL163279, AC007226, AC009516, AF168787, AC005399, AL133353, AC005057, AC009247, AF275948, AL009181, AL022165, AC007707, AC006454, AL357497, and AC006121.
HAJAY92	28	845601	1 - 2331	15 - 2345	AI208943.
HAJBV67	29	866415	1 - 2522	15 - 2536	BG252656, BF732416, AV713753, BE905485, BF062374, BF445098, BF110352, BG252894, BE620095, BG249923, BE867752, AW606977, BG171028, AW576585, BE868698, BF671587, AW860769, BF941584, BF986308, AW305358, BF037687, BE541890, AW958924, AW974216, BF105260, AL048954, BF434917, AA057428, AW860733, BF664978, AI040432, BF984881, BF114918, BE872774, BE349491, AW263003, BF697715, BF382321, BE938703, AI378631, BF447674, AA446149, AA044378, BG114831, BF815345, BF085497, BF815237, BF210190, AA579908, BF132467, AA437015, AW860753, AI741531, AI742016, AI963805, AV748930, AA457625, BF815346, N31845, AI927889, BF699623, AA587067, AA831367, AI038411, AA442844, AI382172, BF084350, AW993684, AW407667, BF029928, AW028681, BE327066, BF887305, AV695738, BE222425, AV696527, BF755168, BE876090, BE167030, AI768063, BE000825, H12700, AV708152, AW001069, H03274, BF063098, BE933732, BF815719, BF594797, AW974217, N93209, N23944, AI290752, BF802746, AA557778, AA604449, Z32781, BE004621, AA910221, AA226865, R78864, BF326913, BG179582, AI370350, BE719765,

				<p> BE768063, BE932712, AA780882, T31498,  AW798498, AI635435, BF088211, BF817478,  Z28655, BF943308, Z24930, BE932705,  BF126152, AI015125, BF981166, AI684725,  T36185, H12701, R37535, AW952059,  AI689130, AA296931, AW798657,  AW364905, R79351, H03275, BE768230,  AW206046, AA081583, AA936681,  BG104571, BE176285, AW993023, W69607,  R31681, AI479514, BE696398, AA716370,  BE463676, AW366456, BE869217,  BF064127, BF001446, AW884802,  AW999085, BG104993, AI039088, R31723,  AW366514, BE871677, AI241206, AI743907,  AA306185, BF037794, D61175, AA852523,  AW365573, BF799275, AA129989,  BF985004, AW838470, BF802748, R36687,  BE001097, AA723997, BE932064,  AW366145, BF230069, AW999007,  AW993306, BF733961, AA508532,  AW972441, AW972636, BE932056,  BE086739, AA164808, BE064535, BF986296,  BF095055, AW408116, BE184804,  BE184805, BE184738, BE695142, BE184803,  BE172976, BE184743, BF984676, BE184732,  BF741954, AI366900, AW082623,  AW118518, AI698391, BF871314, AI954504,  BF753053, AI679312, BE967260, BF207979,  AL515195, AW050850, AW089844,  AW151136, AL515191, BE965599, AI619607,  AI687568, AI540674, AI345688, AL513817,  AI590043, BG032036, AA806028, AL043168,  AA329665, AI923989, AI670002, AA641818,  AI866770, AI679321, AI591420, AI473451,  AI445165, AW051088, BF812961, AL514093,  AI521560, AI633125, AL514871, AI915291,  AW152182, AI247082, AI582932, AI889189,  AI587121, BE875959, AA743012, BF814412,  AW193894, AL515413, BF911554, AI918449,  AF269150, AK000756, A91337, AF116347,  AF269151, A78983, AF160213, A78984,  A78985, AF124819, AF269149, E12747,  I89947, AB048975, I32738, AL137478,  AF287051, AF113694, AL133560, AK027129,  U42766, AJ005690, AL080118, AR068753,  X66871, AL355719, AK026927, I68732,  AB047878, AF113690, AK025857, I48978,  X63162, AL137529, AK000323, AF267849,  AF119899, AF111849, M92439, AL122100,  AL353956, AF130110, AL137557, AF132676,  AL133640, AF061836, E06743, AL137533,  AK027164, AF130055, AL049430,  AK027173, S82852, S83440, AF176651,  AL389935, AL137560, L04504, AF031147,  AR079032, AL137555, AK026649, A08907,  AK000206, AR038854, AK026057,  AF177401, AL110225, AL133623, AF116631,  X81464, AF078844, AB016226, AB050407,  AF118094, AL049938, A77033, A77035,  A08913, AF200464, AF090903, I09499, </p>
--	--	--	--	---

					AL050138, AL137550, D83032, Z97214, S75997, AK027113, AK024588, AL137258, AL390184, A08912, AK000310, AL096744, Y16645, AL389939, AF028823, AL110196, AB033884, AF079763, AK025435, AB046642, AB050431, AK024944, AK000414, AF119843, AB052191, AL117435, AL110218, AF116682, AF130059, AK026534, S36676, AL049283, A76337, AF087943, AF069506, AF141289, AF130099, AK025113, AJ010277, S76508, A08910, AK000160, U72621, AK026857, A08909, AK027096, X99257, AR029490, AL137271, AL137459, S77771, AF036268, AF079765, AL157433, AF260436, AK024622, AB049852, A08908, AK027082, AK026749, AB044547, AL110224, AK027111, AF119875, AL157482, AL122104, AB050410, A90844, AF113699, AF111851, A47357, AL080140, AL442082, AL137488, AF056191, AK025465, AL122050, AL133606, AL133559, AC008250, A18777, L13297, AK027116, AK026547, AK027121, AX014095, AF232009, AK025312, AL133016, AL080074, AL359620, AL050172, A17115, A18079, A18788, AK026462, E01573, E02319, AL117635, S83456, AK027114, AL050277, AR075044, AK026959, AK000083, AK027160, AK000618, AF026124, AF130082, AL133568, Y07905, AL050393, AF227198, AK000653, A08911, AL049347, AX015915, I00734, U75932, AL162002, AL137480, AL162079, A41575, Y11254, AL390154, AL110296, E00617, E00717, E00778, AL080148, AL122121, AL133112, AK026593, AF130104, AB049892, AL122110, AK026542, AF100781, AK024992, AK026894, X83544, Z72491, AK026541, AF183393, AB051158, AF106697, AK000421, AK000257, AF090900, AK027188, AL023657, I03321, A12297, AL137292, D16301, I48979, AF077051, I89931, X63410, AL133637, AL049324, S78453, AF113677, AF116614, E04233, AF116639, AJ299431, and A76335.
HAJCH70	30	827275	1 - 2168	15 - 2182	AL031294, AP001101, AC007677, AP001759, AC006038, AC009430, AL139824, AC007344, AC005939, AC007225, AL009177, AC002538, AF108083, AL031686, AC005486, AL109622, and AL139343.
HAOAG15	31	852204	1 - 5129	15 - 5143	BE349027, AA460958, AA460959, AI291926, AA461266, N58870, N72734, AI277327, AA461265, H43656, BF927468, H44722, BE835623, BE835624, Z33537, AI903814, BF366858, AW968357, AA465480, AA485068, AA430219, AF112345, AF074015, and AF172723.
HAQAI92	32	688037	1 - 593	15 - 607	AL522436, AL524148, AA513002, AI735602, AA772397, AW014080, AI799589, AI818675, BE617237, AA478326, AI217776,

					AW409592, BE646171, AI005409, AA552147, AW177019, H42123, R00846, H42122, BE795741, AA362478, AI491808, AI769438, AI560335, BE613582, AI336126, AI371050, R01499, AI610208, BF089287, BF095098, AW613379, AI275309, BF752321, and AL118502.
HAQCE11	33	633730	1 - 582	15 - 596	N45328, H29603, F10900, AW949645, D80045, AW965158, AW949642, AV738340, AV742732, AV724520, AW949643, AV741220, AW964468, AW966389, AV718489, AW966330, AV699550, D58283, AV718692, AW975618, AV742048, AW978634, D81030, AV718844, AW959570, AV720203, D80043, D59619, D80195, D80210, D80240, D51423, D51799, D80391, D80253, AV719822, D80227, AV719324, AV719783, AV718800, AW966531, D80188, AV720211, AV720464, AV718770, AV720731, D80219, AV699447, AV722801, D80196, AV723927, AV699927, C14429, D59927, D80038, D80212, D80193, D80022, AV719468, AW949632, F13647, AW949641, D80366, D59889, AV700889, AV720812, AV721386, AV723097, AW973447, AW949656, C15076, AW949629, AW966062, D59275, AW966053, AW964532, AV701335, AV742001, AV701043, AV701332, AV701017, AV701248, AW959628, AV701431, AW949653, D80378, T03269, AW949633, AW949658, C75259, AV744690, D50995, AW949657, AV719188, AV719557, AW949631, AV742667, AV701125, AV701166, AV700229, AV699746, C14014, AW973307, AW975621, AV702035, D80134, AW949646, AV701419, AV699479, AV701154, D80164, AW959582, AV701443, AV744012, AV720028, D59502, D81026, AV719628, AV718707, D80166, AW966534, AV701130, AV701149, AV701422, D59859, D80268, AV718440, AW965177, AV741221, AW949618, AV645344, D51250, D80269, AW960553, AW966054, D80168, D59467, AV701428, D80949, AV701415, AV701344, AV719000, D59787, AV745080, AV645389, D80522, AV681510, AV681491, AV701021, AV645343, AW959202, AW960414, AV681505, AV681504, AW949654, AV721784, AW973541, AW966013, AV720654, D58253, AW966041, AV719913, AV718674, AV705134, C14227, AW959597, AW964737, C14331, AV720791, AV701004, AW978661, AW973470, D50979, AW966050, AW752082, AV720220, AV700622, AV699669, D80024, AV743601, AV681529, AW966043, AV704180, AV705869, AV706229, AV701224, AW949655, AV719049, AV701330, AV681468, AV720607, AW960465, AW973334, AW964488, AV681474, AV741012, D57483,

095008 09101

				<p>           AV742671, AW973485, AV744773,            AV718633, AW975605, AV681526,            AV727978, D59610, AI905856, AV699682,            AV718931, AV744934, AV645383,            AV645393, AW959799, AV681528,            AV645339, AV681472, AW978648,            AV681507, AV681465, AV681525,            AV681495, AV681486, AV681464,            AV681487, AV681506, AV681523,            AV742720, AW966029, AV742430,            AW966022, AV738928, AV726812,            AW966023, AA285331, AV706147, D59695,            D81111, AV744768, AW949620, D80241,            AW958992, AW964756, AV681514,            AV681477, AV681512, AV681497,            AV681483, AF035294, AB028947, A62298,            AX047063, A62300, AX047062, AX047064,            AR070327, AX020191, A84916, AX020190,            AX033851, AX035434, A82595, Y17188,            AX027925, AR092424, AR018138, D34614,            A67220, AX021518, X67155, A25909,            D26022, AR074545, AJ302649, AJ132110,            X68127, AR025207, A78862, D89785,            AJ294956, AR016808, I19525, D88547,            X82626, AR088705, AX042372, A85396,            AR074141, AB012117, AR066482,            AF260572, A44171, A85477, A86792,            AF058696, Y17187, AR062871, X93549,            AF135125, AX026824, AR008278,            AR077702, A94995, AR087649, AB028859,            AX015396, AX028130, A30438, AJ287395,            Y12724, AB037923, AR016514, AR064240,            A43190, AR060385, AB002449, AR050070,            AR008443, AX014811, X55486, AR038669,            AR071754, A82593, U79457, AR091537,            I50126, I50132, I50128, I50133, Y08991,            U94592, AX009343, AX009341, AX009337,            AX009335, AX009345, AX009342,            AX009344, AX009340, AX009339,            AX009336, AX009338, AX009334, S69292,            AR066488, AR060138, A45456, A26615,            AR052274, AX035429, Z30183, AX035428,            AX035426, AX026823, AR025466, A84772,            AR074139, A43601, Y09669, A84775,            A43192, AR066490, A43189, I18371, X89963,            A43188, A20702, A84776, A84773,            AR062872, AR008277, AR008281,            AR066487, AR074136, AR062873, A84774,            A20700, AF006072, AR016691, AR016690,            U46128, AR054175, A70867, I18367, I14842,            D88507, AR017907, AR067731, AB023656,            AR067732, A58522, D50010, A91750,            AX035462, AB033111, A63261, S78798,            AR051191, Z32749, X92518, AR087528,            AR008408, AR093385, I15997, AR037157,            X98248, D13509, AX046743, A80951,            A64136, A68321, AR060133, AR051190,            AX040581, U45328, I79511, X60736,            AX006825, AX006826, AX006822,            AX001082, AF217994, U87247, AR079804,         </p>
--	--	--	--	---

					AR000511, M60510, AR023705, AB035274, AX006821, AF123263, AB030204, AR032065, and X93535.
HATBI94	34	839468	1 - 1366	15 - 1380	AW966285, AV720255, AW293337, AV725342, AA297446, AA729088, AA296991, and AA334584.
HATCB45	35	631172	1 - 889	15 - 903	AV703043, W80623, AW137678, AW070465, AI922560, BF432932, BE348452, BF941632, BF475288, BE675766, AW003656, BF110027, AI768642, AW470985, AA708150, BE048489, AI689677, AI373102, BE540247, BF063606, AI635762, BE501611, AI587410, AI767805, BE929680, R44963, W78741, AI266349, AI187875, AL533462, AI762852, R19308, R27786, R27875, C01853, AA492441, D63024, AI796430, AA167435, BE669639, BE464050, AA167436, BF110272, BG150184, AC009307, and AC006501.
HATCD80	36	826098	1 - 1795	15 - 1809	AW936395, AA382841, and BF380111.
HATCI03	37	580805	1 - 920	15 - 934	BF802634, AA744060, AW270385, BF804385, AI696854, AW963489, AA700943, AA583394, AI537407, AI453210, AW571963, AW976024, BF530611, AW468372, AA747757, AA225406, AW958962, BE169870, AW855803, AI554471, BF914416, AA687730, AA559219, AA586667, AW272294, AL137119, AJ007690, AC002369, AC002300, AC010494, Z95114, AL135927, AC007227, AC005837, AC004867, AL161731, AC004821, AC011450, AL445248, AC005015, AC004166, AL035587, AL049872, AC025594, AP001760, AC005080, AP001711, AP002028, U07562, AC020908, AC007011, AL354696, AL450224, AL137802, Z97876, AL163279, AC005098, AC004859, AC004815, AL442167, AC005000, AC011455, AL049538, AL137039, AC009276, AC026432, AF168787, AC005899, AP000208, AP000130, AC009244, AL132639, AC002395, AL138787, AF207550, AL139353, AL049795, AC005067, AC005215, AC004967, AC018758, AC005378, Z98941, AC020552, AC002426, AC007057, AL450226, AC074331, AC007685, Z82190, AC005972, AF134726, AC004755, AC083874, AC007488, AC003101, AC004832, Z69710, AC007383, AK023134, AC006088, AC005919, AC006538, AC011495, AL353748, AJ251973, AC007842, AL161937, AL163285, AC004098, AP000096, AC024239, AC068999, AJ009616, AL136170, AL049830, U73643, AC004836, AL356057, J00083, AF064861, AL121891, Z82215, U91323, AC004531, AC067968, Z85986, Z85987, AC007263, AC005412, AL121583, AP000240, AC011811, AC006388, Z82901, AC002316, U80017, AF051976, AC006251, AF228703, AL049646, AC005746, AC005933,

					AC008403, AC005102, AC002470, AP001718, AL137012, AC005516, AL135918, AC008155, AL109976, AC005529, Z95152, AC008521, AC011479, AC006333, AC006285, AC006050, AL158823, AC005775, AL133295, AL121655, AL117336, AL138702, AL109984, AL023583, AC007845, AL118525, AF243527, AC005874, AF134471, AP000503, AC008266, AL033528, U47924, AC010489, AP001609, AC005702, AC008745, AP000553, AC005231, AC005736, AC007066, AC004874, AC004158, Z97352, AC006011, Z93015, AP000065, AC016816, AC008474, Z94056, AC027319, AC010267, AL049873, AC004852, AL109627, AL133211, AL157791, AP001610, AC009087, AC004966, AC022212, AL109798, AC006530, AC005069, AC005694, AC005480, AL391122, AC005004, AP001725, AC011442, Z82214, AC011490, AC004707, AC010422, AL023693, AL035071, AF001549, AC005907, AP001710, AC020904, AC004883, AL117382, AL139396, AL391071, AL159977, AL121914, AC006236, Z98742, AL161725, AL137800, AC002425, AC008126, AC004526, AL138752, AP000103, AC024561, AL034553, AP000338, AP000212, AP000134, AC004985, and AC011480.
HATEH20	38	836056	1 - 836	15 - 850	AW978851, AI686323, AI767653, AV747166, AA829515, BF512171, AA034240, AA053933, AA737691, AA533167, AW261869, AA835698, AA447216, AI623248, W92607, AA835700, Z21891, AA599963, AW893940, W95232, T20153, T20152, R57454, AC006207, and AB020865.
HBAGD86	39	838799	1 - 1699	15 - 1713	AI658681, BE466145, AI806836, AI653272, AA004211, BE302094, BF970406, BE018485, AA418617, AA594901, AI580148, BF589715, AI804211, AI669907, AI342168, AI810310, AA506350, AW022528, H10330, AA721162, AA452114, W03931, AW953290, AI262137, R61309, AA680147, N62384, H10331, AI264925, AA765972, BF086698, AW275301, AA485210, C15277, N79353, AA350799, AI867727, AI474438, AI129224, AA093047, D60782, AI535847, AA897480, AA350798, AV714899, AW956763, and AV728867.
HBCJL35	40	130078 5	1 - 706	15 - 720	AW167360, AI678459, AI817750, AI559589, AA909847, AA101311, AA868436, Z44341, AW661978, BE974135, T71507, D51829, BF109466, AI267409, BE246888, BF803712, BE247681, AA825376, BF795799, and AB007930.
HBDAB91	41	789532	1 - 673	15 - 687	AI167963, AW081006, AA782398, N75825, AW195519, BE858969, AI701657, R45349, R23362, AI468816, BE043035, W07319, AW858522, AW577199, AL135012, BE927373, AW601637, BF084778, AL134110, AL134524, AW577201,

					AL045327, AL045494, AL042523, AW577192, AL045328, and AL042420.
HBDAB91	42	864374	1 - 993	15 - 1007	AI167963, AA782398, BE043035, AW081006, W07319, R23362, N75825, AW195519, BE858969, AI701657, R45349, AI468816, AW858522, AW577199, AL135012, BE927373, AW601637, BF084778, AL134110, AL134524, AW577201, AL045327, AL045494, AL042523, AW577192, AL045328, BF910726, and AL042420.
HBGBC29	43	691473	1 - 1842	15 - 1856	BF223021, BF036281, AI341667, AA180986, AU153625, AU151704, AI093197, BE855464, BE018834, BE616741, BF684563, AI694268, AA031711, AI469856, N63041, N50125, AI150599, AI597740, AI985206, AI671591, W72535, BF431270, AI741942, AA037642, AA180865, AA031648, AA436065, AI800796, AA129939, BF056140, AW002265, AU157670, AI074205, AA830493, BF063800, AI056532, AI656721, W00519, AI275143, AI337739, AW172525, AA443349, AA043021, AA446926, AI655558, AI769027, AA101851, AA917703, W93307, AA526333, AI689128, AA777090, AW002829, BE295568, AW139517, AI128702, AI276137, AW801873, AA873711, AW892754, N98234, W76109, AI631104, AA856832, W92810, AA042939, H87505, AA129938, AI688779, AA693329, AI676108, T87624, AA570072, AA037641, AI186390, AW515672, AA031685, AA037500, R82703, AA037234, AW380430, AA985191, AU131994, BE302396, H87506, AA938640, AI926907, AU118291, AI696069, T74071, AA102060, AW057528, AI671894, AI962374, AI695458, AA046964, BE869607, BF814627, F12449, AA725452, AI968837, AA917824, AA054749, BF437316, F10070, AA917678, BE218382, BE669660, AI916503, AW612381, AA683581, AI984598, AA937814, AI932475, AA046963, AA053281, AI801723, BE858841, AI499751, AA031686, AI074981, AI341558, AI478279, BF735972, AR074859, AK001006, AF038662, AB024436, AF022367, AF158746, and AF142672.
HBGNC72	44	892131	1 - 788	15 - 802	AL526130, AL524570, AW003889, AI935768, AW440485, AI936267, AA713525, AW272919, AI796977, AI951842, AW014081, AI760160, BF941209, AI263194, BF475772, AA496533, AW514179, AA724851, AA496454, AI799782, BF589971, AA496526, BE646016, BE563432, H41355, AW264331, AA515579, AI582716, AI581108, AI208124, AA927044, AI695535, AI638313, BG170255, AI147521, AA199585, AW264237, AW248758, and AB033019.
HBHAA05	45	603174	1 - 676	15 - 690	AI572680, AW631267, BF970107, AA632355, AI433952, AI753969, AA629668, AA493546, AU158457, BF589864, AL044966,



0950082-09101

				<p>           AW518882, AI570067, AI828721, BF028225,            M77888, AI884404, AI547110, BF724416,            AI434103, AV683406, AW836225,            BE391183, N55076, AA610644, AV731938,            AA313025, AA748071, AV743067, AI065031,            AW148964, AI280566, AI732690, AA601376,            AI311796, AI268465, T03928, AU158814,            AW504667, AW880986, AI819419,            AA018258, AA524800, AW971342,            AI791659, AW020612, AV759022,            AV712092, AA935827, AA773560,            AA425283, AI376687, AA493245, AA847341,            BF942991, BF944618, AW303052, AI174703,            BE392753, BG034698, BG223498, BE152006,            AI683079, AA826166, AI590404, AI285651,            AC004531, AL049780, AC006013,            AC005971, AC005522, AL138713,            AC011445, AC008403, AP001725,            AC010458, AP001726, AP001715, AL022323,            AC008569, AL138976, AC005015,            AC005081, AL022316, AC008738,            AC004967, AC004991, AC020552,            AP001716, AL109965, AC006211, AC011514,            AJ003147, AC018755, AC011510, AC006057,            AC016995, AC006334, Z97054, AC006449,            AC005940, AC004383, AL034422,            AL133229, AL079342, AC004051,            AC018719, AP001712, AP001724, AC073316,            AC024561, AC010422, AC020908,            AC000353, AC006487, AC020906,            AC008066, AC006530, U95742, AC007731,            AP000289, AC005500, AC002527, AP000042,            AP000110, AF001549, AL135905, AC007637,            Z98941, AP000688, AC005377, AL050335,            AF053356, AC008623, AC011490,            AC006040, AC004821, AC007216,            AC005881, AL133545, AL109825, AL138756,            AC004703, AL354696, AC005529, AF168787,            AC003982, AC021019, AC005914,            AC004032, AB003151, AC005052,            AC005291, AC004867, AL023583,            AC008752, AL023693, AC004887,            AL078639, AL031651, AC005098,            AC008551, AL136304, AL023807,            AC005921, AC005874, AF134471, AP001714,            AC004906, AP001748, AC004166,            AC020744, AL031118, AL139109,            AC011895, AC006312, AL049537, AP001630,            AC007383, AL109797, AC006970,            AC005399, AL020997, AC005632,            AC005578, AC004522, AF243527,            AC010605, AP000563, AC006511,            AC004796, AC004000, AP001711,            AC002472, U52112, AP000211, AP000133,            AC004263, AJ400877, AL022330, AL161670,            AC008812, AL117336, AC007298,            AL139343, AF207550, AC004963, AL096840,            AC025593, AL035089, AC008969,            AL359382, AC011495, AC005562,            AC002565, AL034449, AL034549,         </p>
--	--	--	--	---

					AC008892, Z98200, AC005102, AC006538, AL445483, AL354720, AL034548, AC002432, AC008982, AC005618, AL109804, AC012085, AC020913, AC022432, AC005520, AC005627, AL021395, AL117380, AC008543, AL158824, AL109935, AL161802, AL031591, AC006536, AC005049, AC002310, AP000512, AC010609, AC019187, AC002350, AC007405, AC005231, AP000696, Z82195, AC004491, AL160175, AL034417, AC009086, AC005523, AF111167, AF111168, AC010789, AL162831, AC005332, AL049697, U91321, AC004126, AC002312, AC007386, AL050307, AL031846, AL049694, AL078477, AC002070, AC006345, AC011456, AL133246, AL133240, AJ277546, AL356299, AL121920, AC008069, AP000107, AC005070, AC008372, AC007225, AC004929, AP001705, AC024082, AC009155, AC022324, AC011462, AC007963, AL157838, AC005057, AL158828, and AL031733.
HBHAA81	46	846465	1 - 1633	15 - 1647	AL138080, AL138081, BG056111, AW117532, AI885223, BF724275, AW051808, AI332809, AI564820, N63569, AW207494, AI866785, AW148811, AI954565, AI199326, AI199105, N38888, AI243422, H09138, AI986175, AW381536, Z25110, AI991904, Z18300, F31192, Z28916, AW381528, Z41646, AI954371, N38887, AI033629, Z28784, AW381480, N94825, F25740, F00372, AW380848, F16581, AW381481, AA197180, AW104762, BF887537, AB042554, AB032999, and AC006059.
HBIAA59	47	806303	1 - 2378	15 - 2392	BG261283, BG029024, AV723100, AV696216, BE931161, BF673030, AV722640, N31939, AI924550, W30681, BF853977, W93554, AV695121, AI693957, AI418000, AI679157, AI271461, AI143084, AI418010, AI186193, BF809099, BF378856, AI199165, AA809478, AI159998, AW250204, AA808416, AW952282, AA593302, AI034450, AI185975, BE677163, AI086931, AI004945, AA776377, N76113, AA588747, AI095797, W68483, AW243597, AA595636, N68971, AA037099, W77975, W48846, AI079380, AI300827, W56274, AI679162, AI168786, AW105640, W92339, BE151400, AI142671, AA573184, W32053, W05603, W88988, W26295, AW263199, W56352, AW263182, AI346401, T68085, AW665393, N99254, M62046, AI284385, N42776, AA443625, BF476230, W46568, W58618, AW189940, AV685659, AA004754, AA779109, BE185831, AI880425, W46651, BF837568, AI081101, N42802, W89081, AI085087, W73911, BE832538, W68301, R67716, BF081619, W92217, T67937,

					W58619, BE185943, AI364156, BF911225, R70823, AW805885, AW089262, N31493, AV722639, H97191, AW140040, H03407, AA010449, AW265360, H60356, BF911333, AI253668, AW835116, BE042937, T12138, AI695815, AA004964, AA331001, T31211, N86748, AW304070, Z43338, AA971863, AA007267, R88101, H59084, R79508, H04108, AA315687, AI139849, AW241439, BE073482, AA338165, R66110, AW608546, AA778276, BE009544, AW241496, AI077770, AA887320, W31537, R79238, R70773, AW516576, W48613, AA338166, BE815610, AI890222, N76384, AA037713, N55230, AI609982, AW967118, AI824249, R45759, AW366951, BF854364, AV742667, AK025783, and AL121929.
HBIAC29	48	831751	1 - 1768	15 - 1782	BE792362, AU119261, AU117435, AU135719, AL522995, AL523062, AV725708, BG257393, AI033807, AL524959, BG258772, BE005784, BF105898, AW157169, AV695633, AI707987, BF965323, BF794704, BG112387, AA877792, AV697750, AU145651, BE268042, AA001404, AI401215, BE568337, AI812036, AW161740, AA428415, AW182986, AI275437, AW160483, AA678033, AI051135, N40528, AI864046, AI292230, BF670616, AA586803, BE564896, AW294845, AA134820, BF245314, AA768585, BF680744, AI816108, AW163143, BF219379, AW161133, AA506157, AA492248, AI298933, AA594861, AA013286, BE673778, AI421629, AA018334, AA905534, AI684501, AA427401, AA082812, BE768733, AA287601, AW157614, BE768540, D79434, BF946436, AI658777, BF541249, AV696430, AA287752, H98178, BF540910, D79454, D79495, AA018335, AV692094, R11115, H98502, BE768718, D79471, BE768713, AV692366, AV694591, AA808656, BE768500, D62307, R84940, AA037183, AA804276, D79442, AA284650, T93079, N46576, D62351, D62288, AV696341, AW440057, D62360, AW163531, AI057422, D79425, R11059, AA359531, AU156237, W27841, AA013043, AA323139, BF219327, AI991658, D79500, AW613472, BE768582, D61951, T93170, AL524958, BE544935, D62369, AV657638, D62200, AI702248, BF948281, AA134819, D62264, D79447, AV656951, D79470, BE043970, D62243, AA934540, N67412, AW118285, AA483825, D62298, AI911842, AI033466, R31032, X93846, D79508, AI371542, R31522, D79465, AW590942, AW466905, AA180979, AA059251, AI689493, AW661808, AW006273, D63025, AW449881, AI866210, AA059003, N71916, AA381297, AW873906, AW590669, N55843, D79405, AW169134,

					BE390691, AA585262, AI754249, BE389098, AA059266, AI459286, AI991649, N84299, AI655153, BE548704, AW016856, AI400471, AI400469, BF218780, AY007166, AL034379, and AK021792.
HBICW51	49	553630	1 - 605	15 - 619	AW976674, AW977839, BE046511, AI143906, AA837255, AA743668, AW768572, AA699707, AA848026, AA805369, AA765722, AA813921, AW299915, BF229206, AA291129, and BF754240.
HBJAB02	50	837309	1 - 1679	15 - 1693	AL529646, AL529645, BE898304, BG112747, BF791411, BG036058, BE392384, BE621757, BE548173, BE895853, BG034671, AA808894, BE901085, BE278873, AW152607, BE795658, AW166898, BG122141, BE782474, BF972826, BE793716, BE140314, AW750993, AA826362, AW517942, BE714673, T59668, BE731030, BF939314, BE732766, BE745104, AI290469, BF477770, AI805651, AI961329, AA581089, BE902575, AW197375, AA974066, AI950259, BF802171, W27729, AV693783, AA877530, AA715365, AI968889, AA885542, AA160748, AA386371, AA335719, BF873961, W73105, BF223151, BE740826, AL120854, BE548914, AA318192, AA501478, BF125073, AI948815, AA581100, AA658457, AI621069, T59802, AA468534, AA503715, BF850755, AW956069, AW841506, AI144504, AA352215, BE897964, BF883404, BF373009, BE090290, BE168997, AW855521, AW820855, BG230749, BF376598, BE622839, AV699089, AV647789, AI567702, AV726156, AW961037, AW411235, AV726058, AW020397, AV706279, AV702427, AV651955, AV702026, BE393551, AV727787, AV660608, AV687176, AW021717, AV698545, AV687909, AV709256, AV708438, AV656903, AV661704, AV696106, AV697196, AW409775, AW951263, AV689111, AV655280, AV728157, AV692345, AV659322, AV654908, AV656478, AV708893, AV709314, AV708381, AV660728, BG168549, AV659536, AV691080, AV706219, AV695545, AV652001, AV705159, AV648263, AV703169, AV728518, AV707541, AW952409, AV709660, AV726624, AV706854, AV729220, AV709604, AV687035, AV696866, AV728997, AV704955, AV726816, AV725920, AV652156, AV701707, AV656283, AV704234, AV708025, AV707933, AV684604, AV729378, AV708980, AV692691, AV701914, AV708723, AV702516, AV693523, AV709407, AV705693, AV708992, AV729263, AV726103, AV708704, AV727029,



					AW262857, AI446447, AA852298, AI654262, AI184300, AI128506, AA552369, AA604635, AA583259, AA530888, AA158420, AI696777, AA968781, BF435939, AA558592, AI679597, BF195650, AU147255, AI184403, AU154997, AA723126, AA828222, BE301243, AA156574, AU144430, AA632393, AU158982, AI191123, AI889491, AU159099, AA662257, AI814093, BE315360, AU146056, AU154857, AU155238, AI888961, AU152381, AU147517, AU155476, AU148092, AA771920, AI129393, AU159433, AU147240, AU147366, AU148271, AU147605, AL047763, AU146002, AA554695, AW273189, AU145732, AW070654, AW276349, AI571606, R54732, AI660822, AI678682, AI818491, AI803888, AA599490, AI814696, AW197992, AI628530, AL135927, M13452, L12401, AC007227, D13181, D49733, AK026584, X66870, X76297, X03444, AC006441, AL080314, AL137141, R23628, R74021, R74022, H02237, and AA158412.
HBJBM12	52	560606	1 - 1121	15 - 1135	AA722101, and R18111.
HBJCR46	53	815649	1 - 3194	15 - 3208	BF980168, BF001800, BF221545, BE180558, BG117357, BG165887, BE747286, BE867206, BE559905, BF029089, BE884542, BE180560, BE882087, BF672818, BE180608, BG255311, BF695020, AI765879, AV701340, AI832097, AV701354, BG164080, BE568492, BF979546, BE268392, BE180559, AI927915, AI675415, BF195785, BF662916, BE673547, AI086866, AI956035, BF671543, AU146956, AA479515, BF063974, W19888, AW992096, BE504075, AW069858, AW992159, AA626631, BE019647, BE221636, AA479513, BF575729, AU144777, AA973047, AA936602, BF671187, BE545703, AV701112, AU144811, AU160319, AI472144, AI263407, BE019684, AU118130, AW614133, AI954073, AI767153, BF445898, AW087744, AI913738, BE397963, AI628089, AI675273, AA447852, AI635143, AW129685, AI287605, AA486193, W00613, AW135604, AI754985, AI334344, BF062454, AW119185, AW576204, BE041839, N54388, BF354864, AI275063, BE175440, AA883965, AI554276, AI082201, BE718001, AW771023, AI274243, AW337565, AA150018, BE175441, AW771580, AA447700, AW052155, AW771203, AA085991, AA085621, AI954014, AW068250, BF725222, AA486299, BF087209, BF834780, AA971016, AW854246, AA150083, H08089, BF735392, BE180609, AV683146, AA677738, AI220075, AV701437, AA740363, AA909807, H71626, R61225, N49411, AV689665, C05160, AA888983, AI026772, AV692963, AW580238, AI816858, AA340276, BE180555, AW862217, R61226, H82142, AW068158, H08090, BF997557, AW862230,

					AI565387, AI824475, T32511, BE169765, BE002988, AI004624, AW514293, AA775740, T31977, AI660017, AA347780, N63302, AW468684, AA908821, AA953180, AA897589, AI217393, AI536640, AA337664, BF364138, BE816022, AA652646, Z19403, C04239, H71627, AA347779, R14260, AW168255, BF833756, Z42097, AV655482, AI431630, BE088874, BF348891, H82048, AA130053, AA781872, BE544862, Z28501, AI434730, BF575085, AA328725, BF084834, R96217, AU155530, Z38367, AA301061, AA370106, AI810366, AW836437, AA371349, AI991790, H54404, AW015692, Z42140, AI905103, AW242449, BF327421, AI824674, AW195816, BE173133, BF812520, R86231, AI274472, H54488, AW379997, BF476783, D62662, AA382472, AW379945, AL079373, AA703490, AW381290, BF328480, AW392784, AI823416, BF961200, BF910517, BE728262, BE408949, BE277648, AA458950, BF797287, BE866243, AA923063, Z24943, AA236098, BE160734, AA397692, C02287, BF812876, BF444939, AF150734, AK000984, AF124434, AL033531, AL031287, Z97876, AC007034, and AL035304.
HBJDS79	54	813588	1 - 2311	15 - 2325	BE531135, BE614809, BF982395, BG122680, BE615130, BF968072, AV706034, AI222785, BE539326, AI598156, AA131643, AA626884, AI571234, BF036649, BE743394, BE542106, AA770214, AA203111, AI608599, BF035790, BE276209, BG253824, AA436283, AI094146, AI085480, BE223034, AA577314, BE615635, AA436157, AI338243, AW081389, AI224507, BE742762, BE616645, BE879846, AA131533, AA631769, AI348255, AI870284, AI278842, AA971726, AI369248, AA934527, AI274146, AA279453, AA021050, R63988, AW952487, AI869674, BG056655, R78856, BF111769, BG056718, BG056695, BE255966, BE018968, AA551934, AA284976, AA678694, AI202603, AI097586, AW151699, N64789, AW135978, AA471054, AA742282, AI340009, AW873159, AI301870, BE206676, AA730072, AI278735, N22046, N52849, AA324335, BE774969, AA707878, AA312623, AI206104, AA922420, AI081594, AI203119, BE717107, AA770407, AW675278, AI263743, AA291551, AA298598, N76200, H95836, BE770889, AA962696, T15486, N80747, H30520, BE673804, AI520678, AI918182, AA283618, R17600, AI474363, BF806759, AA939087, AI242964, R69061, AA532553, BE770975, BG055182, C03365, T46915, BG029466, R23915, D25896, AA872629, AW860570, AA323822, AW860432, BG011166, AW860428, AA551996, BF898533, BE161547, AW131992, AW862507, AW860442, BF769056, AA551781,

					BE696376, AW860588, AW860592, AA330149, T34728, R69060, AA419231, AW860586, R49598, AA425955, T74490, H05532, AI718310, AI805179, AW610364, BF840958, R79345, R63904, R23914, AA296953, AW292997, BF338486, BE208400, C02983, AA365446, T71450, C02552, AW607020, AW071145, AW610371, BF825608, BG010487, AW603550, AW371102, BE717176, AI587180, AA578420, AI651804, H95835, AI074093, AW363067, BE903315, AW952592, AV707094, AW952361, AW963627, AW952051, AV661490, AW960297, AV645577, AV701728, AV661494, AV654287, AV705143, AV704833, AW954145, AV704504, AW955632, AW958455, AV706272, AV706289, AV725807, AV726142, AW966044, AW959346, AW949447, AV704144, AW963622, AW952181, AV726831, AW952298, AW951445, AW960663, AW951711, AW945159, AV707503, AW950095, AV645936, AW952167, AW961409, AW958088, AV705115, AV703628, AV704342, AV705002, AV706158, AV706028, AV704330, AV705199, AW952418, AW965813, AW963231, AV709660, AW963026, AV702120, AV706714, AW959755, AW960821, AV706446, AV703833, AV703425, AW961597, AW952031, AV706432, AV729597, AV648461, AW949354, AV725900, AV729388, AV707238, AV706910, AV687519, AV658743, AW953804, D50992, AW951951, AV724996, AW967188, AV709751, AV725369, AV707196, AV706850, AX018173, AX018065, AL096840, AL021155, A62300, Z30183, U94592, A62298, A82595, A82593, AX047063, AX047062, AX009343, AX009341, AX009337, AX009335, AX009345, AX009342, AX009344, AX009340, AX009339, AX009336, AX009338, AX009334, AR050070, AX047064, AF006072, AX040581, U45328, AF320628, AB030204, AX046743, and AR025466.
HBJDW56	55	520401	1 - 623	15 - 637	AC005532, AL031319, and AL354933.
HBJEL16	56	847030	1 - 736	15 - 750	AI279501, BE867835, AL528252, AA569392, AW856935, AA071326, AA587712, AA258409, AI341817, BE898008, BE696253, AW576885, AA837880, AW576895, AI584147, AL525748, AW383278, AA071368, BF330803, AW383120, AI393286, AL513864, C00710, AW857093, AA644480, AL528253, AW499908, BF326342, AW749039, BF771813, AA193585, AW383268, BG031591, AL040224, AW383242, BE904616, AW751656,



FD-2760-28005663

					AW383266, AW383144, R15553, AW383205, AW383131, BF931485, AA258753, AW383275, AW383146, AW886371, AW997055, AW996855, BE929916, BE005368, AA188924, BF350669, BF327032, Z99943, AL035308, AF087020, AL035302, AF092425, AF095727, AF092424, AX013163, and AF239756.
HBJFK45	57	531919	1 - 529	15 - 543	AA714560.
HBJIG20	58	866159	1 - 623	15 - 637	AI114569, AI174851, AL538129, BE890567, AL525022, BE877813, BE880157, AV725294, AI133486, AV725311, AI174902, AI114699, AI111183, AA639334, AI114553, AV702695, AI133672, BE880648, AI147501, AI133302, AV706307, BE880425, BE873842, AL048390, BE877322, AV706023, BE879304, AI749163, BE881359, AV715862, AA642199, AI114736, AV713805, AV727892, BE881076, AI750078, AV752193, AI133348, AV709557, BE872112, AV763516, BE877234, AV703758, AV715160, BE876719, AL037681, AV701396, BE879580, BE880140, BE875398, AA583348, AV711558, AA196384, AA723026, BE876143, AV723395, AV706568, BE879909, AV707919, BE875361, AI279442, AA736456, BE876043, AI133323, AV705912, BE877386, BE879135, BE875253, BE872677, AV710081, BE880711, BE877647, BE881218, AV763759, BE875792, BE876085, BE873841, BE873343, AV702923, AA640938, BE875858, AI207615, AV705995, BE879968, BE874846, AV704327, AI721239, AI609232, AV716218, AA737196, BE877268, AA929066, AV709305, BE876531, BE875412, BE879974, AV708006, BE881478, BE880128, BE881645, BE878122, BE880497, BE874647, BE877686, AV700257, AI720842, AI720161, BE874823, BE874885, BE892487, BE875889, AL513828, BE891700, BE878894, BE879695, BE898937, AV723029, AV717787, BE867307, AI833114, AV706277, BE891387, BE878086, AV729201, AI709043, BE881222, BE877757, BE877150, BE876704, AV710239, BE870155, BE880602, BE876425, AI814650, BE881146, BE878626, BE878178, AI750175, BE875408, AV717612, AI833042, BE880690, AV711143, BE878330, AI720252, BE897096, AV726346, BE874215, BE877004, BE874734, AL038791, BE879775, BE870577, BE880762, BE876080, BE879410, AV716025, AA736459, BE874520, AW027357, AI832465, AW081665, BE877131, BE877496, BE876651, AV648404, BE877242, AA195996, BE873478, AV729465, AI985879, AA652417, BE874662, BE896548, AA574324, BE890719, BE875038, AV728251, AV706779, BE874747, BE875536, BE877390, AI832606, BE898560, BE873954, BE880948, BE875601, AV759353, AV710539, AI766340, AV705883, AI832709,

					BE899424, BE876550, AV706266, AI065146, BE880722, BE880442, AI832520, BE877488, BE880237, BE877598, AV708807, AI833048, AV716989, BE879895, BE875795, AA653037, BE876098, AI133444, AV729314, BE879455, AA583501, BE879522, BE878593, AA641024, AI719037, AV752966, BE875905, BE894635, BE875968, BE896837, BE874696, AA745376, AA738012, AV729229, BE898750, BE878213, BE867317, BE875261, BE869133, BE874445, AI708900, BE879302, BE880950, AI924816, BE896982, BE877497, AV693283, BE898873, BE874003, AW027020, BE878079, AI444653, AV701133, BE880568, AV724988, BE868527, BE878210, AV725316, L08441, AR083056, I27367, AX039612, X62996, V00662, X93334, X54629, D38112, D38116, X93335, D38113, AL359334, X93347, D38114, X97707, X84075, Y10129, AF134583, AF004341, U35430, AX026544, Z79403, Z79419, Z79459, AJ252237, and I07623.
HBJKD16	59	853358	1 - 1615	15 - 1629	BG259133, BG258754, BE895955, BE535182, AU119776, BE296370, AW361272, AA205862, BE079707, AI743764, BF669591, BF102652, AA768863, AA767455, AA683506, BF374311, W39021, AA583062, W94893, BE737722, AI128320, AA703242, AA402965, H12229, AA027163, AI338954, W92057, AA769377, AA811137, AW470052, AA027162, AV713020, AI680487, R19758, AA283191, AA531492, AI700367, BF946853, AW514119, AA644413, AI004120, AW876599, BE868122, AI831977, AI219655, AA197283, H17722, AA890197, N70828, AA164346, AV750338, BF791719, AA164345, AA767095, AA393969, AA765421, H17611, BF247968, AI254347, AA253013, AA534905, AA470446, AA182675, H12230, H43795, T32973, AW386765, AU146033, AA078964, R51414, AI269757, T33350, R45177, AI203452, Z44609, AI041094, AV749975, H43709, BF336945, AI268058, T30703, R85203, R51302, R51996, BE738484, W84649, R51995, AA824604, W01425, AA907096, BF801374, BE544754, AI469616, AU156273, AW297202, T83337, H14279, BE929477, AA252973, BE242053, T18899, AA248529, H14306, BE698663, T33352, BG059012, AA361817, AA236020, AA810717, BE243263, AA876139, BF240162, BF893858, BE217859, F13420, H53830, F08935, AA078866, AV727864, AA721692, Z40480, AI383780, H52715, H48241, AW751348, T83485, AI567913, AI247225, BE694869, BE940539, BE713625, BF858355, BF801742, BF824816, BE713728, BE713771, BE713469, AA463921, C01352, AW608887, BE714022,

0950082-091201

					AW876620, AW876628, N89830, BE002346, AA192537, AA319174, BE714013, AV709029, AW991462, AI080442, BF993596, N57976, AA034034, BF799178, BF818732, AW392976, AW876544, BE568245, AW835734, AK000087, AK021899, AF217983, AC008074, and AK024658.
HBMBM96	60	561935	1 - 1062	15 - 1076	AI888795, AA047754, AI561027, BF676343, AA047704, AI187148, AA314069, AA536040, AW976024, AA704393, AI754653, BE973547, AV762633, BF857849, BE897079, AU144320, BF681619, AV757032, AW972919, AW819125, AW151824, AV763457, BF854308, AI306232, AI251576, AA904211, BF589788, AA812058, BE245576, AL042667, AL042670, AI521525, AI891080, AW961593, AI583466, AW274191, BE878926, AW020150, AI459904, T74524, AI280266, AI459943, AA653139, AW572721, F16345, AV729669, AA515728, BF805088, BE350953, AI601229, AA297802, AA747757, AA297145, AI926102, AA629540, AI436433, AI679221, AA084609, BF724838, AW270385, AA164955, H59737, BG029528, AI473995, AI340641, AW504667, AW969831, AA805049, AI858691, AI749893, H07953, AC000057, AC011484, AC005920, AC005225, AC006011, AC006126, AC005837, AL031658, AL109825, U85195, AC005940, AE000658, AL109804, AL049766, AC004089, AC002352, AC005015, AC004797, Z93930, AC009756, AP001711, AL035422, AC005519, AC008403, AL109935, AC004878, AC005696, AL159977, AP001725, AL121983, AC005907, AC009060, AL353807, AF111169, AC005231, AL049760, AP000547, AC002425, AC005544, AC011442, AL031228, AC004526, AF243527, AC005081, AL035681, AC005291, AF196972, AC004019, AC020550, AC003982, AL034423, AL031311, AC000025, AC007664, Z93241, AC009032, AC005944, AC002039, AC003690, AC005756, AF196779, AC008760, AC008265, AJ295844, AC004150, Z99716, AC022392, AC005399, AL353194, AC011895, AC027319, AF168787, AC002470, U91326, AC011479, AC005098, AF228703, AC003104, AC006013, AC010422, AC008569, AL158830, AL118520, Z83845, AC024075, AC007536, AC005332, AL117348, AC004686, AC010677, AC018720, AC006211, AC008745, AL445435, AC004383, U91321, AC004971, AF001549, AC007225, AL162615, AC010271, AC020906, AC010748, AL139396, AC006121, AP001727, AC020934, AC011445, AL096701, AP000045, AL031597, AL024508, AC007690, AP001747, AC011495, AP000300, AC020913, AC002549, AC006536, AC005821, AC004893,

					AC005529, AC005071, AC006965, AL157938, AP000692, AP001728, AL445263, AC007957, AP000347, AC009314, U91322, AL137918, AL022316, AC009506, U95090, AC007055, AC009086, AC007151, AL022476, AC004965, AC011497, AP000113, AC005052, AC004890, AC020904, AL050341, AC006077, AC008474, AC020916, AC008623, AC008102, AP000553, AC008551, AP001718, AL162272, AC006451, AX039602, AC006023, AC004813, U91323, AL008721, AP000117, AL365505, AL031721, AC004821, AC005701, AC006441, AP000212, AP000134, AC010618, AL049569, AP000193, AC004824, AC005088, AL049757, AC022436, AC004882, AP000050, AC004477, AC010519, AL445483, AC006210, AC010412, AC087093, AF207550, AL109758, AC018758, AC021999, AC020754, AL132855, AC004867, AC006511, AP001717, AC011464, AC009247, AC008072, AC005823, AC008273, AC004166, AL121932, AL009181, AC007386, AC008753, AC004841, AC008750, AC003029, AC006071, AC007193, AL121928, AC008687, AC006452, AC005412, AL139353, AC009470, AC004253, AC011489, AP001752, AP001760, AC005952, AF176815, AC008521, AC000378, Z85987, AC023344, AC006483, AC025430, AL133350, AC083863, AC005228, AL158040, AC002472, AL033529, AC006538, AC005379, AC006254, AL138680, AC005300, AL118502, AC006241, AF030453, AC018755, Z98048, and AC002563.
HBMBX01	61	705047	1 - 1638	15 - 1652	AW961140, AV653628, AA284390, AW298801, AA325985, R05810, AV647470, AV647506, AV646943, AV646939, BE079602, AI435248, T91124, BE079603, AI217083, R05895, R05360, T84795, AA903614, AA640495, AA654849, AA002218, AA292719, and AC004236.
HBMTM11	62	589515	1 - 1625	15 - 1639	AU117599, AU133725, BF329048, AU135386, BF215887, AU134241, AL134928, AU144387, AA551031, BG254665, AW961387, AU151757, BG251511, AA713483, N22769, BE675412, N34769, AA166919, AI890079, AI361889, AA668981, AI209020, AV651304, AI588918, AV651702, AI240990, AV651711, AV651263, AW664145, AV650032, AI200221, AA741144, AA490899, AA830016, AW589574, W96201, BF028879, AV650101, N76715, BE816932, N76708, H95025, AI886459, BE090531, AU155873, H04255, AA628554, W96170, AW243986, AA557749, AA770526, R68746, AV651888, R68691, H02957, AA328145, BE816925, AV655159,

095002-0101

					AW809606, AI081357, AV651866, AW818352, AW818356, AW295753, Z38626, AV651172, BF185552, N36348, R33453, N94910, BE887218, R33554, Z45697, R31757, R36180, R31707, AA742488, AI286227, AA693346, AA622811, R27711, N59241, W23502, H94002, N59247, AV651196, AI269719, AU135179, AA897483, AW856145, AA873441, AW388097, AA166811, AC005412, AK001838, AC020717, AC022014, AC026168, and Y13537.
HBMTX26	63	695704	1 - 1294	15 - 1308	AW861714, AV720211, AW973541, AV719822, AW975623, AI061313, BE138594, AW973992, AA833875, AA833896, BG236628, AV756491, AA468505, BE139267, AA284247, AA644090, BE062478, AI733856, AW502873, BE139358, AW410409, AV760508, AI216990, AW237905, AL079734, AW855643, AI753672, AI076228, AW502237, AI755202, AW963463, BF725844, AV700760, AA420546, AI066646, AW970896, AA502991, AA610509, AI912401, AW505253, BE062476, AV758870, AI754653, AW328331, AI284543, AI675615, BF923365, AV718487, AV720729, BE178481, AI625604, BF526964, BE139139, AI250552, AA535216, AL135377, AI053696, AA632744, AI887235, AI251034, AW851816, AA595499, AW243793, AI254779, AW327624, AW271977, BE968744, AA837035, N23913, AI345695, AA693484, AW023111, AI251284, AI251203, AI440117, AV734149, H07953, AA593537, AV720616, AV754716, AW845366, AV719632, AA856961, AA524616, AW301771, T05118, AW328000, BG000961, AI362442, T74524, AA704393, AI753113, AW275432, AV699480, BF725761, AA630854, AI003611, AL138759, AF205588, AC006120, AL132671, AC002289, Z82899, AL133247, AL117337, AL121835, AL049697, AP001671, AF235098, AC004002, AL160231, AF095725, AP001728, AL359272, AL031176, AL049874, AC007488, AC005741, AL022151, AC000120, AC005534, AL031673, AC018769, AP002534, AC018719, AL049778, U52111, AC008482, AC008929, AC006115, AC003013, AL157951, AL354696, AC008602, AP001678, AL022400, AC024581, AC003667, AC008462, AC006440, AP001712, AC004492, AC002994, AF109076, AL360078, AP002533, Z83843, AL022146, AL359846, AC008498, AL137013, AP000517, AL049765, AP001669, AL136101, AL136297, AJ271735, AC048352, AC006010, AC002302, AL157827, AC003966, AL355838, AC006313, AC007335, AP001412, AC003035, AL121901, AL136972, AL354811,

0950082-09101

					AL078604, AC009487, AC004075, AL136168, AL161771, AC002412, AC007282, AC003987, AL138815, AL355136, AP000152, AL132994, AC006966, AL365271, AC078889, AL359378, AC009318, AL356969, AB023054, AC064878, AL356473, AC005541, AC010223, AL138702, AC006924, AC008543, AL122057, AC002038, AL157360, AL353140, AC009404, AL158198, AL132773, AC004189, AC007000, AL390035, AL109956, AC002357, AC005156, AL035402, AL049835, U80460, AL357153, AC012081, AC005095, AC004673, AC025226, AC012558, AL136419, AL031671, AC022148, AP001660, AC005692, AC068999, AL030995, AC009073, AL356795, AL022577, AC004074, AC005191, AL023876, AL390121, AC012077, AL358293, AL121984, AC004981, AB020877, AC007132, AC005536, AL035698, AL137191, AC000119, AL078591, AC008613, Z98946, AL121999, AC002086, AL121781, AC016831, AC009508, Z83846, AP002898, AC002072, AC007262, AC021012, AL355093, AC015592, AF011889, AP001870, AL139350, L11910, AC083862, AL024507, AC005216, AC011890, AP002532, AL031347, AC016395, AF117829, AL031433, AL163207, AC008716, AL050308, AL032821, AC004049, AL049838, AC012085, AL163218, AC009230, AL136531, AL117344, AC008806, AC007486, AC007458, AL390039, AC008372, Z76735, AL358336, AC009953, AC006322, AC024082, AF241726, AL121893, AL122023, AC012150, AL353660, AL109759, AC006430, AL110505, AL356098, AC022401, AL133153, U96629, AL031228, AC062033, AF121898, AL049550, AL355836, AL441883, AL049834, AL035552, AL354750, AL133325, AC005537, AC007037, AL031387, AC073054, AC000114, AL391986, AC016594, AL139109, AC012082, AP001684, AL023284, U91322, AC003668, AL078623, Z97180, AC004063, AL138783, AL354942, AL021368, AP001720, AL355296, AL031114, AC002524, and AL353732.
HBMTY48	64	637521	1 - 1877	15 - 1891	AV714718, BG177227, BF669308, BF209846, BE748253, BF243099, BF209247, BE677537, BF243862, BF215098, AW402403, AA814374, AA557486, AI445815, AW500029, AA128592, AI610201, AV701116, BF870951, AW974932, AI859834, AL037910, AA828047, AW973400, BF761328, AA720732, AL036283, AI859946, AW975634, AV761519, AW271904, AW302709, AV762047, AV733328, AV758989, AA720702, BE910362, BF475757,



					AC006965, AB023054, AL135818, AL136124, U73023, AC005940, AC010719, Z97054, AC002310, AC011500, AC020916, AL096840, AC005529, AL078581, AL136300, AC004801, AC004016, AC004867, AC005070, AC000120, AC006254, AL096870, AC011464, AC002366, AC002073, AC002996, AC005484, AC007282, AL157372, AC005103, AC005874, AF134471, AF047825, AC004996, AC005740, AF196971, AL163281, AL133448, AL032822, AC005837, AC005899, AL109797, AC006967, AC015550, AL121926, Z99572, AL138716, AC011455, AC004858, AL050318, AC020663, AC006539, and AC002400.
HBMUH74	65	866160	1 - 712	15 - 726	AI633540, BE999936, AL529110, AI911597, AW016785, AA479308, AI381011, AI057451, AI283542, AI224172, AI025510, BF929951, AW589256, AU156824, AU155569, BF063133, R43074, R25758, BF818086, AL529111, BE567017, BE077233, H09061, AA479409, AK001927, AK001324, and AC009318.
HBMWE61	66	778066	1 - 1104	15 - 1118	AL530335, AW182591, BF435671, AA776879, BF435138, BF435606, AL523150, AA707339, BF055381, BE312352, BF888738, AW513106, AL119508, AI684324, AI827310, AI538166, AI457932, AI420719, AI889349, AW026348, AA456678, AI567105, AW316798, AI362960, AI313262, AL523149, AW189249, R42782, AA564318, AA995148, AA598982, AA161330, N30438, N30445, AW771697, AA448673, AI334337, AA194840, AW149443, T99604, AW969668, AI915277, AU145646, AA214246, AW085918, AA161331, AA501503, T57340, F36146, BE700927, AA528196, AW956355, AA436448, AA436494, AW339369, AW467983, F10251, BF057117, AI049659, N67950, BF934197, AA653249, AI805953, Z39922, BE700856, AW137398, BF927130, N94961, R56027, AA377531, W38702, H39550, AW878949, AL049732, AB029037, AK001300, AR027928, and U04811.
HBNAX40	67	834801	1 - 2779	15 - 2793	BF966078, BF792338, BF034911, BF217973, BE883387, BF947401, BF574197, BF060683, BE220005, BE645102, AI808818, AU158323, BE222311, BE467629, BF985268, AA203305, BE504175, BE612371, BE504478, AI890286, BF514573, AW173142, AI674096, BE301797, AW962903, AI674111, AI935063, AW958697, BE931820, AI431629, AI418384, AU157624, AW958686, H10461, AW995348, AW511978, AA864829, N29528, AI287632, AU157306, AW206871, AI381961, BE018315, AW238878, R61198, H80193, AA531283, AA565321, AW073280, AW026572, AW243789, AI819460, AI913516, BE503173, AA305587, AA305897,



					AW952734, BE815642, C01749, H10462, R58932, AI040839, T17118, R62976, H17748, H11865, R63031, AA970617, AI247608, R63338, T07723, AA774553, AA371079, BF084654, AA329699, BF802622, AW139568, AW027833, R61199, H11505, N57467, R19912, BE695240, R63337, R06902, AA337757, R33944, R14158, T55659, AA247547, AU140145, AA329948, AI872648, H17635, T55578, AI802966, N67275, R40453, AA318909, AA039614, AI565257, AI540752, AW865932, N46626, R06946, N40466, T25153, AI444969, AI262121, H78817, AW150803, AI003014, AI457343, N34900, AK023746, AF305081, and AX029397.
HBNBJ76	68	810332	1 - 1960	15 - 1974	AL534356, AL529732, AL534050, AL534051, BE745824, BF966566, BF966920, AU143029, BE250074, BE743214, BE293428, AA521297, BE293836, BE890115, BE273039, BG029074, AW576992, BF920823, BE299201, AU147440, BE262643, BE297473, AU160197, AA237063, AW749171, BE899113, AI818118, AA477060, AL522323, AI863128, AI094964, AL526791, BE541518, BG001429, AI073714, AA557526, BE312979, AI688523, BE880788, BF693443, AI818673, AA476949, AA504527, AA237017, AI754204, AI073509, AA236864, BF846722, AW129564, AI075855, BE305107, AI078504, AA503761, AW953337, AA410682, AI906296, BF921511, AI620391, BF930664, H28108, BF934123, BF930777, T80380, BF934139, BF921439, AI669584, AI074509, BF921504, AV696486, BF920987, BF921181, BF930672, BE619914, BF921183, AI540579, AI087186, BF930838, AL526822, AI342200, AI190396, N25006, BF930678, D53101, H15668, R91002, AW189172, BF921415, BF920975, AI445693, AA504456, BF568918, BF512279, AA235460, T33078, H38314, N23507, H15669, H41352, AI688360, T33077, BF921110, AW168851, AA932279, W28365, H06522, T16220, R50974, BF929910, H93073, R66792, H49516, AA432099, BF809362, BE787105, BE243707, H17715, F24147, BF939372, BE047893, AA835754, BG003276, AA235568, AA584011, BE398065, T33748, W99346, BG004624, H95616, AA938420, AA292524, AI372738, H09967, AI569025, BF921193, H30757, BF802828, AA719210, AA322159, H97134, BF930773, N23060, R88745, BF877775, R43985, AL135592, D45278, H98793, W99304, H06577, H17097, AI541278, T34955, H23692, Z39225, H88727, BF894389, T32634, H95901, BE207940, W16965, AA831137, BF926660, N55137, F32307, C15500, AA191241, R81752, T30362, F10931, R35432, AA362861, AA852184, BF946139, R13653, T31151,

					H95902, Z42100, N89617, AW731979, R85881, BF946142, F01633, AW674315, AW882084, AI372739, F10929, AA322320, AA337191, AA446785, AA360857, Z43144, R38598, AA938068, R39608, AW795181, T31175, T09252, BF946140, R38830, BF812774, D52713, AA350382, AA251259, T33747, T53545, T03354, AA505949, R49230, R13752, BF946137, BF818030, BF798116, AA429879, BG007235, AA830961, C15188, H93938, R84492, T33382, AI372737, R14418, AI025547, AA716021, AI005001, AI372403, AW380183, BF880404, AA331873, BF924213, BE543234, AI208872, BF526157, BF762225, AX015352, AK026126, AC004453, AB028991, AK025555, and AK024226.
HBQAB79	69	810542	1 - 1317	15 - 1331	BE926412, W27043, BG006701, AW500368, and AB020689.
HBQAC57	70	793814	1 - 2097	15 - 2111	AI087837.
HBSAK32	71	856387	1 - 578	15 - 592	AI740936, AI742064, AI832483, BE856354, W89126, AI741855, AA552666, AL525133, AW293469, AI032044, AI769344, AI199155, AL537059, AA769290, AA481420, AA425849, AA968823, R73406, AW290963, AA653956, AA481658, AA244354, BF477489, AI278115, BF664060, N92264, AI014386, N45235, AA723656, AI354229, BE041734, W24441, BE350121, BG109716, R73405, BF690465, AI675727, AL530882, AA570628, AA992527, AW089841, BE858139, C21531, AA029467, AA029534, AI951077, BG004006, AK026029, and AL442086.
HBXCM66	72	639039	1 - 996	15 - 1010	AW970983, AA311432, AA515629, AA515360, AW973992, BF804385, AL046519, AV703785, AA502991, BE968744, AA631507, AI206841, AV763026, AV763058, AW613805, AW340905, AI431303, AI366555, AW873061, AI284640, AW270258, AW341903, AA809546, AI821716, AI613389, AV704541, AI499954, AA533025, AW438542, AL079734, AV655282, AL079645, AL046409, AW023672, N23504, AW419262, AV759632, AI061313, AW872736, AI189682, AV738383, AA533176, AW193265, AW069227, AW023111, BF725844, BE139139, AI284543, AI254770, AW979087, AW026305, AI421950, AI419337, AW872676, AW970981, AL038936, AI634187, AA515351, AV762633, AW327624, AI251034, AI223626, AW407578, AI635440, AI003611, AI801482, AI250552, AI754170, AW574899, AI251284, AI251203, AW130188, AW303196, BF827410, AV755512, AI355559, AV763584, AW845366, AW979241, AI891080, AF123761, AP001711, AL122015, AL035249, AC009247, AL133545, AP001725, AP000252,

095008280550

				AL133453, AL117258, AB023050, U80017, AC004560, AP000557, AC011444, AC005015, AP000212, AP000134, AC011742, AJ003147, AC007957, AC018644, AL121936, AL049713, AC002527, AC004910, AC008012, AC023490, AC007270, AL035420, AL136305, AC005911, AL365505, AC006077, AC000353, AC011491, AC016554, AC016995, AC012077, AC020934, AC007686, AC006530, AC003098, AL356020, AC004999, AC011480, AC011479, AD000092, AL162390, Z81364, AP000031, AF053356, AC013434, AL357497, AC005800, AB023049, AC006121, AB003151, AC002544, AC005288, AC004841, AC004166, AP000689, AL023553, AC005839, AC005731, AL121749, AC006167, AL050335, AC004965, AL022723, AL035089, AP000556, AL021939, AC005080, AL137230, AF165926, AC004972, AP001748, AC004605, AC010620, AL121893, AL353777, Z98200, AC004033, AL132825, AL445483, AC008555, AL139100, AC004912, AC011477, AC006483, AP000555, AC006443, AC004966, AC002504, Z94802, AC007114, AL035415, AC005921, AC008543, AL022328, AC010129, AC006486, AF196972, AL133353, AL135744, AP001752, AL137798, AL035588, AC087225, AP001732, AP000513, AL137019, AC008736, AC009506, AC008747, AC006329, AC009756, AP000511, AC011500, AL034555, AC005037, Z99716, AF200465, AL121580, AC007216, AC002319, AL049569, AC003108, AC005666, X53550, AC005682, AC009470, AC008379, AC018641, AJ246003, AF111168, AC012627, AL022318, AL158167, AL161670, AC004041, AL109627, AB001523, AC020898, AL162740, AL136179, AP001760, AC006511, AC008040, AL021155, AL161937, AC007546, AF077058, AC006153, AC004583, AL139396, AL035587, AC008519, AL031276, AP000553, AC025435, AL137061, AL365444, AC005228, Z97180, AC008739, AC006030, AL033529, AC007249, AC006111, AC004904, AC002407, AC011811, AC003070, AC011526, AC012076, AC005231, AC068799, AL031133, AC004089, AL157938, AC007690, AC006345, AC036103, AC005529, AC004859, AL096791, AL050321, AP002534, AC006449, AP000692, AC005920, AC084693, AC007277, AL035422, AC004386, AL022316, AC005995, AC006344, AC004685, AL034553, AC004382, AF168787, AC024952, AL034379, AC000052, AC004837, AP001753, AP000501, AL133246, AP000133, AP000211, U57005,
--	--	--	--	--



				<p> AW753067, AW965175, AW959202,  AW973482, AW958993, AW959136, D51060,  D80269, D80024, D50979, AW965158,  AW949632, T03269, AW965196, D80378,  AW960454, AW960473, AW949654,  AW958992, AV699682, D80241, AW959062,  AW964477, AW956434, AW964488,  AW962082, AV701004, AW964532,  AW965176, D50995, C14014, AW966043,  C75259, AW956397, AW966030, AW966032,  AW960532, AW177501, AW949633,  AW177511, AW753053, AW966023,  AV718530, AV742048, C14407, AV742001,  AV700229, AV738928, AV699669,  AW960564, AV701335, D81026, AV742667,  D51022, AW975623, AV701043, AV701332,  AV719049, AV701017, AV701248,  AV701431, AV701149, AW752082,  AW959469, AV719628, AW960504,  AV720150, Z21582, AV645389, AV645344,  AV720533, AV720654, D80134, AW960570,  AW378532, AW178893, AV701130,  AV701419, AW352117, D58253, AV719913,  AV701125, AV701166, AA305578,  AW960414, AV744690, AV681510,  AV681491, AW973465, AV699866,  AW179328, D80949, AV699652, D80248,  AV719000, AW960474, AW973490,  AW973445, AW966368, AV703738,  AV645343, AW973470, D80133, AV745080,  AV701422, AV742732, D51250, AV701154,  AV701428, AW960514, AV701443,  AX047063, A62298, A84916, A62300,  Y17188, AX033851, AR018138, AX047064,  AX047062, AR070327, AX027925, AJ132110,  AJ302649, A82595, AX021518, X67155,  AF058696, AX020190, D26022, D34614,  AX035434, AX020191, A67220, AR008278,  D89785, A25909, A78862, AB028859,  AR087649, AR092424, AX028130, X82626,  AR077702, D88547, Y12724, Y17187,  AR025207, AJ294956, X68127, A94995,  AR060385, AB002449, AF260572, A30438,  AR008443, AR074545, AB012117,  AX015396, I50126, I50132, I50128, I50133,  AR008277, AR008281, A85396, AR088705,  AR066488, AR016514, AX042372, A85477,  AR060138, AJ287395, A86792, A45456,  A26615, AR052274, A44171, AR074139,  A43192, A43190, AR038669, AR074141,  AR066482, I14842, Y09669, I19525, U79457,  AR066487, AR074136, X93549, AR066490,  I18367, AR054175, AX014811, D50010,  AR016691, AR016690, U46128, D88507,  A63261, AR008408, AF135125, AR062872,  AR091537, A70867, AR093385, A64136,  A68321, D13509, AR060133, I79511,  AX035426, AX035429, AX035428,  AR032065, AF123263, AJ000347, X72378,  and AR008382. </p>
--	--	--	--	---

HCE1G78	76	761204	1 - 1882	15 - 1896	AW025289, AI935720, BG222525, AA724676, BF844613, BE707252, AW385203, AW580449, AW243018, BE932090, R15390, BF436472, BF351100, AW014134, AA074234, R18788, BF819553, BE162530, H14886, BF087139, AA772066, F35935, R42130, R40003, AI628487, BE169397, R13943, AI540418, BE167881, BF800299, BE142196, AI804744, BE931587, BE166493, AW890237, AL036574, AI675744, R88613, AW607153, BF082899, AW935303, U45975, AB032551, and AC005005.
HCE2H52	77	847007	1 - 1262	15 - 1276	AI821394, F08688, AW812688, AA826667, BE089841, H54483, AV714369, AW812690, U49973, AC016395, AC010386, AC004828, AL160237, AL137787, AL353643, AC002456, AL049838, AC008598, AC004941, AL136168, AC018719, AL136419, AC006211, AL121944, AC005544, AP001818, AL136146, AL158093, AL121865, AC008806, Z99497, AC019106, AC011299, AL121894, AL133257, AB020862, AC006213, AL359382, AC018360, AC004241, AL021408, AC002556, AC005872, AC083862, AC004067, and AL161450.
HCE3B04	78	831151	1 - 1793	15 - 1807	BF341078, BG028747, BE879916, BF059108, AW300205, AI634862, BE048884, BF195876, AI636211, AW117753, BF573148, BF197549, AW954937, N91173, AW168897, AA983273, AI374834, AW002887, AI435122, AI674869, BE673355, BE671667, AW081459, AW271351, AW237603, AI818463, AI025174, BF691042, AW952156, BF832239, BF089340, AI559577, AA758512, N48695, BF790935, BF211459, AI492924, AW168956, AA291263, BF738867, AI476602, AA209287, AI953330, AI702174, AI590318, N29813, AA653205, BE645737, AA908587, W19735, AI679742, AA255954, BF666339, N49753, AW087559, BF571795, M86083, AI303020, AA148623, N89992, T31216, T16818, N72208, AA642349, N45545, AL044337, AW515018, W19616, AA256117, AI276869, N52681, T86722, D61438, N59844, AW391658, N51450, AA319376, W31671, BF839545, AI702072, AI623267, BF381609, AI692792, AI014575, AW151467, AW389355, BF751083, D57869, N22895, BF752070, BF748108, AW449444, N55976, N90029, BE565819, BF908853, AW601227, BE074229, W17143, BF751076, BF751211, BE869711, BF154561, AF020762, and AK025520.
HCE5F78	79	838101	1 - 1718	15 - 1732	AK025051.
HCEDR26	80	771144	1 - 1405	15 - 1419	AW809560, BF822291, AW805745, T06675, T41328, AW809450, BF884442, BF773357, BF738231, BE163588, BF998055, H00095, BF900030, AA346118, AA644090, BF725844, BF725688, AI919265, AI801505, AW103406, AW855803, BF673854, AA833896,

				AW958962, AA630854, AI798521, AW855730, AV702109, AA833875, BF725884, AA513851, AW814024, AI537020, AW474825, AW243793, AW275432, AW970064, AC010326, AC010645, AC010522, AL353748, AC003006, AL136418, AP000501, AP001760, AC004840, AC018738, AC008264, AC018641, AC005102, AC007383, AC020908, AC011497, AP000338, AC004526, AL137072, AP000216, AC006011, AL024498, U62293, AC004913, AC002044, AC022148, AL022316, AL354864, AL035400, AC005529, AL109797, AC004975, AC004895, AC002472, AC005881, AF258547, AC002350, AL049776, AP002028, AC002558, U91321, AC008745, AC020913, AC004859, AF243527, AL031311, L44140, AL138741, AC003048, AL137012, AL117352, AC005081, AL109801, AL109798, AL163279, AL035587, AC010422, AC010789, AF045555, AC011480, AC004491, AC011470, AC007283, AP001711, AC005409, AC008812, AC008569, AF168787, AC005082, AL118505, AL135927, AC007227, AC020917, Z98949, Y14768, AL117382, AC004156, AP001694, AC007051, Z82215, AC004796, AC007664, AC020916, AP000505, AC010328, AC004804, AJ400877, AF111167, AC004167, AC007536, AC007216, AP001052, AL139099, AC004755, AL034420, AC000052, AC003043, AL121891, AC002314, AC005920, AL159977, AD001527, AL021808, AC005702, AL355476, AC011455, AC009244, Z85986, Z83844, AC004150, AL136980, AL158823, AL158040, AC000353, AC005899, AC011489, AL035086, AC005971, AL136087, AL445483, AL133387, AP001752, AC007993, AL353804, AL031767, AC009314, AC006252, AC007666, AL121886, AC005800, AC004815, AL136179, AC008536, AC008482, AL109921, AC011442, AC006333, AL158830, AC003029, AC012099, AL138733, AC005412, AC027319, AL133294, AC007919, AC005291, AC005011, AC005098, AL356244, AL137230, AF235097, Z93015, AP001718, AC004019, AP001720, AP001725, Z82901, U52111, AC002492, AC004253, Z82190, AL008718, AL031657, AL133448, AC007225, AL353653, AC007390, AC008543, AC004738, AC006211, AC007739, AL133545, AF111169, AL121601, AC008474, AC004882, AC002301, AC008072, AF196779, AC005372, AC008806, AF207550, AC009600, AC007404, AC012384, AC008132, AC004089, AL050335, AF053356, AF038458, AC005057, AL163303, AC004813,
--	--	--	--	--

					AC004797, AC005484, AL133229, AL356379, AP001724, AL121926, AC008372, AC006483, AF064861, AL035071, AC020663, AC006329, AC022073, U95742, AC005231, AC005399, AC004000, AC007620, AC007066, AC006111, AL356575, AC011462, AL022476, AL121652, AF030876, AL138752, AC005736, AC005089, AC008551, AC006511, AP001670, AC005225, AF111168, AL132713, AC002352, AP000688, AJ009616, AL359457, and AC011475.
HCEEE79	81	560609	1 - 1038	15 - 1052	
HCEEQ25	82	531784	1 - 978	15 - 992	AW444547, BF514399, AL534267, AI567447, BE747694, BG152517, AW298411, AW865264, AA807579, AA554958, BE889430, AA612578, BF798462, AI078409, AU157259, AI819391, AA643770, AU120121, H77386, AW438907, A87682, U78181, U78180, A98491, U94403, A87681, A87684, AC003687, AC002094, AC007220, AL136984, AC020750, AL031666, AL136110, AL161781, AC020744, AL031672, AL162424, AC002425, AL022721, Z75887, Y10196, AL139340, AC021863, AC025464, AL353812, AC007956, AL117382, AC005789, AL049555, AL035086, AC008403, AP001754, AL035684, AC007204, AL021395, AC020552, AL157701, AC010605, AL137139, AC083866, AL035695, AC007021, AC010412, AL021707, AL135927, AC007227, AF109907, AC008651, AL035587, AL133259, AL136418, AC010200, AL353194, AL034405, AL020997, AL049539, U91327, AC008551, AC004816, AF123462, Z95330, AL121914, AC002120, Z97876, AC004846, and AC005189.
HCEEU18	83	688041	1 - 1215	15 - 1229	AL045384, AL042668, AI525108, T85422, AL046089, BE843928, H08562, AA921935, AA815292, AW972431, F23282, BE794230, U91320, AC008469, AB018295, AL117630, AC009032, AC003043, AC008745, AC007405, AC004867, AL117381, AC004967, AC013429, AC005098, AC016395, AL050335, AC005088, AC020913, AF001548, AC004876, U91321, AC005279, AL355392, AC011497, AL031658, AC005412, AC005231, AC005089, AC009600, AC011490, AL109897, Z98051, AC011495, AC009087, AL133215, AC010271, AL136305, AC004125, AC007052, AC004815, AC005944, AC004703, AC004019, AC004813, AC011479, AF168787, AC012384, AC004797, AC005052, AC007282, AL080243, AL031680, Z84469, AC006116, Z84466, AP001717, AC007956, AC006014, AL034549, AF030453, AC006454, AL049761, AC005821, AL133551, AL157938, AL009181, AL133353, AL031670, AC005488,



					AC000025, AF205588, AC012076, AL355094, AC011455, AC020934, AL157372, AC087094, AC083863, AC005527, AC022515, AC004966, AC009086, AC005056, AL049795, AC004841, AL121825, AC002425, AL121900, AP000687, U80017, AC022392, AL132713, AC008403, AC018644, AC036103, AC020663, AC018719, AC027319, AL118505, AC002039, AC011484, AC008040, AC007240, AP001670, AL031685, AC002565, AL031230, AC021999, AC011444, AF111168, AL391114, AC005874, AF134471, AC008392, AC002287, AC004491, AC005722, AC025593, AC004166, U63721, AC011491, AC020550, AC007383, AC011465, AL356652, AC005792, AC004854, AC009123, and AC006388.
HCEFZ82	84	831745	1 - 1797	15 - 1811	BF981465, BF688419, BF969763, BG178653, BE730527, AI672493, N21040, BE395792, AW386160, BE858812, AI672483, BF530193, AI693512, AV751914, BG180158, AI138621, BG104179, AA778387, AA173791, BF939691, AW615384, AW960851, AW594109, BF091657, AI022755, AA209239, AI077708, AI824069, AI936432, AI038303, N39250, AI927782, AI457926, AI436138, AI056772, AI079503, N58793, AI016045, AA210850, AI096581, AA062719, W88815, AA725072, AI375410, AA669791, BE300887, BF431891, AA173843, W31742, W88816, AI740977, BE727603, AI086937, AA704681, AI190844, AI341909, AI365029, N46695, BF590052, AV749863, AI086941, AI676179, AA826493, AA554932, AA789007, BF111593, AA917998, R08679, AA889734, W04647, AA321894, AI912831, AV750240, AI239655, BF592139, H71960, AI368377, AA992261, BE277655, H78240, H78440, AI470391, R37067, AV694383, AI700804, R44781, AW612991, R10835, H96434, N77482, AA314780, R44068, AV751269, R08587, AV697548, AI419628, BE218690, N90646, H65409, BF530646, AA836620, W26811, R10834, AV660888, AV747670, AA905784, AI086303, H84253, AI086248, AV723953, BE881061, BG110517, BE047952, BG180996, AV682466, BF107905, AI312428, BE876038, AW051059, AI538885, AV757598, BF752170, BG113385, BF968903, BG028873, BG113847, AW301865, AL036802, BG033199, AV682875, BG178911, AI345612, AV732936, BF924882, AW827285, BE966634, BG120492, AI345415, AW827206, BG164371, AW827214, AW827276, BF971336, AA568405, BG026714, AL118781, BE965758, BE965192, BE875407, AI581033, AL041573, BG260037, BF835240, AV682330, AI343059, AV760102, BG058150, AI361701, BF816811, AV756838,

095008-0101

				AI345416, BF338002, AA614183, AI349933, AI340519, AI349937, AI340603, AW129264, AV681885, BE887488, BF341801, AV757639, BE881134, AW022636, AV682787, AV682672, AL036631, BE907151, AL040169, BG034564, BF108279, BG110797, BG029053, AV709517, AA572758, BE966699, AV757018, BE963838, BE875868, BE905726, AW403717, BG114990, BG110660, BG116387, BG036846, BF980991, AI859991, BG119329, BG260187, AV729627, BF915208, AI241901, AW161579, BF911517, AI815232, BE895585, AL036396, BF792050, AI064830, AW673679, BG111377, AV734765, BF814450, AW935969, AW806761, AL045413, AL119836, AI698391, BE536058, BF968533, BE964614, AW302988, AW827289, BF764538, AW778801, AL039086, AV706987, AI433157, BE910005, BE048135, AL513943, AV757943, BE892325, BE965621, BE967307, AI349645, BE874133, BF791952, BG033267, BE965481, BG105812, BF527014, BF816685, AW834302, AI683395, N71199, AV733682, AI312152, BF909758, AF225418, AL133067, AF130066, I48978, AL049300, AF116631, AL110196, AL122050, AL096744, AF090934, AF100931, AL137529, AK025958, U42766, AL133565, AF218031, AK026784, A08916, AL117457, AK026434, U35846, AF119875, AL389935, I48979, AL162006, S61953, I89947, AF116644, AF113694, AF097996, AF218014, AF090900, AB047631, AB047941, AL050393, S78214, AL357195, AL133557, A08910, AF207829, AF130059, AX019230, AL133080, AL133640, AF210052, AL359601, AK026518, AL137459, AK025015, AL035458, AF078844, AF116691, I89931, AF130075, AL050116, AB052200, AL442072, AX019229, AF113690, A08909, AK026353, U87620, AF130068, AF111847, AL133606, A03736, AK026542, AK024992, S36676, AL049314, AL050146, AL442082, X72387, AK026506, AL137527, Y07905, AF113013, AB049892, AF022813, U95114, AF118070, AL049382, AK027161, AL359583, AK026741, AB048974, A08913, AK026927, AL157431, AL162085, AF090896, AL122121, AL162002, AF116639, AX046603, AK026583, AL049430, AK026797, AF017152, AX020124, AL133016, AK000323, AF158248, AL122118, X93495, AL137283, Y16645, AF090943, A65341, AK026480, AK026522, AK026630, AK025708, AK025484, AF175983, AR087170, AF118064, AK026086, AL133093, I26207, Y11587, AK000432, AK025339, AF130087, AF146568, AL137705, AL122123, AK000618, AF106827, AK026855, AX017991, AB052191,
--	--	--	--	--



					BE879597, AA643500, AI090381, BE876617, AW055002, AI744096, AA714840, AI148138, AA598703, AW340721, BE559510, AI125728, AI830384, AU151986, AI885716, BF432640, AI475597, AW339888, AI377299, AI309247, AW005497, AW771450, BE622365, AI139390, AI422379, N95665, AI144110, BF341713, AI919264, AI475648, AI189926, BF809564, AI559248, BE396630, AI817016, AI640708, AA825735, AV662025, BF526593, AI708507, AA935563, AW939178, AW939138, AW939190, AA961207, N66071, N69365, AA603786, AW939128, AI285062, AI826293, AW951255, BF512843, AA620317, AW605529, N80991, AI285338, AW381729, BF956527, AI031873, AW381749, BG056758, AW371517, AW381756, R78288, AI434319, AI282698, AI343784, AW002119, AW169595, N48713, AW371499, AI537993, N26561, R52366, AI916353, R93746, BE828699, AA835499, AW072300, W70306, AA907306, AW748173, N68294, AI468129, BG032430, H89920, AI080137, AA326474, H69097, AA826574, AW303330, AI858652, AI804227, AA322294, AI342977, AA291513, N35677, AI824463, T86711, H49562, Z19236, AI500205, AW966976, AI187820, R78289, AL525884, W76006, BE828722, N81088, F37932, R31642, AA905077, AA922535, AA877249, BF957167, AA780926, BG055657, AW087260, AI202070, AW129497, AA911554, AA045568, R32359, AA204681, AA057054, AA364763, R63145, AI277392, AI886719, BE616036, AW238934, AI082383, BG164867, BG150873, BE548567, BG035187, N91393, BF957863, H69096, R88359, AW328569, BF743483, BF742356, AC005089, X89985, and AJ223979.
HCFLT90	87	788578	1 - 896	15 - 910	BG164543, BG254195, BE569122, BF701728, BE883133, BG164607, BE564055, BE621682, BE621082, BE548066, BG164527, BE439646, BF212991, BF669801, BF679424, BF699188, AW958367, BF696977, AI300093, BF104175, BE896610, BF382702, N73920, AI301065, BE327267, AV714574, BF382383, AA131247, AA417682, AA053860, BF681596, BF242641, BE909206, BE327067, W80481, BF666642, AW263496, BF696681, AI479737, AI499701, W56158, AA922117, AA781414, AA131515, AW104450, AA001425, AV754612, AW630854, AW274584, T66256, AW173171, BG252902, AA258280, AA026671, BF679441, AA256918, AW071970, AW817587, AA019198, BF695475, AA054011, AA026632, AA954721, AI926512, AI393574, AA742324, AA972701, AI685317, W47268, BE856240, AA434451, AW630034, AV713327, AI305153, BF673772, AA678370, AW613806, AI356800, AI280399, AI200095, AI749979, AI334386, BE262340, AA837589,

					AA164369, AW627592, N30835, AI491703, AI073728, AI422897, N48017, AV744241, AI218584, AA885442, AI367984, H16890, BF029129, AI282124, AA054252, W47269, N36665, AA053948, BE439592, AA731678, AI273274, AW166832, R94765, AW338258, BG027442, BE710524, AA256494, AV764328, AI086693, BF381741, AW022998, AI347082, R21151, AA743354, AI200245, H84116, AI280442, R73589, AI698561, BF029639, F12140, AA054003, R72975, AA480586, H03499, W78713, AW182912, BF208960, D79117, AA026691, AA713938, BE958068, R69294, AI381988, BF765409, R69295, BF765699, AI948431, H03608, R34494, AA379320, BF091606, AA019166, AA026702, H84115, AA095757, BE708262, BF197074, AA365458, C01000, AA044911, AA417566, and AA877878.
HCHAB84	88	834326	1 - 1345	15 - 1359	X84712, BF526942, BF036429, BF035689, BF034330, BE878646, BG117306, BE906856, BE871642, AV703538, AW955111, BE729985, BE958344, BE271782, BF698225, BE568321, BG250080, BF826293, BE156569, AW579884, BF977502, AA587630, BE621946, BF512422, BF695706, AV764128, AA448786, AI032411, AA477231, BF695587, AA641139, BE621432, H02682, H02590, H29948, AW972521, AA186733, BF909586, R22544, BE673152, AW405966, AI358557, R22543, BG169787, BF813006, BF742389, AW673871, AA327923, AW392393, AA311766, BF916201, BF742334, R70413, AV763358, AA505606, AA477230, AW794624, AW674083, AA595661, AA579044, AW265468, AA807704, AA642809, AW021674, AA618263, AA405570, BG180320, AA533066, AI702049, AI061313, AI254267, AA084439, AA491767, BG059139, AL042667, AL042670, AA187682, AV763460, BF131490, AA313025, AL121039, AA557945, AW148821, BF901147, AW402784, AA693484, AV758870, AW410844, AW873417, AA601376, AL119909, AI251024, AI444575, BF725761, AW963489, BF857486, AI141202, AA776665, AW069110, AA601290, AW469462, AW270385, AI572680, AW028376, BF817511, AA809116, BF739035, AI064968, AA640310, AA535216, AI679759, AI753113, BF030482, AA600863, AI185160, AW192930, AL138262, BG118544, AW675677, AW023390, AW672927, R67038, AI445699, AI312267, AA659832, AV647070, BF804385, AA610644, AI821901, AV764383, AW105463, BF770715, R83577, AU157209, AI039257, AA602906, AA809546, AW820105, AA502991, AV762112, AI252611, AI567676, AI476049, AA015948,

06505660 "6070"

				<p> AF126023, AF126024, X96618, U02057,  AC002492, AP001760, Z68276, AC008649,  AC007842, AP001711, AC021752,  AC007405, AP000133, AP000211, AC004922,  AL034451, AL139039, AF168787, AC010463,  AC003102, AL139353, AC006261,  AC009756, AP000563, AC002310,  AC008068, U47924, Z69719, AL050349,  AC003048, AL391839, AL031846, AL121655,  Z95331, AL389875, AC005207, AP000252,  AC006111, AP001692, AC007333,  AC005821, AL035684, AC005081,  AC008745, AL136300, AC005902,  AL034418, AJ295844, AC020904, AF139813,  AC005519, AL445483, AC020906, L78810,  AC008567, Z83844, AC008821, AC073898,  AC009311, AC004542, AC002306,  AL031311, AC016683, AC023105,  AC006449, AL139100, AC016594,  AC008044, AL133517, AC002554,  AL020993, AC008403, AC005355,  AL121675, AL359695, AC005261, AP000511,  AL096712, AC024561, AL117348,  AC010627, AC005288, AC010271,  AP001745, AL135783, AC007055, L44140,  AL031427, AC007637, U85195, AL035086,  AC012085, Z84466, AF053356, AC008009,  AC009247, AL133211, AL158830,  AC002551, AL109984, AC002477,  AL159168, AF205588, AC007383, AP000146,  AC010491, AC006483, AE000658,  AC004883, AC025264, AL133230,  AC007390, AF042090, AL035455, AC005280,  AC008897, AC011495, AC005038,  AC005914, AL136228, AP001709, Z97181,  AL022323, Z98048, AL133453, AF111168,  AC010422, AP001630, AC010458, AL109935,  Z94160, AL157372, AC005080, AC005412,  AC009123, AP000356, AL137787, AC004854,  AL121886, AC003684, AC027319,  AC005759, AC004263, AC020552,  AC004686, AL049636, AC007957, AJ243213,  AL139343, AL117381, AC005527,  AC008521, AC011891, AL138836,  AC005529, AC004975, AL442166,  AB023048, AC008555, AC011445,  AC005781, AL121712, AC020934,  AC008372, AL031587, AL121601, AL133355,  Z84488, AL133451, AL133448, AL163284,  AC008543, AL096701, AC010203,  AL157818, AC005180, AC010620, AF038458,  AC004659, AC007277, AC000353,  AP000557, AL031575, AC078844, AC006581,  U91327, AC005619, AC004491, AC002401,  AC007216, AC009298, AC004000,  AL049758, AL163282, AL020997, AC005585,  AL022727, AP001748, AC020750, AC004076,  AL121933, AL033520, AC007919, AF064861,  AL031594, AL158172, AC012627,  AC007237, AC011494, AC002375, Z82182, </p>
--	--	--	--	---

					AL121926, AC010328, AC006241, AC002394, AL109797, AC011469, AL136418, Z83838, AC007666, AL049780, AC005755, AC006571, AL133288, AC011480, AL158040, AC007358, AL390374, AC009086, AC008083, AC017111, AC005365, AC006597, AL022315, AC006349, AL096775, AC004971, AF109907, AC005911, U91323, AL021877, AC005602, AC004846, AC022150, AL033521, AC084693, AC007179, and AC006512.
HCMXS51	89	788643	1 - 2239	15 - 2253	AL520206, AL522291, AL520207, BG115714, BG023953, BF343959, AU133571, BE839880, AW954438, BE264316, BG261277, BE879757, AU131026, BE265959, BE278903, BF725639, AW246741, AA864833, AU148856, BF111640, AA706935, AA431813, N38742, BE857705, BF476344, AU152863, AU125122, AA480041, AW170367, AI094797, AU154528, N48379, BF913004, N26479, AI803158, AL120744, N35219, AW245159, AI089912, AI927351, N20323, AL046695, AA476664, N35530, AI078494, AA015687, AI016568, BE857202, AI587317, AA446620, AW629254, AI433184, AA548282, W03412, N27597, W00855, AA825427, AI128747, AI082265, H38927, BF970202, R48359, AI569253, N29410, N44883, AW014479, AA934555, N41471, AW974179, R15948, AA573084, AA233832, AA017058, W16680, AI312737, R60804, N67483, R15949, Z43237, BE811896, N45053, BE832888, T11764, Z43099, AA431409, R48260, AI933045, AW874096, AW105691, AA336676, AA044969, BF362640, AW893387, AW892550, AW892516, R60299, N35043, T49574, Z41630, AA013473, N35211, AA738419, AA223632, H86402, T49573, AA017209, BF941569, AA635071, AA054652, H86066, AI351292, F02575, N79527, T11765, T35773, AW993110, AW194575, BE893541, AA448030, AI086309, BF737533, AF001690, AF029231, U96629, AB007042, AB011091, T66574, and T66575.
HCNCO11	90	775086	1 - 732	15 - 746	BF926420, BF926408, BF875996, AV705104, AV726755, AW964429, AW950395, AV703435, AV707451, AV707628, AW961373, AV705453, AW964210, AW964423, AV704361, AW952896, AW961510, AV726887, AV729165, AW963643, AV707705, AW963965, AV707556, AV702814, AW963219, AV704916, AV706906, AV703045, AW950229, AV690921, AV704674, AV728297, AV702810, AW960535, AI557262, AW963644, AV708024, AV701594, AV727806, AV727803, AW957298, AV650843, AW957682,

095005660

				AV704283, AV708829, AV701751, AW950078, AW950079, AW949946, AW961329, AW954386, AW954962, AW957974, AW952228, AV725024, AW960663, AW957083, AW950256, AV649266, AV704144, AV703160, AW963857, AW966775, AW958568, AW964298, AW958569, AW966684, AW951998, AV704342, AV703361, AV704848, AV703833, AV703425, AV705771, AV653846, AV728884, AW960406, AW954104, AW945153, AV650865, AV705189, AW958320, AW958316, AV729457, AV728929, AV729285, AV726743, AW949454, AW953619, AW955397, AW949863, AV701787, AW945196, AV708035, AV702901, AW961052, AV701953, AW963108, AV729170, AW958127, AV703284, AV706964, AV728355, AW963612, D50992, AV703367, AV706742, AV703862, AV706047, AV709139, AV652860, AW955394, AV702749, AV708590, AV707020, AW951793, AW951816, AW967188, AV705481, AV706133, AV702435, AV702958, AW955616, AW954194, AW962970, AV726728, AV705420, AV726770, AV695489, AV691061, AW963234, AW955139, AV649942, AW962367, AW964203, AW954242, AW954413, AV727101, AV707189, AW966146, AW951740, AW960600, AW952132, AV703632, AW961431, AW964278, AW958290, AW959722, AV659467, AV706802, AV705998, AV709332, AV705516, AV728913, AV728341, AV709281, AW958045, AW950597, AV707266, AW950411, AW960545, AV705267, AV704712, AV704401, AV702819, AV702458, AV702187, AW965827, AV727386, AW949731, AV707117, AV702298, AV701626, AV727268, AV703465, AW954783, AW953992, AW963581, AW958104, AW950254, AV705154, AW962980, AW957286, AW962378, AW958093, AW963811, AW954221, AV727618, AW963652, AV652536, AW962929, AV704033, AW950520, AW952361, AV701614, AV705518, AV728428, AV729392, AV702035, AV709623, AV702315, AV703266, AV707197, AW945183, AV727756, AV707238, AV706910, AW961377, AW945164, AW950681, AV647033, AV647066, AV647129, AW957628, AW952419, AV707649, AW957779, AV647144, AV702975, D59751, AV650924, AV693604, AW960143, AW963486, AV704876,
--	--	--	--	--





					AI535676, AW579265, AI955386, AA279581, AA983814, BF185409, BE622718, N59886, AI859864, AI498376, BG171039, BG116650, W01363, AI699807, AA824487, T86598, AA994605, AW044013, T85108, AA489144, AW768600, AA811658, AW271482, BE895361, AI631722, AW021293, R64514, T77523, AA736753, H44608, AI955411, AV689519, N90263, AL119283, H94626, T77559, AL119309, T86597, AI909117, AW376940, N79005, N77027, AW105078, N62828, AI701272, AI334730, T07505, AW848643, AW243861, AI909110, BG007292, BE162291, BE162293, AA532611, AL022153, AC004804, AC006840, AC008125, and AL355871.
HCQCJ56	94	832157	1 - 1273	15 - 1287	AI674974, AI217307, AA813576, BE302346, AI824976, AA994749, AI244904, AI262935, AA020796, AA234517, BF432061, AA443035, AA463478, AW079079, AA694400, AI005463, AA776532, R00437, and R00438.
HCQCM24	95	845070	1 - 1915	15 - 1929	AI905478, AW963993, AV701494, AA528253, W37762, BF970457, AW771888, AI905529, BE817678, W37779, BE817676, BE817671, AI245339, AI650320, BG164304, AA524462, AW241742, BE539666, AA532816, AI261199, AW024681, BE817677, BE540718, AI073889, AA634509, AA430047, BG105129, AI301532, BE817544, BE567771, AI284328, W37840, AA946995, AI261595, AA430256, AI557710, AI367914, AW802496, AW470760, AI254217, BF856116, AW183037, BF673560, BE770611, BF855193, AI886042, BE817655, AW802501, AA587825, W31581, AI632720, AI952748, AI204507, AW451863, AI659684, R21943, BE176747, BE003678, BE003673, BE837515, AW866502, AA854919, BG169910, AI608737, BE003667, BE003665, AA902893, BE817476, BE817652, BE176749, BE817470, AW176042, AW866504, BE003676, AA112330, BE817478, BE536935, AA284617, H92723, BF445303, R28123, BF222392, BE003682, AA626191, BF475670, BE817645, AI797583, AA377172, AA579168, T90137, AW363872, AW070545, AI796563, AA069314, AW960665, H57105, AA614369, R28013, BE081148, AI922451, AI824371, BF818434, BF818436, BE837483, BE817451, BF229149, AA314230, BF846051, AW994489, AI972085, AA809584, AA594344, AW299894, BE769260, BE827354, AW299537, AW148344, BE067485, AW148392, BE817444, BE086250, BE855526, AW802489, AW603877, BE817576, BE772109, AW603875, AA345280, BF819869, AA886302, BE817574, BG008318, AI694024, BE837418, BF677528, BF435329, BE928914,